

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2003 (03.01.2003)

PCT

(10) International Publication Number
WO 03/000235 A1

(51) International Patent Classification⁷: A61K 9/14, 9/16

(21) International Application Number: PCT/IB02/01783

(22) International Filing Date: 10 May 2002 (10.05.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/300,255 22 June 2001 (22.06.2001) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/300,255 (CON)
Filed on 22 June 2001 (22.06.2001)

(71) Applicant (for all designated States except US): **PFIZER PRODUCTS INC.** [US/US]; Eastern Point Road, Groton, CT 06340 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FRIESEN, Dwayne, Thomas** [US/US]; 60779 Currant Way, Bend, OR 97702 (US). **GUMKOWSKI, Michael, Jon** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **KETNER, Rodney, James** [US/US]; 1615 N.W. Trenton Avenue, Bend, OR 97701 (US). **LORENZ, Douglas, Alan** [US/US]; 61332 S.E. King Jehu Way, Bend, OR 97702 (US). **NIGHTINGALE, James, Alan, Schriver** [US/US]; 62900 Santa Cruz, Bend,

OR 97701 (US). **SHANKER, Ravi, Mysore** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **WEST, James, Blair** [US/US]; 1511 N.W. Jacksonville Avenue, Bend, OR 97701 (US).

(74) Agents: **LUMB, J., Trevor** et al.; Pfizer Inc., 201 Tabor Road, Morris Plains, NJ 07950 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/000235 A1

(54) Title: PHARMACEUTICAL COMPOSITIONS OF DISPERSIONS OF DRUGS AND NEUTRAL POLYMERS

(57) Abstract: In one aspect, pharmaceutical compositions comprising dispersions of an acid-sensitive drug and a neutral dispersion polymer are disclosed. The acid-sensitive drug has improved chemical stability relative to dispersions of the drug and acidic polymers. In another aspect, pharmaceutical compositions of low solubility drugs and amphiphilic, hydroxy-functional vinyl copolymers are disclosed.

PHARMACEUTICAL COMPOSITIONS OF DISPERSIONS OF DRUGS AND NEUTRAL POLYMERS

5 This application claims the benefit of priority of provisional Patent Application Serial No. 60/300,255 filed June 22, 2001, which is incorporated herein by reference in its entirety for all purposes.

BACKGROUND OF THE INVENTION

10 This invention relates to pharmaceutical compositions comprised of amorphous dispersions of drugs and neutral polymers that provide either improved chemical stability, concentration-enhancement, or both improved chemical stability and concentration-enhancement.

15 It is sometimes desired to form a solid amorphous dispersion of a drug and a polymer. One reason for forming dispersions is that the aqueous concentration of a poorly soluble drug may be improved by forming an amorphous dispersion of the drug and a polymer. For example, Curatolo, et al., EP 0 901 786 A2 disclose forming pharmaceutical spray dried dispersions of sparingly soluble drugs and the polymer hydroxypropyl methyl cellulose acetate succinate. The spray dried dispersions disclosed in Curatolo et al. provide 20 superior aqueous concentration relative to dispersions formed from other methods and relative to the crystalline drug alone.

Similarly, others have recognized the enhancement in aqueous concentration afforded by dispersing a drug in a polymer. Nakamichi, et al., U.S. Patent No. 5,456,923 disclose solid dispersions formed by twin-screw 25 extrusion of low solubility drugs and various polymers.

Another reason for forming an amorphous dispersion is that it may be desired to use a particular process for forming a pharmaceutical composition, such as a spray-coating or wet granulation process which results in the formation of amorphous drug, in whole or in part, rather than pure crystalline 30 drug. Thus, amorphous dispersions may be formed of drugs which are not low-solubility drugs.

However, regardless of whether the drug is poorly soluble, the inventors have determined that for some drug and polymer dispersions, the drug is not chemically stable in the dispersion. In particular, the inventors have observed 35 that for dispersions containing certain drugs and polymers, the drug degrades in

the dispersion over time, resulting in a loss of potency for the composition. The inventors have found this problem to arise especially for acid-sensitive drugs.

Drug degradation within the dispersion is a particular problem for low-solubility, acid-sensitive drugs, since the increase in aqueous concentration of the drug provided by the dispersion is offset by decreasing drug purity. In general, the greatest concentration-enhancement is often observed through the use of acidic dispersion polymers, especially acidic, cellulosic enteric polymers. However, the use of such acidic polymers within the dispersion is precluded due to the acid-sensitive nature of the drug.

Accordingly, there is still a need for pharmaceutical compositions of dispersions containing acid-sensitive drugs that are chemically stable over time. Likewise, there is also a continuing need to provide concentration-enhancement for low-solubility drugs.

BRIEF SUMMARY OF INVENTION

The present invention relates to, in a first aspect, pharmaceutical compositions comprising a solid amorphous dispersion of an acid-sensitive drug and a neutral dispersion polymer, wherein said composition provides improved chemical stability relative to a control acidic dispersion comprising an equivalent quantity of said drug and an acidic polymer.

In a preferred embodiment, the acid-sensitive drug has one or more functional groups selected from the group consisting of sulfonyl ureas, hydroxamic acids, hydroxy amides, carbamates, acetals, hydroxy ureas, esters, and amides.

In another preferred embodiment, the acid-sensitive drug when present in a control acidic dispersion and stored for a period of six months at 40°C and 75% relative humidity has a degree of degradation of at least 0.01%, preferably at least 0.1%.

In another preferred embodiment, the drug is quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3- fluorobenzyl-2(S),7-dihydroxy-7-methyl-octyl]amide; quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide; quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide; (+)-N-{3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl}-N-

hydroxyurea; omeprazole; etoposide; famotidine; erythromycin; quinapril; lansoprazole; or progabide.

In another preferred embodiment, the drug in said composition has a relative degree of improvement in chemical stability of at least 1.25, preferably at least 3, more preferably at least 10. Preferably, drug in said composition has a relative degree of improvement in chemical stability of at least 1.25 when stored at 40°C and 75% relative humidity for a period of six months.

In another preferred embodiment, the dispersion polymer is ionizable, non-ionizable, cellulosic, or non-cellulosic. Preferred cellulosic dispersion polymers include hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and hydroxyethyl ethyl cellulose. Preferred non-cellulosic dispersion polymers include vinyl polymers and copolymers having one or more substituents comprising hydroxyl-containing repeat units, alkylacyloxy-containing repeat units, or cyclicamido-containing repeat units; polyvinyl alcohols that have at least a portion of their repeat units in the unhydrolyzed form; polyvinyl alcohol polyvinyl acetate copolymers; polyethylene glycol, polyethylene glycol polypropylene glycol copolymers, polyvinyl pyrrolidone; polyethylene polyvinyl alcohol copolymers, and polyoxyethylene-polyoxypropylene block copolymers. Preferably, the non-cellulosic dispersion polymer comprising a vinyl copolymer having: (1) hydroxyl-containing repeat units; and (2) hydrophobic repeat units.

In another preferred embodiment, the acid-sensitive drug is also a low-solubility drug and said dispersion polymer is concentration-enhancing. Preferably, the drug has a minimum solubility in aqueous solution in the absence of said dispersion polymer of less than 1 mg/mL at any pH of from about 1 to about 8, more preferably the minimum aqueous solubility is less than 0.01 mg/mL.

In another preferred embodiment, the drug has a dose-to-aqueous-solubility ratio of at least 10 mL.

In yet another preferred embodiment, the dispersion polymer is present in an amount sufficient to provide a maximum concentration of said acid-sensitive drug in a use environment that is at least 1.25-fold, preferably at least 2-fold, that provided by a second control composition comprising an equivalent quantity of said acid-sensitive drug and free from said dispersion polymer.

In another preferred embodiment, the dispersion polymer is present in a sufficient amount so that said dispersion provides in a use environment an area under the concentration versus time curve for any period of at least 90 minutes between the time of introduction into the use environment and about 270 minutes following introduction to the use environment that is at least 1.25-fold, preferably at least 2-fold, that provided by a second control composition comprising an equivalent quantity of said acid-sensitive drug and free from said dispersion polymer.

In another preferred embodiment, the dispersion polymer is present in a sufficient amount so that said dispersion provides a relative bioavailability that is at least 1.25, preferably at least 2, relative to a second control composition comprising an equivalent quantity of said acid-sensitive drug and free from said polymer.

In yet another preferred embodiment, the drug is base-sensitive and said dispersion polymer is non-ionizable.

In another preferred embodiment, the dispersion comprising a buffer. Preferred buffers include sodium acetate, ammonium acetate, sodium carbonate, sodium bicarbonate, disodium hydrogen phosphate and trisodium phosphate. Preferably, the buffer comprises at least 10 wt% of said dispersion.

In another preferred embodiment, the dispersion has a pH from about 6 to about 10. Preferably, the composition comprising a base and the dispersion has a pH of from about 6 to about 10.

In still another preferred embodiment, the acid-sensitive drug is a low-solubility drug and the composition comprising a second polymer, said dispersion is free from at least a portion of said second polymer, and said second polymer is concentration-enhancing. Preferably, the second polymer has at least one hydrophobic portion and at least one hydrophilic portion. More preferably, the second polymer is a cellulosic, ionizable polymer, comprising hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, hydroxyethyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, hydroxyethyl methyl cellulose acetate succinate, hydroxyethyl methyl cellulose acetate phthalate, carboxyethyl cellulose, carboxymethyl cellulose, cellulose acetate phthalate, methyl cellulose acetate phthalate, ethyl cellulose acetate phthalate, hydroxypropyl cellulose

acetate phthalate, hydroxypropyl methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, hydroxypropyl methyl cellulose acetate succinate phthalate, hydroxypropyl methyl cellulose succinate phthalate, cellulose propionate phthalate, hydroxypropyl cellulose butyrate phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate, ethyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate, hydroxypropyl methyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate succinate, cellulose propionate trimellitate, cellulose butyrate trimellitate, cellulose acetate terephthalate, cellulose acetate isophthalate, cellulose acetate pyridinedicarboxylate, salicylic acid cellulose acetate, hydroxypropyl salicylic acid cellulose acetate, ethylbenzoic acid cellulose acetate, hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl phthalic acid cellulose acetate, ethyl nicotinic acid cellulose acetate, and ethyl picolinic acid cellulose acetate. Other more preferable second polymers are non-ionizable cellulosic polymers comprising hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and hydroxyethyl ethyl cellulose. Still other more preferably second polymers are ionizable, non-cellulosic polymers comprising carboxylic acid functionalized polymethacrylates, carboxylic acid functionalized polyacrylates, amine-functionalized polyacrylates, amine-functionalized polymethacrylates, proteins, and carboxylic acid functionalized starches.

Yet other more preferable second polymers are non-ionizable, non-cellulosic polymers comprising vinyl polymers and copolymers having one or more substituents such as hydroxyl-containing repeat units, alkylacyloxy-containing repeat units, or cyclicamido-containing repeat units; polyvinyl alcohols that have at least a portion of their repeat units in the unhydrolyzed form; polyvinyl alcohol polyvinyl acetate copolymers; polyethylene glycol, polyethylene glycol polypropylene glycol copolymers, polyvinyl pyrrolidone polyethylene polyvinyl alcohol copolymers, and polyoxyethylene-polyoxypropylene block copolymers. Within these vinyl copolymers, preferably the second polymer has (1) hydroxyl-containing repeat units; and (2) hydrophobic repeat units.

In another preferred embodiment, the drug in said composition has a relative degree of improvement in chemical stability of at least 1.25, preferably at least 3.

In another preferred embodiment, the dispersion is mixed with said second polymer. Another preferred embodiment, the dispersion and said second polymer occupy separate regions of said composition.

5 In still another preferred embodiment, the second polymer is present in an amount sufficient to provide a maximum concentration of said acid-sensitive drug in a use environment that is at least 1.25-fold, preferably at least 2-fold, that of a control composition comprising an equivalent quantity of said dispersion and free from said second polymer.

10 In another preferred embodiment, the second polymer is present in a sufficient amount so that said composition provides in a use environment an area under the concentration versus time curve for any period of at least 90 minutes between the time of introduction into the use environment and about 270 minutes following introduction to the use environment that is at least 1.25-fold, preferably at least 2-fold, that of a control composition comprising an
15 equivalent quantity of said dispersion and free from said second polymer.

In another preferred embodiment, the second polymer is present in a sufficient amount so that said composition provides a relative bioavailability that is at least 1.25, preferably at least 2, relative to a control composition comprising an equivalent quantity of said dispersion and free from said second
20 polymer.

A second aspect of the present invention relates to methods for treating a condition in an animal comprising by administering to an animal in need of such treatment a therapeutic amount of the above-described composition.

A third aspect of the present invention relates to methods of
25 administering a pharmaceutical composition comprising co-administering to a patient: (a) a solid amorphous dispersion comprising an acid-sensitive drug and a neutral polymer; and (b) a second polymer, wherein said dispersion is substantially free from said second polymer and said second polymer is concentration-enhancing.

30 In one preferred embodiment, the second polymer is cellulosic, such as hydroxypropyl methyl cellulose acetate succinate, cellulose acetate phthalate, hydroxypropyl methyl cellulose phthalate, methyl cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate and cellulose acetate isophthalate.

In another preferred embodiment, the dispersion is administered separately from said second polymer. Preferably, the dispersion and said second polymer are administered at about the same time.

5 A fourth aspect of the present invention relates to pharmaceutical compositions that a solid amorphous dispersion of a low-solubility drug and a neutral dispersion polymer, wherein said neutral dispersion polymer comprising a vinyl copolymer having hydrophilic hydroxyl-containing repeat units and hydrophobic repeat units.

10 In one preferred embodiment, the hydrophobic repeat units include ester-linked alkylate or arylate substituents. More preferably, the hydrophobic repeat unit is an alkylate such as acetate, propionate, and butyrate. More preferably, the hydrophobic repeat units comprise the acetylated form of the hydroxyl-containing repeat units. Preferably, the acetylated form of the hydroxyl-containing repeat units comprise 0.5 to 30% of the repeat units of the polymer.

15 In another preferred embodiment, the hydroxyl-containing repeat unit is vinyl alcohol.

In another preferred embodiment, the dispersion polymer is a vinyl alcohol/vinyl acetate copolymer. Preferably, from about 0.5% to about 30% of the repeat units of said polymer are vinyl acetate.

20 A fifth aspect of the present invention relates to pharmaceutical compositions that include a solid amorphous dispersion comprising a low-solubility drug, a neutral dispersion polymer, and an excipient such as a base or a buffer.

25 In a preferred embodiment, the neutral dispersion polymer is concentration enhancing. Preferred neutral dispersion polymers are present in an amount sufficient to provide a maximum concentration of said low-solubility drug in a use environment that is at least 1.25-fold, preferably at least 2-fold, that provided by a second control composition comprising an equivalent quantity of said low-solubility drug and free from a concentration-enhancing polymer.

30 In another preferred embodiment, the neutral dispersion polymer is present in a sufficient amount so that said composition, when introduced to a use environment, provides an area under the concentration-versus time curve for any period of at least 90 minutes between the time of introduction to the use environment and about 270 minutes following introduction to the use environment that is at least 1.25-fold, preferably at least 2-fold, that of a second
35

control composition comprising an equivalent quantity of said low-solubility drug and free from a concentration-enhancing polymer.

In another preferred embodiment, the neutral dispersion polymer is present in a sufficient amount so that said composition provides a relative
5 bioavailability that is at least 1.25, preferably at least 2, relative to a second control composition comprising an equivalent quantity of said low-solubility drug and free from a concentration-enhancing polymer.

In another preferred embodiment, the drug in said composition has a relative degree of improvement in chemical stability of at least 1.25.

10 In yet another preferred embodiment, the composition provides improved chemical stability relative to a control composition, wherein said control comprising an equivalent quantity of a dispersion of said drug and said neutral dispersion polymer but free from said base and said buffer.

As described more fully below, the term "use environment" may refer to
15 the *in vivo* environment of the gastrointestinal (GI) tract of an animal, particularly a human, or the *in vitro* environment of a test solution, such as phosphate buffered saline (PBS) or model fasted duodenal (MFD) solution.

The composition may be dosed in a variety of dosage forms, including both immediate release and controlled release dosage forms, the latter
20 including both delayed and sustained release forms. The composition may include blends of polymers.

The various aspects of the present invention provide one or more of the following advantages. For those embodiments comprising an acid-sensitive drug, the inventors have recognized that a problem with forming dispersions of
25 acid-sensitive drugs is that for some dispersions, the drug does not remain chemically stable in the dispersion over time. The inventors have found that acid-sensitive drugs dispersed in an acidic polymer, such as hydroxypropyl methyl cellulose acetate succinate (which has carboxylic acid functional groups), have a tendency to chemically degrade over time. It is believed that
30 the presence of acidic ionic groups on the acidic polymer may either catalyze degradation of the drug or react directly with the drug. In any event, regardless of the particular degradation mechanism, the inventors have substantially reduced, if not eliminated the problem of drug degradation, by forming dispersions using neutral polymers, i.e. polymers that do not include acidic
35 functional groups. Thus, the present invention is able to realize the advantages

of forming dispersions of acid-sensitive drugs by improving the chemical stability of the acid-sensitive drug in the dispersion.

In addition, the invention in some embodiments further provides enhanced aqueous concentration and bioavailability for low-solubility drugs while at the same time minimizing the loss in potency and generation of impurities in the composition resulting from reaction or degradation of the drug when in the presence of an acidic species, such as an acidic dispersion polymer.

For those aspects comprising neutral vinyl copolymers of the present invention having hydroxyl-containing repeat units and hydrophobic repeat units, the compositions provide surprisingly effective concentration-enhancement. The polymers may be used with any low-solubility drug to improve the concentration of the drug in a use environment.

The foregoing and other objectives, features, and advantages of the invention will be more readily understood upon consideration of the following detailed description of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The pharmaceutical compositions of the present invention comprise solid amorphous dispersions of a drug and a neutral dispersion polymer. The present invention finds utility anytime it is desired to improve either the chemical stability of an acid-sensitive drug, to improve the concentration or bioavailability of a low-solubility drug, or both. In one embodiment of the invention, the dispersions of the present invention improve the chemical stability of acid-sensitive drugs. Thus, the dispersions may be used to prevent degradation of the drug due to interactions with dispersion polymers, acidic dispersion species, or other acidic excipients present in the composition. In another embodiment, the dispersions improve the concentration or bioavailability of a low-solubility drug. Suitable acid-sensitive drugs, low-solubility drugs, neutral polymers and methods for making the dispersions are discussed in more detail below.

ACID-SENSITIVE DRUGS

The term "drug" is conventional, denoting a compound having beneficial prophylactic and/or therapeutic properties when administered to an animal, especially humans. In one embodiment of the invention, the drug is an acid-sensitive drug, meaning that the drug either chemically reacts with or otherwise

degrades in the presence of acidic species. Acid-sensitive drugs often include functional groups which are reactive under acidic conditions, such as sulfonyl ureas, hydroxamic acids, hydroxy amides, carbamates, acetals, hydroxy ureas, esters, and amides. Drugs which include such functional groups may be prone to reactions such as hydrolysis, lactonization, or transesterification in the presence of acidic species.

Acid-sensitive drugs may be identified experimentally by determining whether the drug chemically reacts or degrades when dispersed in an acidic polymer. In particular, as used herein, the term "acid-sensitive drug" refers to a drug which, when dispersed in a "control acidic dispersion," degrades when stored under controlled aging conditions either for long storage times at ambient storage conditions or for short storage times under elevated temperature and relative humidity conditions.

The "control acidic dispersion" used to determine whether a drug is acid-sensitive is a dispersion of the drug and a pharmaceutically acceptable acidic polymer. A convenient pharmaceutically acceptable acidic polymer for use as the acidic dispersion polymer is the HF grade of hydroxypropyl methyl cellulose acetate succinate (HPMCAS), which is an acidic, or so-called enteric polymer having carboxylic acid functional groups, and a pK_a of about 5. The HPMCAS in the control acidic dispersion should have a minimum degree of substitution of succinate groups ($O(CO)CH_2CH_2(CO)OH$) of at least 4 wt% (or at least about 100 milliequivalents of carboxylic acid functional groups per mole of polymer). Alternatively, other acidic polymers which are at least as acidic as the HF grade of HPMCAS may be used, such as hydroxypropyl methyl cellulose acetate phthalate (HPMCP) or cellulose acetate phthalate (CAP). As with HPMCAS, such polymers should have at least about 100 milliequivalents of carboxylic acid functional groups per mole of polymer. The amount of acidic polymer present in the control acidic dispersion may vary, but should comprise at least 50 wt% or more of the control acidic dispersion. A drug is an acid-sensitive drug if it meets the drug degradation criteria described below in at least one control acidic dispersion comprised of 25 wt% drug and 75 wt% of the dispersion polymer HPMCAS, HPMCP or CAP.

In general, drug degradation may be measured using any conventional method for measuring the purity or potency of drug in a pharmaceutical composition. For example, the amount of active drug present in a dispersion

may be measured initially using high-performance liquid chromatography (HPLC) or any other analytical technique well known in the art. Alternatively, the amount of drug initially present may be calculated from the amount of drug present in the dispersion formulation. The potency of the dispersion may then
 5 be measured after storage at controlled temperature and humidity conditions for an appropriate period of time. A decrease in potency indicates that a chemical reaction has occurred, leading to a decrease in the amount of active drug present in the dispersion, and is an indication of poor chemical stability.

An alternative method used to evaluate chemical stability is to analyze
 10 the rate of increase in the amount of drug degradant(s) in the dispersion, which would indicate reaction of the drug to form the degradant(s). An HPLC or other analytical technique may be used to determine the concentration of drug degradant(s) in a dispersion. The amount of the degradant(s) is measured before and after storage under controlled aging conditions. The amount of
 15 increase in the drug degradant(s) may be used to determine the amount of decrease in "percent drug purity." The "percent drug purity" is defined as 100 times the total amount of drug present divided by the total amount of drug initially present. Thus, a percent drug purity at a given time may be calculated by the formula

$$\begin{array}{l} \text{20} \qquad \qquad \qquad \text{total amt. of} \\ \qquad \qquad \qquad \text{drug present} \\ \text{wt\% drug purity} = \left(\frac{\text{total amt. of}}{\text{drug init. present}} \right) * 100 \end{array}$$

25
 When the drug purity is calculated from the total amount of impurities, "percent drug purity" may be calculated by assuming that the "total amount of drug initially present," given in wt%, is equal to 100 wt% minus the wt% of total initial impurities, and that "total amount of drug present" is equal to 100 wt%
 30 minus the wt% of total impurities after storage, that is, at some later time. This method is equivalent to calculating "percent drug purity" by the formula

$$\begin{array}{l} \text{35} \qquad \qquad \qquad \text{total amt. of impurities} \\ \text{wt\% drug purity} = \left[1 - \left(\frac{\text{total amt. of drug}}{\text{init. present}} \right) \right] * 100 \end{array}$$

The rate at which drug degradation occurs is generally dependent on the storage conditions. The drug, when formulated as a composition of the present invention, should be stable at ambient temperature and humidity conditions (e.g., relative humidities of 20% to 60%) for long periods of time, such as months or years. However, to expedite testing, the storage conditions may employ elevated temperature and/or humidity to simulate longer storage times at ambient conditions. The storage time may vary from a few days to weeks or months, depending on the reactivity of the drug and the storage conditions.

A "degree of degradation" of drug following storage may be determined by subtracting the final percent drug purity (either determined by measuring the decrease in drug present or an increase in the amount of drug degradants present) from the initial percent drug purity. For example, for a dispersion initially containing 100 mg drug, and no measurable impurities, the initial percent drug purity is 100 wt%. If, after storage, the amount of drug in the dispersion decreases to 95 mg, the final percent drug purity would be 95 wt% and the "degree of degradation" is 5 wt% (100 wt% - 95 wt%). Alternatively, if 100 mg of drug substance were found to initially have 1 mg of impurities present, it would have an initial "percent drug purity" of 99 wt%. If, after storage, the total impurities present had increased to 6 wt%, the final percent drug purity would be 94 wt% and the "degree of degradation" would be 5 wt% (99 wt% - 94 wt%).

Alternatively, "degree of degradation" can be determined by subtracting from the amount of one or more specific degradants initially present from the amount of that specific drug degradant present after storage. Such a measure is useful where there are several drug degradants, of which only one (or a few) is of concern. The degree of degradation may be calculated on the basis of only those degradants that are of concern, rather than all of the degradant. For example, if a drug initially contained a specific degradant at a concentration of 1 wt% and after storage the concentration of that degradant was 6 wt%, the degree of degradation would be 5 wt% (6 wt%-1 wt%).

Returning now to the determination of an acid-sensitive drug, as used herein an "acid-sensitive" drug is one in which the degree of degradation of the drug in a control acidic dispersion (as defined above) is at least 0.01 wt% in a six month period at 40°C and 75% relative humidity (RH), or alternatively, in which the degree of degradation is at least 0.01 wt% in a one year period at

30°C at 60% RH, or as defined below with respect to degradation of drug in an acidic aqueous solution. These storage conditions are conventional and well-known in the art, and are the ICH recommendations for storage conditions to be used to evaluate the stability of a drug after storage for two years at ambient temperature and humidity. A relative degree of improvement may become apparent within a shorter time, such as three to five days, and shorter storage times may be used for some very acid-sensitive drugs.

The need for the present invention will generally be greater when the drug's reactivity with or sensitivity to acidic species increases. Dispersions of the present invention are preferred for acid-sensitive drugs having a degree of degradation that is greater than the minimum level when in the presence of an acidic dispersion polymer. Thus, the dispersions of the present invention are preferred for "very acid-sensitive drugs." Very acid-sensitive drugs are those which, when dispersed in a control acidic dispersion, have a degree of degradation of at least 0.1 wt% when stored at 40°C/75% RH for six months. The dispersions are even more preferred for drugs which have a degree of degradation of at least 1 wt%, and are most preferred for drugs which have a degree of degradation of at least 5.0 wt% when stored at 40°C/75% RH for six months.

In addition, the acid-sensitive drug when present in the control acidic dispersion degrades at a rate that is greater than the rate at which pure drug in the amorphous form, or if the amorphous form is physically unstable, the crystalline form, degrades. A relative degree of degradation of the drug in a control acidic dispersion to the pure form of the drug may be determined by taking the ratio of the degree of degradation of the drug in the control acidic dispersion and the degree of degradation of the pure form of the drug under the same storage conditions for the same storage time period. For example, where the degree of degradation of the drug in the control acidic dispersion is 5 wt% and the degree of degradation of the pure amorphous form of the drug is 0.1 wt%, the relative degree of degradation of the drug in the control acidic dispersion to the pure form of the drug is 50 (5 wt%/0.1 wt%). An acid-sensitive drug is one in which the relative degree of degradation of the drug in the control composition to the pure form of the drug is at least 2. The invention finds increasing utility as the relative degree of degradation increases above this minimum level. Thus, the invention is preferred for drugs having a relative

degree of degradation of about 3 or more, and more preferred for drugs having a relative degree of degradation of about 5 or more.

Alternatively, another test to determine whether a drug is an acid sensitive drug as used herein is to administer the drug to an acidic aqueous solution and plot drug concentration versus time. The acidic solution should have a pH of from 1-4. Drugs which are acid sensitive are those for which the drug concentration decreases by at least 1% within 24 hours of administration of the drug to the acidic solution. If the drug concentration changes by 1% in the 6-24 hour time period, then the drug is "slightly acid-sensitive." If the drug concentration changes by 1% in the 1-6 hour time period, then the drug is "moderately acid-sensitive." If the drug concentration changes by 1% in less than 1 hour, then the drug is "highly acid-sensitive." The present invention finds increasing utility for drugs which are slightly acid-sensitive, moderately acid-sensitive and highly acid-sensitive.

Specific examples of acid-sensitive drugs deliverable by the invention are set forth below, by way of example only. Each named drug should be understood to include the neutral form of the drug, pharmaceutically acceptable salts, and prodrugs. Examples of acid-sensitive drugs include quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3-fluorobenzyl-2(S),7-dihydroxy-7-methyloctyl]amide; quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide; quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide; (+)-N-[3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl]-N-hydroxyurea; omeprazole; etoposide; famotidine; erythromycin; quinapril; lansoprazole; and progabide.

LOW-SOLUBILITY DRUGS

In another embodiment of the invention, the drug is a "low-solubility drug," meaning that the drug may be either "substantially water-insoluble," which means that the drug has a minimum aqueous solubility at physiologically relevant pH (e.g., pH 1-8) of less than 0.01 mg/mL, "sparingly water-soluble," that is, has an aqueous solubility up to about 1 to 2 mg/mL, or even low to moderate aqueous-solubility, having an aqueous-solubility from about 1 mg/mL to as high as about 20 to 40 mg/mL. In general, it may be said that the drug has a dose-to-aqueous solubility ratio greater than 10 mL, and more typically greater than 100 mL, where the drug solubility (mg/mL) is the minimum value observed

in any physiologically relevant aqueous solution (e.g., those with pH values between 1 and 8) including USP simulated gastric and intestinal buffers, and dose is in mg. The dose-to-aqueous-solubility-ratio may be determined by simply dividing the dose (in mg) by the aqueous solubility (in mg/mL).

5 Preferred classes of drugs include, but are not limited to, antihypertensives, antianxiety agents, anticlotting agents, anticonvulsants, blood glucose-lowering agents, decongestants, antihistamines, antitussives, antineoplastics, beta blockers, anti-inflammatories, antipsychotic agents, cognitive enhancers, anti-atherosclerotic agents, cholesterol-reducing agents, 10 antiobesity agents, autoimmune disorder agents, anti-impotence agents, antibacterial and antifungal agents, hypnotic agents, anti-Parkinsonism agents, anti-Alzheimer's disease agents, antibiotics, anti-depressants, and antiviral agents, glycogen phosphorylase inhibitors, and cholesterol ester transfer protein inhibitors.

15 Each named drug should be understood to include the neutral form of the drug, pharmaceutically acceptable salts, as well as prodrugs. Specific examples of antihypertensives include prazosin, nifedipine, amlodipine besylate, trimazosin and doxazosin; specific examples of a blood glucose-lowering agent are glipizide and chlorpropamide; a specific example of an anti-impotence agent 20 is sildenafil and sildenafil citrate; specific examples of antineoplastics include chlorambucil, lomustine and echinomycin; a specific example of an imidazole-type antineoplastic is tubulazole; a specific example of an anti-hypercholesterolemic is atorvastatin calcium; specific examples of anxiolytics include hydroxyzine hydrochloride and doxepin hydrochloride; specific 25 examples of anti-inflammatory agents include betamethasone, prednisolone, aspirin, piroxicam, valdecoxib, carprofen, celecoxib, flurbiprofen and (+)-N-{4-[3-(4-fluorophenoxy)phenoxy]-2-cyclopenten-1-yl}-N-hydroxyurea; a specific example of a barbiturate is phenobarbital; specific examples of antivirals include acyclovir, nelfinavir, and virazole; specific examples of vitamins/nutritional 30 agents include retinol and vitamin E; specific examples of beta blockers include timolol and nadolol; a specific example of an emetic is apomorphine; specific examples of a diuretic include chlorthalidone and spironolactone; a specific example of an anticoagulant is dicumarol; specific examples of cardiotonics include digoxin and digitoxin; specific examples of androgens include 17- 35 methyltestosterone and testosterone; a specific example of a mineral corticoid is

desoxycorticosterone; a specific example of a steroidal hypnotic/anesthetic is alfaxalone; specific examples of anabolic agents include fluoxymesterone and methanstenolone; specific examples of antidepressant agents include sulpiride, [3,6-dimethyl-2-(2,4,6-trimethyl-phenoxy)-pyridin-4-yl]-(1-ethylpropyl)-amine, 3,5-dimethyl-4-(3'-pentoxy)-2-(2',4',6'-trimethylphenoxy)pyridine, pyroxidine, fluoxetine, paroxetine, venlafaxine and sertraline; specific examples of antibiotics include carbenicillin indanyl sodium, bacampicillin hydrochloride, troleandomycin, doxycycline hyclate, ampicillin and penicillin G; specific examples of anti-infectives include benzalkonium chloride and chlorhexidine; specific examples of coronary vasodilators include nitroglycerin and miflozine; a specific example of a hypnotic is etomidate; specific examples of carbonic anhydrase inhibitors include acetazolamide and chlorzotamide; specific examples of antifungals include econazole, terconazole, fluconazole, voriconazole, and griseofulvin; a specific example of an antiprotozoal is metronidazole; specific examples of anthelmintic agents include thiabendazole and oxfendazole and morantel; specific examples of antihistamines include astemizole, levocabastine, cetirizine, decarboethoxyloratadine and cinnarizine; specific examples of antipsychotics include ziprasidone, olanzapine, thiothixene hydrochloride, fluspirilene, risperidone and penfluridol; specific examples of gastrointestinal agents include loperamide and cisapride; specific examples of serotonin antagonists include ketanserin and mianserin; a specific example of an anesthetic is lidocaine; a specific example of a hypoglycemic agent is acetohexamide; a specific example of an anti-emetic is dimenhydrinate; a specific example of an antibacterial is cotrimoxazole; a specific example of a dopaminergic agent is L-DOPA; specific examples of anti-Alzheimer's Disease agents are THA and donepezil; a specific example of an anti-ulcer agent/H2 antagonist is famotidine; specific examples of sedative/hypnotic agents include chlordiazepoxide and triazolam; a specific example of a vasodilator is alprostadil; a specific example of a platelet inhibitor is prostacyclin; specific examples of ACE inhibitor/antihypertensive agents include enalaprilic acid and lisinopril; specific examples of tetracycline antibiotics include oxytetracycline and minocycline; specific examples of macrolide antibiotics include erythromycin, clarithromycin, and spiramycin; a specific example of an azalide antibiotic is azithromycin; specific examples of glycogen phosphorylase inhibitors include [R-(R'S')]5-chloro-N-[2-hydroxy-3-{methoxymethylamino}-3-

oxo-1-(phenylmethyl)propyl-1H-indole-2-carboxamide and 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-3-oxypropyl]amide; and specific examples of cholesterol ester transfer protein (CETP) inhibitors include [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester, [2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester, [2R, 4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester.

The invention is not limited by any particular structure or group of CETP inhibitors. Rather, the invention has general applicability to CETP inhibitors as a class, the class tending to be composed of compounds having low solubility. Compounds which may be the subject of the invention may be found in a number of patents and published applications, including DE 19741400 A1; DE 19741399 A1; WO 9914215 A1; WO 9914174; DE 19709125 A1; DE 19704244 A1; DE 19704243 A1; EP 818448 A1; WO 9804528 A2; DE 19627431 A1; DE 19627430 A1; DE 19627419 A1; EP 796846 A1; DE 19832159; DE 818197; DE 19741051; WO 9941237 A1; WO 9914204 A1; WO 9835937 A1; JP 11049743; WO 200018721; WO 200018723; WO 200018724; WO 200017164; WO 200017165; WO 200017166; EP 992496; and EP 987251, all of which are hereby incorporated by reference in their entireties for all purposes.

The invention is useful for CETP inhibitors that have sufficiently low aqueous solubility, low bioavailability or slow rate of absorption such that it is desirable to increase their concentration in an aqueous environment of use. Therefore, anytime one finds it desirable to raise the aqueous concentration of the CETP inhibitor in a use environment, the invention will find utility. The CETP inhibitor is "substantially water-insoluble" which means that the CETP inhibitor has a minimum aqueous solubility of less than about 0.01 mg/mL (or 10 μ g/ml) at any physiologically relevant pH (e.g., pH 1-8) and at about 22°C. (Unless otherwise specified, reference to aqueous solubility herein and in the claims is determined at about 22°C.) Compositions of the present invention find greater utility as the solubility of the CETP inhibitors decreases, and thus are preferred for CETP inhibitors with solubilities less than about 2 μ g/mL, and even more preferred for CETP inhibitors with solubilities less than about 0.5 μ g/mL.

Many CETP inhibitors have even lower solubilities (some even less than $0.1 \mu\text{g/mL}$), and require dramatic concentration enhancement to be sufficiently bioavailable upon oral dosing for effective plasma concentrations to be reached at practical doses.

5 In general, it may be said that the CETP inhibitor has a dose-to-aqueous solubility ratio greater than about 100 mL, where the solubility (mg/mL) is the minimum value observed in any physiologically relevant aqueous solution (e.g., those with pH values from 1 to 8) including USP simulated gastric and intestinal buffers, and dose is in mg. Compositions of the present invention, as
10 mentioned above, find greater utility as the solubility of the CETP inhibitor decreases and the dose increases. Thus, the compositions are preferred as the dose-to-solubility ratio increases, and thus are preferred for dose-to-solubility ratios greater than 1000 mL, and more preferred for dose-to-solubility ratios greater than about 5000 mL. The dose-to-solubility ratio may be determined by
15 dividing the dose (in mg) by the aqueous solubility (in mg/mL).

Oral delivery of many CETP inhibitors is particularly difficult because their aqueous solubility is usually extremely low, typically being less than $2 \mu\text{g/mL}$, often being less than $0.1 \mu\text{g/mL}$. Such low solubilities are a direct consequence of the particular structural characteristics of species that bind to
20 CETP and thus act as CETP inhibitors. This low solubility is primarily due to the hydrophobic nature of CETP inhibitors. Clog P, defined as the base 10 logarithm of the ratio of the drug solubility in octanol to the drug solubility in water, is a widely accepted measure of hydrophobicity. In general, Clog P values for CETP inhibitors are greater than 4 and are often greater than 5 to 7.
25 Thus, the hydrophobic and insoluble nature of CETP inhibitors as a class pose a particular challenge for oral delivery. Achieving therapeutic drug levels in the blood by oral dosing of practical quantities of drug generally requires a large enhancement in drug concentrations in the gastrointestinal fluid and a resulting large enhancement in bioavailability. Such enhancements in drug concentration
30 in gastrointestinal fluid typically need to be at least about 10-fold and often at least about 50-fold or even at least about 200-fold to achieve desired blood levels. Surprisingly, the dispersions of the present invention have proven to have the required large enhancements in drug concentration and bioavailability.

In contrast to conventional wisdom, the relative degree of enhancement
35 in aqueous concentration and bioavailability generally improves for CETP

inhibitors as solubility decreases and hydrophobicity increases. In fact, the inventors have recognized a subclass of these CETP inhibitors that are essentially aqueous insoluble, highly hydrophobic, and are characterized by a set of physical properties. This subclass exhibits dramatic enhancements in aqueous concentration and bioavailability when formulated using the compositions of the present invention.

The first property of this subclass of essentially insoluble, hydrophobic CETP inhibitors is extremely low aqueous solubility. By extremely low aqueous solubility is meant that the minimum aqueous solubility at physiologically relevant pH (pH of 1 to 8) is less than about 10 $\mu\text{g/ml}$ and preferably less than about 1 $\mu\text{g/ml}$.

A second property is a very high dose-to-solubility ratio. Extremely low solubility often leads to poor or slow absorption of the drug from the fluid of the gastrointestinal tract, when the drug is dosed orally in a conventional manner. For extremely low solubility drugs, poor absorption generally becomes progressively more difficult as the dose (mass of drug given orally) increases. Thus, a second property of this subclass of essentially insoluble, hydrophobic CETP inhibitors is a very high dose (in mg) to solubility (in mg/ml) ratio (ml). By "very high dose-to-solubility ratio" is meant that the dose-to-solubility ratio has a value of at least 1000 ml, and preferably at least 5,000 ml, and more preferably at least 10,000 ml.


A third property of this subclass of essentially insoluble, hydrophobic CETP inhibitors is that they are extremely hydrophobic. By extremely hydrophobic is meant that the Clog P value of the drug, has a value of at least 4.0, preferably a value of at least 5.0, and more preferably a value of at least 5.5.

A fourth property of this subclass of essentially insoluble CETP inhibitors is that they have a low melting point. Generally, drugs of this subclass will have a melting point of about 150°C or less, and preferably about 140°C or less.

Primarily, as a consequence of some or all of these four properties, CETP inhibitors of this subclass typically have very low absolute bioavailabilities. Specifically, the absolute bioavailability of drugs in this subclass when dosed orally in their undispersed state is less than about 10% and more often less than about 5%.

Turning now to the chemical structures of specific CETP inhibitors, one class of CETP inhibitors that finds utility with the present invention consists of oxy substituted 4-carboxyamino-2-methyl-1,2,3,4-tetrahydroquinolines having the Formula I

5



Formula I

and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

wherein R_{1-1} is hydrogen, Y_1 , W_1-X_1 , W_1-Y_1 ;

wherein W_1 is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_1 is $-O-Y_1$, $-S-Y_1$, $-N(H)-Y_1$ or $-N-(Y_1)_2$;

wherein Y_1 for each occurrence is independently Z_1 or a fully saturated, partially unsaturated or fully unsaturated one to ten membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with Z_1 ;

wherein Z_1 is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six

membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said Z_i substituent is optionally mono-, di- or tri-substituted independently with halo, (C_2-C_6) alkenyl, (C_1-C_6) alkyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxyl, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxyl, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl substituent is also optionally substituted with from one to nine fluorines; R_{1-3} is hydrogen or Q_i ;

wherein Q_i is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_i ;

wherein V_i is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_i substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carbamoyl, mono-N- or di-N,N- (C_1-C_6) alkylcarbamoyl, carboxyl, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxyl, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl or

(C₂-C₆)alkenyl substituents are also optionally substituted with from one to nine fluorines;

R_{1,4} is Q_{1,1} or V_{1,1}

5 wherein Q_{1,1} is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{1,1};

15 wherein V_{1,1} is a partially saturated, fully saturated or fully unsaturated three to six membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen;

20 wherein said V_{1,1} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, amino, nitro, cyano, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-substituted with oxo, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines;

25 wherein either R_{1,3} must contain V₁ or R_{1,4} must contain V_{1,1}; and R_{1,5}, R_{1,6}, R_{1,7} and R_{1,8} are each independently hydrogen, hydroxy or oxy wherein said oxy is substituted with T₁ or a partially saturated, fully saturated or fully unsaturated one to twelve membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with T₁;

35 wherein T₁ is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of

two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said T₁ substituent is optionally mono-, di- or tri-substituted independently with halo, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines.

Compounds of Formula I and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,140,342, United States Patent No. 6,362,198, and European Patent publication 987251, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula I:

[2R,4S] 4-[(3,5-dichloro-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-dinitro-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(2,6-dichloro-pyridin-4-ylmethyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-methoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-methoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-ethoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid 2,2,2-trifluoroethyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid tert-butyl ester;


[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-6-trifluoromethoxy-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester,

[2R,4S] (3,5-bis-trifluoromethyl-benzyl)-(1-butyryl-6,7-dimethoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-carbamic acid methyl ester;

[2R,4S] (3,5-bis-trifluoromethyl-benzyl)-(1-butyl-6,7-dimethoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-carbamic acid methyl ester; and

[2R,4S] (3,5-bis-trifluoromethyl-benzyl)-[1-(2-ethyl-butyl)-6,7-dimethoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-carbamic acid methyl ester, hydrochloride.

Another class of CETP inhibitors that finds utility with the present invention consists of 4-carboxyamino-2-methyl-1,2,3,4,-tetrahydroquinolines, having the Formula II



Formula II

and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

wherein R_{II-1} is hydrogen, Y_{II} , $W_{II}-X_{II}$, $W_{II}-Y_{II}$;

wherein W_{II} is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_{II} is $-O-Y_{II}$, $-S-Y_{II}$, $-N(H)-Y_{II}$ or $-N-(Y_{II})_2$;

wherein Y_{II} for each occurrence is independently Z_{II} or a fully saturated, partially unsaturated or fully unsaturated one to ten membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with Z_{II} ;

Z_{II} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said Z_{II} substituent is optionally mono-, di- or tri-substituted independently with halo, (C_2-C_6) alkenyl, (C_1-C_6) alkyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl is also optionally substituted with from one to nine fluorines;

R_{II-3} is hydrogen or Q_{II} ;

wherein Q_{II} is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{II} ;

wherein V_{II} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_{II} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxamoyl, mono-N- or di-N,N- (C_1-C_6) alkylcarboxamoyl, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino or said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituents are optionally substituted with from one to nine fluorines;

R_{II-4} is Q_{II-1} or V_{II-1}

wherein Q_{II-1} a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{II-1} ;

wherein V_{II-1} is a partially saturated, fully saturated or fully unsaturated three to six membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen;

wherein said V_{II-1} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, amino, nitro, cyano, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-substituted with oxo, said (C_1-C_6) alkyl substituent is optionally substituted with from one to nine fluorines;

wherein either R_{II-3} must contain V_{II} or R_{II-4} must contain V_{II-1} ; and

R_{II-5} , R_{II-6} , R_{II-7} and R_{II-8} are each independently hydrogen, a bond, nitro or halo wherein said bond is substituted with T_{II} or a partially saturated, fully saturated or fully unsaturated (C_1 - C_{12}) straight or branched carbon chain wherein carbon may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon is optionally mono-substituted with T_{II} ;

wherein T_{II} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said T_{II} substituent is optionally mono-, di- or tri-substituted independently with halo, (C_1 - C_6)alkyl, (C_2 - C_6)alkenyl, hydroxy, (C_1 - C_6)alkoxy, (C_1 - C_4)alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1 - C_6)alkyloxycarbonyl, mono-N- or di-N,N-(C_1 - C_6)alkylamino wherein said (C_1 - C_6)alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1 - C_6)alkoxy, (C_1 - C_4)alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1 - C_6)alkyloxycarbonyl, mono-N- or di-N,N-(C_1 - C_6)alkylamino, said (C_1 - C_6)alkyl substituent is also optionally substituted with from one to nine fluorines; provided that at least one of substituents R_{II-5} , R_{II-6} , R_{II-7} and R_{II-8} is not hydrogen and is not linked to the quinoline moiety through oxy.

Compounds of Formula II and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,147,090, United States Patent Application No. 09/671,400 filed September 27, 2000, and PCT Publication No. WO00/17166, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula II:

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-7-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-chloro-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-chloro-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

5 [2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2,6,7-trimethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-diethyl-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;


10 [2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-ethyl-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester; and

15 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester.

Another class of CETP inhibitors that finds utility with the present invention consists of annulated 4-carboxyamino-2-methyl-1,2,3,4,-tetrahydroquinolines, having the Formula III

20



Formula III

and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

25

wherein R_{III-1} is hydrogen, Y_{III} , $W_{III}-X_{III}$, $W_{III}-Y_{III}$;

wherein W_{III} is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_{III} is $-O-Y_{III}$, $-S-Y_{III}$, $-N(H)-Y_{III}$ or $-N-(Y_{III})_2$;

Y_{III} for each occurrence is independently Z_{III} or a fully saturated, partially unsaturated or fully unsaturated one to ten membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with Z_{III};

wherein Z_{III} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said Z_{III} substituent is optionally mono-, di- or tri-substituted independently with halo, (C₂-C₆)alkenyl, (C₁-C₆) alkyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl optionally substituted with from one to nine fluorines;

R_{III-3} is hydrogen or Q_{III};

wherein Q_{III} is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{III};

wherein V_{III} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_{III} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxamoyl, mono-N- or di-N,N- (C_1-C_6) alkylcarboxamoyl, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino or said (C_1-C_6) alkyl or (C_2-C_6) alkenyl are optionally substituted with from one to nine fluorines; R_{III-4} is Q_{III-1} or V_{III-1} ;

wherein Q_{III-1} a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{III-1} ;

wherein V_{III-1} is a partially saturated, fully saturated or fully unsaturated three to six membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen;

wherein said V_{III-1} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, amino, nitro, cyano, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-substituted with oxo, said (C_1-C_6) alkyl substituent optionally having from one to nine fluorines;

wherein either R_{III-3} must contain V_{III} or R_{III-4} must contain V_{III-1} ; and

R_{III-5} and R_{III-6}, or R_{III-6} and R_{III-7}, and/or R_{III-7} and R_{III-8} are taken together and form at least one four to eight membered ring that is partially saturated or fully unsaturated optionally having one to three heteroatoms independently selected from nitrogen, sulfur and oxygen;

5 wherein said ring or rings formed by R_{III-5} and R_{III-6}, or R_{III-6} and R_{III-7}, and/or R_{III-7} and R_{III-8} are optionally mono-, di- or tri-substituted independently with halo, (C₁-C₆)alkyl, (C₁-C₄)alkylsulfonyl, (C₂-C₆)alkenyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with
10 hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent optionally having from one to nine fluorines;

provided that the R_{III-5}, R_{III-6}, R_{III-7} and/or R_{III-8}, as the case may be, that
15 do not form at least one ring are each independently hydrogen, halo, (C₁-C₆)alkoxy or (C₁-C₆)alkyl, said (C₁-C₆)alkyl optionally having from one to nine fluorines.

Compounds of Formula III and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,147,089, United
20 States Patent No. 6,310,075, and European Patent Application No. 99307240.4 filed September 14, 1999, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula III:

25 [2R, 4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-2,3,4,6,7,8-hexahydro-cyclopenta[g]quinoline-1-carboxylic acid ethyl ester;

[6R, 8S] 8-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-methyl-3,6,7,8-tetrahydro-1H-2-thia-5-aza-cyclopenta[b]naphthalene-5-
30 carboxylic acid ethyl ester;

[6R, 8S] 8-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-methyl-3,6,7,8-tetrahydro-2H-furo[2,3-g]quinoline-5-carboxylic acid ethyl ester;

[2R, 4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-3,4,6,8-tetrahydro-2H-furo[3,4-g]quinoline-1-carboxylic acid ethyl ester;

optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with Z_{IV} ;

5 wherein Z_{IV} is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

10 wherein said Z_{IV} substituent is optionally mono-, di- or tri-substituted independently with halo, (C_2-C_6) alkenyl, (C_1-C_6) alkyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl substituent is also optionally substituted with from one to nine fluorines;

15 R_{IV-2} is a partially saturated, fully saturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with oxo, said carbon is optionally mono-substituted with hydroxy, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo; or said R_{IV-2} is a partially saturated, fully saturated or fully unsaturated three to seven membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen, wherein said R_{IV-2} ring is optionally attached through (C_1-C_4) alkyl;

25 wherein said R_{IV-2} ring is optionally mono-, di- or tri-substituted independently with halo, (C_2-C_6) alkenyl, (C_1-C_6) alkyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, oxo or (C_1-C_6) alkyloxycarbonyl;

with the proviso that R_{IV-2} is not methyl;

R_{IV-3} is hydrogen or Q_{IV} ;

wherein Q_{IV} is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons
5 other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo,
10 said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{IV} ;

wherein V_{IV} is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of
15 two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_{IV} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxamoyl, mono-N- or
20 di-N,N- (C_1-C_6) alkylcarboxamoyl, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl or
25 (C_2-C_6) alkenyl substituents are also optionally substituted with from one to nine fluorines;

R_{IV-4} is Q_{IV-1} or V_{IV-1} ;

wherein Q_{IV-1} a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons,
30 other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo,
35

said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with

V_{IV-1} ;

5 wherein V_{IV-1} is a partially saturated, fully saturated or fully unsaturated three to six membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen;

wherein said V_{IV-1} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, amino, nitro, cyano, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein
10 said (C_1-C_6) alkyl substituent is optionally mono-substituted with oxo, said (C_1-C_6) alkyl substituent is also optionally substituted with from one to nine fluorines;

wherein either R_{IV-3} must contain V_{IV} or R_{IV-4} must contain V_{IV-1} ;
 R_{IV-5} , R_{IV-6} , R_{IV-7} and R_{IV-8} are each independently hydrogen, a bond, nitro or halo wherein said bond is substituted with T_{IV} or a partially saturated, fully
15 saturated or fully unsaturated (C_1-C_{12}) straight or branched carbon chain wherein carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally
20 mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon is optionally mono-substituted with T_{IV} ;

wherein T_{IV} is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected
25 independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said T_{IV} substituent is optionally mono-, di- or tri-substituted
30 independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, $(C_1-$

C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines; and

wherein R_{IV-5} and R_{IV-6}, or R_{IV-6} and R_{IV-7}, and/or R_{IV-7} and R_{IV-8} may also be taken together and can form at least one four to eight membered ring that is partially saturated or fully unsaturated optionally having one to three heteroatoms independently selected from nitrogen, sulfur and oxygen;

wherein said ring or rings formed by R_{IV-5} and R_{IV-6}, or R_{IV-6} and R_{IV-7}, and/or R_{IV-7} and R_{IV-8} are optionally mono-, di- or tri-substituted independently with halo, (C₁-C₆)alkyl, (C₁-C₄)alkylsulfonyl, (C₂-C₆)alkenyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines; with the proviso that when R_{IV-2} is carboxyl or (C₁-C₄)alkylcarboxyl, then R_{IV-1} is not hydrogen.

Compounds of Formula IV and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,197,786, United States Application Serial No. 09/685,3000 filed 10/10/00, and PCT Publication No. WO 00/17164, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula IV:

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-isopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-chloro-2-cyclopropyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 2-cyclopropyl-4-[(3,5-dichloro-benzyl)-methoxycarbonyl-amino]-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;


[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid tert-butyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-3,4,6,7,8,9-hexahydro-2H-benzo[g]quinoline-1-carboxylic acid propyl ester;

5 [7R,9S] 9-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-methyl-1,2,3,7,8,9-hexahydro-6-aza-cyclopenta[a]naphthalene-6-carboxylic acid ethyl ester; and

[6S,8R] 6-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-8-methyl-1,2,3,6,7,8-hexahydro-9-aza-cyclopenta[a]naphthalene-9-carboxylic acid ethyl ester.

10 Another class of CETP inhibitors that finds utility with the present invention consists of 4-carboxyamino-2-substituted-1,2,3,4,-tetrahydroquinolines, having the Formula IV



15

Formula IV

and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

wherein R_{IV-1} is hydrogen, Y_{IV} , $W_{IV}-X_{IV}$ or $W_{IV}-Y_{IV}$;

20 wherein W_{IV} is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_{IV} is $-O-Y_{IV}$, $-S-Y_{IV}$, $-N(H)-Y_{IV}$ or $-N-(Y_{IV})_2$;

wherein Y_{IV} for each occurrence is independently Z_{IV} or a fully saturated, partially unsaturated or fully unsaturated one to ten membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is

25

[2R,4R] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

5 [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclobutyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

10 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methoxymethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

15 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid 2-hydroxyethyl ester;

20 [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;


[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

25 [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester; and

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester.

30 Another class of CETP inhibitors that finds utility with the present invention consists of 4-amino substituted-2-substituted-1,2,3,4,-tetrahydroquinolines, having the Formula V

38



Formula V

and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said
5 compounds;

wherein R_{V-1} is Y_V , W_V-X_V or W_V-Y_V ;

wherein W_V is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_V is $-O-Y_V$, $-S-Y_V$, $-N(H)-Y_V$ or $-N-(Y_V)_2$;

wherein Y_V for each occurrence is independently Z_V or a fully saturated,
10 partially unsaturated or fully unsaturated one to ten membered straight or
branched carbon chain wherein the carbons, other than the connecting carbon,
may optionally be replaced with one or two heteroatoms selected independently
from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-
substituted independently with halo, said carbon is optionally mono-substituted
15 with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is
optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or
di-substituted with oxo, and said carbon chain is optionally mono-substituted
with Z_V ;

wherein Z_V is a partially saturated, fully saturated or fully unsaturated
20 three to eight membered ring optionally having one to four heteroatoms selected
independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of
two fused partially saturated, fully saturated or fully unsaturated three to six
membered rings, taken independently, optionally having one to four
heteroatoms selected independently from nitrogen, sulfur and oxygen;

25 wherein said Z_V substituent is optionally mono-, di- or tri-substituted
independently with halo, (C_2-C_6) alkenyl, (C_1-C_6) alkyl, hydroxy, (C_1-C_6) alkoxy,
 (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl,
mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is
optionally mono-, di- or tri-substituted independently with halo, hydroxy, $(C_1-$

C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines;

5 R_{V,2} is a partially saturated, fully saturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with oxo, said carbon is optionally mono-substituted with hydroxy, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo; or said R_{V,2} is a partially saturated, fully saturated or fully unsaturated three to seven membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen, wherein said R_{V,2} ring is optionally attached through
10 (C₁-C₄)alkyl;
15

wherein said R_{V,2} ring is optionally mono-, di- or tri-substituted independently with halo, (C₂-C₆)alkenyl, (C₁-C₆) alkyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, oxo or (C₁-C₆)alkyloxycarbonyl;
20 R_{V,3} is hydrogen or Q_V;

wherein Q_V is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_V;
25
30

wherein V_V is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six
35

membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_V substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxamoyl, mono-N- or di-N,N- (C_1-C_6) alkylcarboxamoyl, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituents are also optionally substituted with from one to nine fluorines;

R_{V-4} is cyano, formyl, $W_{V-1}Q_{V-1}$, $W_{V-1}V_{V-1}$, (C_1-C_4) alkylene V_{V-1} or V_{V-2} ;

wherein W_{V-1} is carbonyl, thiocarbonyl, SO or SO_2 ,

wherein Q_{V-1} a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{V-1} ;

wherein V_{V-1} is a partially saturated, fully saturated or fully unsaturated three to six membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_{V-1} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, hydroxy, oxo, amino, nitro, cyano, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-substituted with oxo, said (C_1-C_6) alkyl substituent is also optionally substituted with from one to nine fluorines;

wherein V_{V-2} is a partially saturated, fully saturated or fully unsaturated five to seven membered ring containing one to four heteroatoms selected independently from oxygen, sulfur and nitrogen;

wherein said V_{V-2} substituent is optionally mono-, di- or tri-substituted independently with halo, (C_1-C_2) alkyl, (C_1-C_2) alkoxy, hydroxy, or oxo wherein said (C_1-C_2) alkyl optionally has from one to five fluorines; and

wherein R_{V-4} does not include oxycarbonyl linked directly to the C^4 nitrogen;

wherein either R_{V-3} must contain V_V or R_{V-4} must contain V_{V-1} ;

R_{V-5} , R_{V-6} , R_{V-7} and R_{V-8} are independently hydrogen, a bond, nitro or halo wherein said bond is substituted with T_V or a partially saturated, fully saturated or fully unsaturated (C_1-C_{12}) straight or branched carbon chain wherein carbon may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen, wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with T_V ;

wherein T_V is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said T_V substituent is optionally mono-, di- or tri-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl substituent also optionally has from one to nine fluorines;

wherein R_{V-5} and R_{V-6} , or R_{V-6} and R_{V-7} , and/or R_{V-7} and R_{V-8} may also be taken together and can form at least one ring that is a partially saturated or fully

unsaturated four to eight membered ring optionally having one to three heteroatoms independently selected from nitrogen, sulfur and oxygen;

wherein said rings formed by R_{V-5} and R_{V-6} , or R_{V-6} and R_{V-7} , and/or R_{V-7} and R_{V-8} are optionally mono-, di- or tri-substituted independently with halo, (C₁-C₆)alkyl, (C₁-C₄)alkylsulfonyl, (C₂-C₆)alkenyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent also optionally has from one to nine fluorines.

Compounds of Formula V and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,140,343, United States Patent Application Serial No. 09/671,221 filed September 27, 2000, and PCT Publication No. WO 00/17165, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula V:

- [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
- [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester;
- [2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid tert-butyl ester;
- [2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
- [2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- [2S,4S] 4-[1-(3,5-bis-trifluoromethyl-benzyl)-ureido]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
- [2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- [2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-methoxymethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester;

[2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

5 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

10 [2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;


[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

15 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester; and

[2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester.

20 Another class of CETP inhibitors that finds utility with the present invention consists of cycloalkano-pyridines having the Formula VI



Formula VI


and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

25 in which

A_{VI} denotes an aryl containing 6 to 10 carbon atoms, which is optionally substituted with up to five identical or different substituents in the form of a halogen, nitro, hydroxyl, trifluoromethyl, trifluoromethoxy or a straight-chain or branched alkyl, acyl, hydroxyalkyl or alkoxy containing up to 7 carbon atoms
30 each, or in the form of a group according to the formula $-BNR_{VI-3}R_{VI-4}$, wherein

R_{VI-3} and R_{VI-4} are identical or different and denote a hydrogen, phenyl or a straight-chain or branched alkyl containing up to 6 carbon atoms,

D_{VI} denotes an aryl containing 6 to 10 carbon atoms, which is optionally substituted with a phenyl, nitro, halogen, trifluoromethyl or trifluoromethoxy, or a radical according to the formula $R_{VI-5}-L_{VI-}$,




or $R_{VI-9}-T_{VI}-V_{VI}-X_{VI}$, wherein

R_{VI-5} , R_{VI-6} and R_{VI-9} denote, independently from one another, a cycloalkyl containing 3 to 6 carbon atoms, or an aryl containing 6 to 10 carbon atom or a 5- to 7-membered, optionally benzo-condensed, saturated or unsaturated, mono-, bi- or tricyclic heterocycle containing up to 4 heteroatoms from the series of S, N and/or O, wherein the rings are optionally substituted, in the case of the nitrogen-containing rings also via the N function, with up to five identical or different substituents in the form of a halogen, trifluoromethyl, nitro, hydroxyl, cyano, carboxyl, trifluoromethoxy, a straight-chain or branched acyl, alkyl, alkylthio, alkylalkoxy, alkoxy or alkoxycarbonyl containing up to 6 carbon atoms each, an aryl or trifluoromethyl-substituted aryl containing 6 to 10 carbon atoms each, or an optionally benzo-condensed, aromatic 5- to 7-membered heterocycle containing up to 3 heteroatoms from the series of S, N and/or O, and/or in the form of a group according to the formula BOR_{VI-10} , $-SR_{VI-11}$, $-SO_2R_{VI-12}$ or $BNR_{VI-13}R_{VI-14}$, wherein

R_{VI-10} , R_{VI-11} and R_{VI-12} denote, independently from one another, an aryl containing 6 to 10 carbon atoms, which is in turn substituted with up to two identical or different substituents in the form of a phenyl, halogen or a straight-chain or branched alkyl containing up to 6 carbon atoms,


R_{VI-13} and R_{VI-14} are identical or different and have the meaning of R_{VI-3} and R_{VI-4} given above, or

R_{VI-5} and/or R_{VI-6} denote a radical according to the formula



R_{VI-7} denotes a hydrogen or halogen, and

R_{VI-8} denotes a hydrogen, halogen, azido, trifluoromethyl, hydroxyl, trifluoromethoxy, a straight-chain or branched alkoxy or alkyl containing up to 6 carbon atoms each, or a radical according to the formula



wherein

R_{VI-15} and R_{VI-16} are identical or different and have the meaning of R_{VI-3} and R_{VI-4} given above, or

R_{VI-7} and R_{VI-8} together form a radical according to the formula $=O$ or $=NR_{VI-17}$, wherein

R_{VI-17} denotes a hydrogen or a straight-chain or branched alkyl, alkoxy or acyl containing up to 6 carbon atoms each,

L_{VI} denotes a straight-chain or branched alkylene or alkenylene chain containing up to 8 carbon atoms each, which are optionally substituted with up to two hydroxyl groups,

T_{VI} and X_{VI} are identical or different and denote a straight-chain or branched alkylene chain containing up to 8 carbon atoms, or


T_{VI} or X_{VI} denotes a bond,

V_{VI} denotes an oxygen or sulfur atom or an BNR_{VI-18} group, wherein

R_{VI-18} denotes a hydrogen or a straight-chain or branched alkyl containing up to 6 carbon atoms or a phenyl,

E_{VI} denotes a cycloalkyl containing 3 to 8 carbon atoms, or a straight-chain or branched alkyl containing up to 8 carbon atoms, which is optionally substituted with a cycloalkyl containing 3 to 8 carbon atoms or a hydroxyl, or a phenyl, which is optionally substituted with a halogen or trifluoromethyl,

R_{VI-1} and R_{VI-2} together form a straight-chain or branched alkylene chain containing up to 7 carbon atoms, which must be substituted with a carbonyl group and/or a radical according to the formula



wherein

a and b are identical or different and denote a number equaling 1, 2 or 3,

R_{VI-19} denotes a hydrogen atom, a cycloalkyl containing 3 to 7 carbon atoms, a straight-chain or branched silylalkyl containing up to 8 carbon atoms, or a straight-chain or branched alkyl containing up to 8 carbon atoms, which is optionally substituted with a hydroxyl, a straight-chain or a branched alkoxy containing up to 6 carbon atoms or a phenyl, which may in turn be substituted with a halogen, nitro, trifluoromethyl, trifluoromethoxy or phenyl or tetrazole-substituted phenyl, and an alkyl that is optionally substituted with a group according to the formula BOR_{VI-22} , wherein


R_{VI-22} denotes a straight-chain or branched acyl containing up to 4 carbon atoms or benzyl, or

R_{VI-19} denotes a straight-chain or branched acyl containing up to 20 carbon atoms or benzoyl, which is optionally substituted with a halogen, trifluoromethyl, nitro or trifluoromethoxy, or a straight-chain or branched fluoroacyl containing up to 8 carbon atoms,

R_{VI-20} and R_{VI-21} are identical or different and denote a hydrogen, phenyl or a straight-chain or branched alkyl containing up to 6 carbon atoms, or

R_{VI-20} and R_{VI-21} together form a 3- to 6-membered carbocyclic ring, and the carbocyclic rings formed are optionally substituted, optionally also geminally, with up to six identical or different substituents in the form of trifluoromethyl, hydroxyl, nitrile, halogen, carboxyl, nitro, azido, cyano, cycloalkyl or cycloalkyloxy containing 3 to 7 carbon atoms each, a straight-chain or branched alkoxycarbonyl, alkoxy or alkylthio containing up to 6 carbon atoms each, or a straight-chain or branched alkyl containing up to 6 carbon atoms, which is in turn substituted with up to two identical or different substituents in the form of a hydroxyl, benzyloxy, trifluoromethyl, benzoyl, a straight-chain or branched alkoxy, oxyacyl or carboxyl containing up to 4 carbon atoms each and/or a phenyl, which may in turn be substituted with a halogen, trifluoromethyl or trifluoromethoxy, and/or the carbocyclic rings formed are optionally substituted, also geminally, with up to five identical or different substituents in the form of a phenyl, benzoyl, thiophenyl or sulfonylbenzyl, which in turn are optionally substituted with a halogen, trifluoromethyl, trifluoromethoxy or nitro, and/or optionally in the form of a radical according to the formula

47




wherein

c is a number equaling 1, 2, 3 or 4,

5 d is a number equaling 0 or 1,

$\text{R}_{\text{VI-23}}$ and $\text{R}_{\text{VI-24}}$ are identical or different and denote a hydrogen, cycloalkyl containing 3 to 6 carbon atoms, a straight-chain or branched alkyl containing up to 6 carbon atoms, benzyl or phenyl, which is optionally substituted with up to two identical or different substituents in the form of halogen, trifluoromethyl, cyano, phenyl or nitro, and/or the carbocyclic rings formed are optionally substituted with a spiro-linked radical according to the formula



wherein

15 W_{VI} denotes either an oxygen atom or a sulfur atom,

Y_{VI} and Y'_{VI} together form a 2- to 6-membered straight-chain or branched alkylene chain,


e is a number equaling 1, 2, 3, 4, 5, 6 or 7,

f is a number equaling 1 or 2,

20 $\text{R}_{\text{VI-25}}$, $\text{R}_{\text{VI-26}}$, $\text{R}_{\text{VI-27}}$, $\text{R}_{\text{VI-28}}$, $\text{R}_{\text{VI-29}}$, $\text{R}_{\text{VI-30}}$ and $\text{R}_{\text{VI-31}}$ are identical or different and denote a hydrogen, trifluoromethyl, phenyl, halogen or a straight-chain or branched alkyl or alkoxy containing up to 6 carbon atoms each, or

$\text{R}_{\text{VI-25}}$ and $\text{R}_{\text{VI-26}}$ or $\text{R}_{\text{VI-27}}$ and $\text{R}_{\text{VI-28}}$ each together denote a straight-chain or branched alkyl chain containing up to 6 carbon atoms or

25 $\text{R}_{\text{VI-25}}$ and $\text{R}_{\text{VI-26}}$ or $\text{R}_{\text{VI-27}}$ and $\text{R}_{\text{VI-28}}$ each together form a radical according to the formula



wherein

W_{VI} has the meaning given above,

g is a number equaling 1, 2, 3, 4, 5, 6 or 7,

R_{VI-32} and R_{VI-33} together form a 3- to 7-membered heterocycle, which contains an oxygen or sulfur atom or a group according to the formula SO, SO₂ or BNR_{VI-34}, wherein

5 R_{VI-34} denotes a hydrogen atom, a phenyl, benzyl, or a straight-chain or branched alkyl containing up to 4 carbon atoms, and salts and N oxides thereof, with the exception of 5(6H)-quinolones, 3-benzoyl-7,8-dihydro-2,7,7-trimethyl-4-phenyl.


10 Compounds of Formula VI and their methods of manufacture are disclosed in European Patent Application No. EP 818448 A1, United States Patent No. 6,207,671 and United States Patent No. 6,069,148, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula VI:

15 2-cyclopentyl-4-(4-fluorophenyl)-7,7-dimethyl-3-(4-trifluoromethylbenzoyl)-4,6,7,8-tetrahydro-1H-quinolin-5-one;
2-cyclopentyl-4-(4-fluorophenyl)-7,7-dimethyl-3-(4-trifluoromethylbenzoyl)-7,8-dihydro-6H-quinolin-5-one;
[2-cyclopentyl-4-(4-fluorophenyl)-5-hydroxy-7,7-dimethyl-5,6,7,8-
20 tetrahydroquinolin-3-yl]-(4-trifluoromethylphenyl)-methanone;
[5-(t-butyldimethylsilyloxy)-2-cyclopentyl-4-(4-fluorophenyl)-7,7-dimethyl-5,6,7,8-tetrahydroquinolin-3-yl]-(4-trifluoromethylphenyl)-methanone;
[5-(t-butyldimethylsilyloxy)-2-cyclopentyl-4-(4-fluorophenyl)-7,7-dimethyl-5,6,7,8-tetrahydroquinolin-3-yl]-(4-trifluoromethylphenyl)-methanol;
25 5-(t-butyldimethylsilyloxy)-2-cyclopentyl-4-(4-fluorophenyl)-3-[fluoro-(4-trifluoromethylphenyl)-methyl]-7,7-dimethyl-5,6,7,8-tetrahydroquinoline; and
2-cyclopentyl-4-(4-fluorophenyl)-3-[fluoro-(4-trifluoromethylphenyl)-methyl]-7,7-dimethyl-5,6,7,8-tetrahydroquinolin-5-ol.

30 Another class of CETP inhibitors that finds utility with the present invention consists of substituted-pyridines having the Formula VII

49




Formula VII

or a pharmaceutically acceptable salt or tautomer thereof,
wherein

5 R_{VII-2} and R_{VII-6} are independently selected from the group consisting of hydrogen, hydroxy, alkyl, fluorinated alkyl, fluorinated aralkyl, chlorofluorinated alkyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, alkoxy, alkoxyalkyl, and alkoxyacetyl; provided that at least one of R_{VII-2} and R_{VII-6} is fluorinated alkyl, chlorofluorinated alkyl or alkoxyalkyl;

10 R_{VII-3} is selected from the group consisting of hydroxy, amido, arylcarbonyl, heteroarylcarbonyl, hydroxymethyl -CHO, -CO₂R_{VII-7}, wherein R_{VII-7} is selected from the group consisting of hydrogen, alkyl and cyanoalkyl; and




15 wherein R_{VII-15a} is selected from the group consisting of hydroxy, hydrogen, halogen, alkylthio, alkenylthio, alkynylthio, arylthio, heteroarylthio, heterocyclylthio, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy and heterocycliloxy, and

20 R_{VII-16a} is selected from the group consisting of alkyl, haloalkyl, alkenyl, haloalkenyl, alkynyl, haloalkynyl, aryl, heteroaryl, and heterocyclyl, arylalkoxy, trialkylsilyloxy;

25 R_{VII-4} is selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, aryl, heteroaryl, heterocyclyl, cycloalkylalkyl, cycloalkenylalkyl, aralkyl, heteroarylalkyl, heterocyclylalkyl, cycloalkylalkenyl, cycloalkenylalkenyl, aralkenyl, heteroarylalkenyl, heterocyclylalkenyl, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, heterocycliloxy, alkanoyloxy, alkenoyloxy, alkynoyloxy,


- aryloyloxy, heteroaryloxy, heterocyclyloxy, alkoxycarbonyl, alkenoxycarbonyl, alkynoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, heterocyclyloxycarbonyl, thio, alkylthio, alkenylthio, alkynylthio, arylthio, heteroarylthio, heterocyclylthio, cycloalkylthio, cycloalkenylthio, alkylthioalkyl, alkenylthioalkyl, alkynylthioalkyl, arylthioalkyl, heteroarylthioalkyl, heterocyclylthioalkyl, alkylthioalkenyl, alkenylthioalkenyl, alkynylthioalkenyl, arylthioalkenyl, heteroarylthioalkenyl, heterocyclylthioalkenyl, alkylamino, alkenylamino, alkynylamino, arylamino, heteroarylamino, heterocyclylamino, aryldialkylamino, diarylamino, diheteroarylamino, alkylarylamino, alkylheteroarylamino, arylheteroarylamino, trialkylsilyl, trialkenylsilyl, triarylsilyl, -CO(O)N(R_{VII-8a}R_{VII-8b}), wherein R_{VII-8a} and R_{VII-8b} are independently selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl, -SO₂R_{VII-9}, wherein R_{VII-9} is selected from the group consisting of hydroxy, alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl, -OP(O)(OR_{VII-10a})(OR_{VII-10b}), wherein R_{VII-10a} and R_{VII-10b} are independently selected from the group consisting of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl, and -OP(S)(OR_{VII-11a})(OR_{VII-11b}), wherein R_{VII-11a} and R_{VII-11b} are independently selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl;
- R_{VII-5} is selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, aryl, heteroaryl, heterocyclyl, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, heterocyclyloxy, alkylcarbonyloxyalkyl, alkenylcarbonyloxyalkyl, alkynylcarbonyloxyalkyl, arylcarbonyloxyalkyl, heteroarylcarbonyloxyalkyl, heterocyclylcarbonyloxyalkyl, cycloalkylalkyl, cycloalkenylalkyl, aralkyl, heteroarylalkyl, heterocyclylalkyl, cycloalkylalkenyl, cycloalkenylalkenyl, aralkenyl, heteroarylalkenyl, heterocyclylalkenyl, alkylthioalkyl, cycloalkylthioalkyl, alkenylthioalkyl, alkynylthioalkyl, arylthioalkyl, heteroarylthioalkyl, heterocyclylthioalkyl, alkylthioalkenyl, alkenylthioalkenyl, alkynylthioalkenyl, arylthioalkenyl, heteroarylthioalkenyl, heterocyclylthioalkenyl, alkoxyalkyl, alkenoxyalkyl, alkynoxyalkyl, aryloxyalkyl, heteroaryloxyalkyl, heterocyclyloxyalkyl, alkoxyalkenyl, alkenoxyalkenyl, alkynoxyalkenyl, aryloxyalkenyl, heteroaryloxyalkenyl, heterocyclyloxyalkenyl, cyano, hydroxymethyl, -CO₂R_{VII-14}, wherein R_{VII-14} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl;

51




wherein $R_{VII-15b}$ is selected from the group consisting of hydroxy, hydrogen, halogen, alkylthio, alkenylthio, alkynylthio, arylthio, heteroarylthio, heterocyclylthio, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, heterocycloxy, aroyloxy, and alkylsulfonyloxy, and

$R_{VII-16b}$ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, arylalkoxy, and trialkylsilyloxy;



wherein R_{VII-17} and R_{VII-18} are independently selected from the group consisting of alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl;




wherein R_{VII-19} is selected from the group consisting of alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, $-SR_{VII-20}$, $-OR_{VII-21}$, and $BR_{VII-22}CO_2R_{VII-23}$, wherein

R_{VII-20} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aminoalkyl, aminoalkenyl, aminoalkynyl, aminoaryl, aminoheteroaryl, aminoheterocyclyl, alkylheteroaryl amino, arylheteroaryl amino,


R_{VII-21} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl,

R_{VII-22} is selected from the group consisting of alkylene or arylene, and

R_{VII-23} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;




wherein R_{VII-24} is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, aralkenyl, and aralkynyl;






5

wherein R_{VII-25} is heterocyclidenyl;




wherein R_{VII-26} and R_{VII-27} are independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;



10


wherein R_{VII-28} and R_{VII-29} are independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;

15





53


wherein R_{VII-30} and R_{VII-31} are independently alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, and heterocyclyloxy; and



5 wherein R_{VII-32} and R_{VII-33} are independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;




wherein R_{VII-36} is selected from the group consisting of alkyl, alkenyl, aryl, heteroaryl and heterocyclyl;



10


wherein R_{VII-37} and R_{VII-38} are independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;



15 wherein R_{VII-39} is selected from the group consisting of hydrogen, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, heterocyclyloxy, alkylthio, alkenylthio, alkynylthio, arylthio, heteroarylthio and heterocyclylthio, and


R_{VII-40} is selected from the group consisting of haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl, haloheterocyclyl, cycloalkyl, cycloalkenyl, heterocyclylalkoxy, heterocyclylalkenoxy, heterocyclylalkynoxy, alkylthio, alkenylthio, alkynylthio, arylthio, heteroarylthio and heterocyclylthio;

20




wherein R_{VII-41} is heterocyclylidenyl;

54





wherein $\text{R}_{\text{VII-42}}$ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl, and

$\text{R}_{\text{VII-43}}$ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl, and haloheterocyclyl;



wherein $\text{R}_{\text{VII-44}}$ is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl;


10



15

wherein $\text{R}_{\text{VII-45}}$ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl, haloheterocyclyl, heterocyclyl, cycloalkylalkyl, cycloalkenylalkyl, aralkyl, heteroarylalkyl, heterocyclylalkyl, cycloalkylalkenyl, cycloalkenylalkenyl, aralkenyl, heteroarylalkenyl, heterocyclylalkenyl, alkylthioalkyl, alkenylthioalkyl, alkynylthioalkyl, arylthioalkyl, heteroarylthioalkyl, heterocyclylthioalkyl, alkylthioalkenyl, alkenylthioalkenyl, alkynylthioalkenyl, arylthioalkenyl, heteroarylthioalkenyl, heterocyclylthioalkenyl, aminocarbonylalkyl, aminocarbonylalkenyl, aminocarbonylalkynyl, aminocarbonylaryl, aminocarbonylheteroaryl, and aminocarbonylheterocyclyl,

25




wherein $\text{R}_{\text{VII-46}}$ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl, and

$\text{R}_{\text{VII-47}}$ is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl; and


30

55




wherein $\text{R}_{\text{VII-48}}$ is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl, and


$\text{R}_{\text{VII-49}}$ is selected from the group consisting of alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, heterocycliloxy, haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl and haloheterocyclyl;



wherein $\text{R}_{\text{VII-50}}$ is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy and heterocycliloxy;



wherein $\text{R}_{\text{VII-51}}$ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl and haloheterocyclyl; and



wherein $\text{R}_{\text{VII-53}}$ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl;


provided that when $\text{R}_{\text{VII-5}}$ is selected from the group consisting of heterocyclylalkyl and heterocyclylalkenyl, the heterocyclyl radical of the corresponding heterocyclylalkyl or heterocyclylalkenyl is other than δ -lactone; and

provided that when $\text{R}_{\text{VII-4}}$ is aryl, heteroaryl or heterocyclyl, and one of $\text{R}_{\text{VII-2}}$ and $\text{R}_{\text{VII-6}}$ is trifluoromethyl, then the other of $\text{R}_{\text{VII-2}}$ and $\text{R}_{\text{VII-6}}$ is difluoromethyl.

Compounds of Formula VII and their methods of manufacture are disclosed in PCT Publication No. WO 9941237-A1, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor of Formula VII is dimethyl 5,5-dithiobis[2-difluoromethyl-4-(2-methylpropyl)-6-(trifluoromethyl)-3-pyridine-carboxylate].

Another class of CETP inhibitors that finds utility with the present invention consists of substituted biphenyls having the Formula VIII



Formula VIII

or a pharmaceutically acceptable salt, enantiomers, or stereoisomers thereof, in which

A_{VIII} stands for aryl with 6 to 10 carbon atoms, which is optionally substituted up to 3 times in an identical manner or differently by halogen, hydroxy, trifluoromethyl, trifluoromethoxy, or by straight-chain or branched alkyl, acyl, or alkoxy with up to 7 carbon atoms each, or by a group of the formula -NR_{VIII-1}R_{VIII-2}, wherein

R_{VIII-1} and R_{VIII-2} are identical or different and denote hydrogen, phenyl, or straight-chain or branched alkyl with up to 6 carbon atoms,

D_{VIII} stands for straight-chain or branched alkyl with up to 8 carbon atoms, which is substituted by hydroxy,

E_{VIII} and L_{VIII} are either identical or different and stand for straight-chain or branched alkyl with up to 8 carbon atoms, which is optionally substituted by cycloalkyl with 3 to 8 carbon atoms, or stands for cycloalkyl with 3 to 8 carbon atoms, or

E_{VIII} has the above-mentioned meaning and

L_{VIII} in this case stands for aryl with 6 to 10 carbon atoms, which is optionally substituted up to 3 times in an identical manner or differently by halogen, hydroxy, trifluoromethyl, trifluoromethoxy, or by straight-chain or branched alkyl, acyl, or alkoxy with up to 7 carbon atoms each, or by a group of the formula

-NR_{VIII-3}R_{VIII-4}, wherein

R_{VIII-3} and R_{VIII-4} are identical or different and have the meaning given above for R_{VIII-1} and R_{VIII-2}, or


5 E_{VIII} stands for straight-chain or branched alkyl with up to 8 carbon atoms, or stands for aryl with 6 to 10 carbon atoms, which is optionally substituted up to 3 times in an identical manner or differently by halogen, hydroxy, trifluoromethyl, trifluoromethoxy, or by straight-chain or branched alkyl, acyl, or alkoxy with up to 7 carbon atoms each, or by a group of the formula

-NR_{VIII-5}R_{VIII-6}, wherein

10 R_{VIII-5} and R_{VIII-6} are identical or different and have the meaning given above for R_{VIII-1} and R_{VIII-2}, and

L_{VIII} in this case stands for straight-chain or branched alkoxy with up to 8 carbon atoms or for cycloalkoxy with 3 to 8 carbon atoms,

T_{VIII} stands for a radical of the formula



R_{VIII-7} and R_{VIII-8} are identical or different and denote cycloalkyl with 3 to 8 carbon atoms, or aryl with 6 to 10 carbon atoms, or denote a 5- to 7-member aromatic, optionally benzo-condensed, heterocyclic compound with up to 3 heteroatoms from the series S, N and/or O, which are optionally substituted up to 3 times in an identical manner or differently by trifluoromethyl, trifluoromethoxy, halogen, hydroxy, carboxyl, by straight-chain or branched alkyl, acyl, alkoxy, or alkoxycarbonyl with up to 6 carbon atoms each, or by phenyl, phenoxy, or thiophenyl, which can in turn be substituted by halogen, trifluoromethyl, or trifluoromethoxy, and/or the rings are substituted by a group

20 of the formula

25

-NR_{VIII-11}R_{VIII-12}, wherein

R_{VIII-11} and R_{VIII-12} are identical or different and have the meaning given above for R_{VIII-1} and R_{VIII-2},

30 X_{VIII} denotes a straight or branched alkyl chain or alkenyl chain with 2 to 10 carbon atoms each, which are optionally substituted up to 2 times by hydroxy,

R_{VIII-9} denotes hydrogen, and

$R_{VIII-10}$ denotes hydrogen, halogen, azido, trifluoromethyl, hydroxy, mercapto, trifluoromethoxy, straight-chain or branched alkoxy with up to 5 carbon atoms, or a radical of the formula


$-NR_{VIII-13}R_{VIII-14}$, wherein

5 $R_{VIII-13}$ and $R_{VIII-14}$ are identical or different and have the meaning given above for R_{VIII-1} and R_{VIII-2} , or

R_{VIII-9} and $R_{VIII-10}$ form a carbonyl group together with the carbon atom.

Compounds of Formula VIII are disclosed in PCT Publication No. WO 9804528, which is incorporated herein by reference in its entirety for all
10 purposes.

Another class of CETP inhibitors that finds utility with the present invention consists of substituted 1,2,4-triazoles having the Formula IX



Formula IX

or a pharmaceutically acceptable salt or tautomer thereof;

15 wherein R_{IX-1} is selected from higher alkyl, higher alkenyl, higher alkynyl, aryl, aralkyl, aryloxyalkyl, alkoxyalkyl, alkylthioalkyl, arylthioalkyl, and cycloalkylalkyl;

wherein R_{IX-2} is selected from aryl, heteroaryl, cycloalkyl, and cycloalkenyl, wherein

20 R_{IX-2} is optionally substituted at a substitutable position with one or more radicals independently selected from alkyl, haloalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkoxy, halo, aryloxy, aralkyloxy, aryl, aralkyl, aminosulfonyl, amino, monoalkylamino and dialkylamino; and

wherein R_{IX-3} is selected from hydrido, -SH and halo;

25 provided R_{IX-2} cannot be phenyl or 4-methylphenyl when R_{IX-1} is higher alkyl and when R_{IX-3} is BSH.

Compounds of Formula IX and their methods of manufacture are disclosed in PCT Publication No. WO 9914204, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula IX:

- 2,4-dihydro-4-(3-methoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(2-fluorophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 5 2,4-dihydro-4-(2-methylphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(3-chlorophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(2-methoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(3-methylphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 4-cyclohexyl-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-thione;
 10 2,4-dihydro-4-(3-pyridyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(2-ethoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(2,6-dimethylphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(4-phenoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 4-(1,3-benzodioxol-5-yl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-
 15 thione;
 4-(2-chlorophenyl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(4-methoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-5-tridecyl-4-(3-trifluoromethylphenyl)-3H-1,2,4-triazole-3-
 thione;
 20 2,4-dihydro-5-tridecyl-4-(3-fluorophenyl)-3H-1,2,4-triazole-3-thione;
 4-(3-chloro-4-methylphenyl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-
 thione;
 2,4-dihydro-4-(2-methylthiophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 4-(4-benzyloxyphenyl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-thione;
 25 2,4-dihydro-4-(2-naphthyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-5-tridecyl-4-(4-trifluoromethylphenyl)-3H-1,2,4-triazole-3-
 thione;
 2,4-dihydro-4-(1-naphthyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(3-methylthiophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 30 2,4-dihydro-4-(4-methylthiophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
 2,4-dihydro-4-(3,4-dimethoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-
 thione;
 2,4-dihydro-4-(2,5-dimethoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-
 thione;

2,4-dihydro-4-(2-methoxy-5-chlorophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;


4-(4-aminosulfonylphenyl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-thione;

5 2,4-dihydro-5-dodecyl-4-(3-methoxyphenyl)-3H-1,2,4-triazole-3-thione;
2,4-dihydro-4-(3-methoxyphenyl)-5-tetradecyl-3H-1,2,4-triazole-3-thione;
2,4-dihydro-4-(3-methoxyphenyl)-5-undecyl-3H-1,2,4-triazole-3-thione;

and

2,4-dihydro-(4-methoxyphenyl)-5-pentadecyl-3H-1,2,4-triazole-3-thione.

10 Another class of CETP inhibitors that finds utility with the present invention consists of hetero-tetrahydroquinolines having the Formula X



Formula X

and pharmaceutically acceptable salts, enantiomers, or stereoisomers or N-oxides of said compounds;

15 in which

A_x represents cycloalkyl with 3 to 8 carbon atoms or a 5 to 7-membered, saturated, partially saturated or unsaturated, optionally benzo-condensed heterocyclic ring containing up to 3 heteroatoms from the series comprising S, N and/or O, that in case of a saturated heterocyclic ring is bonded to a nitrogen function, optionally bridged over it, and in which the aromatic systems mentioned above are optionally substituted up to 5-times in an identical or different substituents in the form of halogen, nitro, hydroxy, trifluoromethyl, trifluoromethoxy or by a straight-chain or branched alkyl, acyl, hydroxyalkyl or alkoxy each having up to 7 carbon atoms or by a group of the formula BNR_x.


20 3R_{x-4},

in which


R_{x-3} and R_{x-4} are identical or different and denote hydrogen, phenyl or straight-chain or branched alkyl having up to 6 carbon atoms, or

30

A_x represents a radical of the formula



5 D_x represents an aryl having 6 to 10 carbon atoms, that is optionally substituted by phenyl, nitro, halogen, trifluoromethyl or trifluoromethoxy, or it represents a radical of the formula



10 in which

R_{X-5} , R_{X-6} and R_{X-9} independently of one another denote cycloalkyl having 3 to 6 carbon atoms, or an aryl having 6 to 10 carbon atoms or a 5- to 7-membered aromatic, optionally benzo-condensed saturated or unsaturated, mono-, bi-, or tricyclic heterocyclic ring from the series consisting of S, N and/or O, in which the rings are substituted, optionally, in case of the nitrogen containing aromatic rings via the N function, with up to 5 identical or different substituents in the form of halogen, trifluoromethyl, nitro, hydroxy, cyano, carbonyl, trifluoromethoxy, straight straight-chain or branched acyl, alkyl, alkylthio, alkylalkoxy, alkoxy, or alkoxycarbonyl each having up to 6 carbon atoms, by aryl or trifluoromethyl-substituted aryl each having 6 to 10 carbon atoms or by an, optionally benzo-condensed, aromatic 5- to 7-membered heterocyclic ring having up to 3 heteroatoms from the series consisting of S, N, and/or O, and/or substituted by a group of the formula BOR_{X-10} , $-SR_{X-11}$, SO_2R_{X-12} or $BNR_{X-13}R_{X-14}$,

25 in which



R_{X-10} , R_{X-11} and R_{X-12} independently from each other denote aryl having 6 to 10 carbon atoms, which is in turn substituted with up to 2 identical or different

substituents in the form of phenyl, halogen or a straight-chain or branched alkyl having up to 6 carbon atoms,

R_{X-13} and R_{X-14} are identical or different and have the meaning of R_{X-3} and R_{X-4} indicated above,

5 or

R_{X-5} and/or R_{X-6} denote a radical of the formula



or

R_{X-7} denotes hydrogen or halogen, and

10 R_{X-8} denotes hydrogen, halogen, azido, trifluoromethyl, hydroxy, trifluoromethoxy, straight-chain or branched alkoxy or alkyl having up to 6 carbon atoms or a radical of the formula

$\text{BNR}_{X-15}\text{R}_{X-16}$,

in which

15 R_{X-15} and R_{X-16} are identical or different and have the meaning of R_{X-3} and R_{X-4} indicated above,

or

R_{X-7} and R_{X-8} together form a radical of the formula $=\text{O}$ or $=\text{NR}_{X-17}$,
in which

20 R_{X-17} denotes hydrogen or straight chain or branched alkyl, alkoxy or acyl having up to 6 carbon atoms,

L_X denotes a straight chain or branched alkylene or alkenylene chain having up to 8 carbon atoms, that are optionally substituted with up to 2 hydroxy groups,

25 T_X and X_X are identical or different and denote a straight chain or branched alkylene chain with up to 8 carbon atoms

or


T_X or X_X denotes a bond,

V_X represents an oxygen or sulfur atom or an BNR_{X-18} -group, in which

30 R_{X-18} denotes hydrogen or straight chain or branched alkyl with up to 6 carbon atoms or phenyl,

E_x represents cycloalkyl with 3 to 8 carbon atoms, or straight chain or branched alkyl with up to 8 carbon atoms, that is optionally substituted by cycloalkyl with 3 to 8 carbon atoms or hydroxy, or represents a phenyl, that is optionally substituted by halogen or trifluoromethyl,

- 5 R_{x-1} and R_{x-2} together form a straight-chain or branched alkylene chain with up to 7 carbon atoms, that must be substituted by carbonyl group and/or by a radical with the formula



- 10 in which a and b are identical or different and denote a number equaling 1,2, or 3,

- 15 R_{x-19} denotes hydrogen, cycloalkyl with 3 up to 7 carbon atoms, straight chain or branched silylalkyl with up to 8 carbon atoms or straight chain or branched alkyl with up to 8 carbon atoms, that are optionally substituted by hydroxyl, straight chain or branched alkoxy with up to 6 carbon atoms or by phenyl, which in turn might be substituted by halogen, nitro, trifluoromethyl, trifluoromethoxy or by phenyl or by tetrazole-substituted phenyl, and alkyl, optionally be substituted by a group with the formula BOR_{x-22} , in which



- 20 R_{x-22} denotes a straight chain or branched acyl with up to 4 carbon atoms or benzyl, or

- 25 R_{x-19} denotes straight chain or branched acyl with up to 20 carbon atoms or benzoyl, that is optionally substituted by halogen, trifluoromethyl, nitro or trifluoromethoxy, or it denotes straight chain or branched fluoroacyl with up to 8 carbon atoms and 9 fluorine atoms,

R_{x-20} and R_{x-21} are identical or different and denote hydrogen, phenyl or straight chain or branched alkyl with up to 6 carbon atoms, or

- 30 R_{x-20} and R_{x-21} together form a 3- to 6- membered carbocyclic ring, and the carbocyclic rings formed are optionally substituted, optionally also geminally, with up to six identical or different substituents in the form of trifluoromethyl, hydroxy, nitrile, halogen, carboxyl, nitro, azido, cyano, cycloalkyl or


cycloalkyloxy with 3 to 7 carbon atoms each, by straight chain or branched
 alkoxy, carbonyl, alkoxy or alkylthio with up to 6 carbon atoms each or by straight
 chain or branched alkyl with up to 6 carbon atoms, which in turn is substituted
 with up to 2 identically or differently by hydroxyl, benzyloxy, trifluoromethyl,
 5 benzoyl, straight chain or branched alkoxy, oxyacyl or carbonyl with up to 4
 carbon atoms each and/or phenyl, which may in turn be substituted with a
 halogen, trifluoromethyl or trifluoromethoxy, and/or the formed carbocyclic rings
 are optionally substituted, also geminally, with up to 5 identical or different
 substituents in the form of phenyl, benzoyl, thiophenyl or sulfonylbenzyl, which
 10 in turn are optionally substituted by halogen, trifluoromethyl, trifluoromethoxy or
 nitro, and/or optionally are substituted by a radical with the formula



in which

15 c denotes a number equaling 1, 2, 3, or 4,
 d denotes a number equaling 0 or 1,

R_{X-23} and R_{X-24} are identical or different and denote hydrogen, cycloalkyl
 with 3 to 6 carbon atoms, straight chain or branched alkyl with up to 6 carbon
 atoms, benzyl or phenyl, that is optionally substituted with up to 2 identically or
 20 differently by halogen, trifluoromethyl, cyano, phenyl or nitro, and/or the formed
 carbocyclic rings are substituted optionally by a spiro-linked radical with the
 formula



25 in which

W_X denotes either an oxygen or a sulfur atom

Y_X and Y'_X together form a 2 to 6 membered straight chain or branched
 alkylene chain,

e denotes a number equaling 1, 2, 3, 4, 5, 6, or 7,


30 f denotes a number equaling 1 or 2,

65

R_{X-25} , R_{X-26} , R_{X-27} , R_{X-28} , R_{X-29} , R_{X-30} and R_{X-31} are identical or different and denote hydrogen, trifluoromethyl, phenyl, halogen or straight chain or branched alkyl or alkoxy with up to 6 carbon atoms each,
or

5 R_{X-25} and R_{X-26} or R_{X-27} and R_{X-28} respectively form together a straight chain or branched alkyl chain with up to 6 carbon atoms,
or

R_{X-25} and R_{X-26} or R_{X-27} and R_{X-28} each together form a radical with the formula



10

in which

W_X has the meaning given above,

g denotes a number equaling 1, 2, 3, 4, 5, 6, or 7,

15 R_{X-32} and R_{X-33} form together a 3- to 7- membered heterocycle, which contains an oxygen or sulfur atom or a group with the formula SO , SO_2 or $-NR_{X-34}$,

in which

R_{X-34} denotes hydrogen, phenyl, benzyl or straight or branched alkyl with up to 4 carbon atoms.

20 Compounds of Formula X and their methods of manufacture are disclosed in PCT Publication No. WO 9914215, which is incorporated herein by reference in its entirety for all purposes.


In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula X:

25 2-cyclopentyl-5-hydroxy-7,7-dimethyl-4-(3-thienyl)-3-(4-trifluoromethylbenzoyl)-5,6,7,8-tetrahydroquinoline;

2-cyclopentyl-3-[fluoro-(4-trifluoromethylphenyl)methyl]-5-hydroxy-7,7-dimethyl-4-(3-thienyl)-5,6,7,8-tetrahydroquinoline; and

30 2-cyclopentyl-5-hydroxy-7,7-dimethyl-4-(3-thienyl)-3-(trifluoromethylbenzyl)-5,6,7,8-tetrahydroquinoline.


Another class of CETP inhibitors that finds utility with the present invention consists of substituted tetrahydro naphthalines and analogous compound having the Formula XI



Formula XI

and stereoisomers, stereoisomer mixtures, and salts thereof, in which


A_{XI} stands for cycloalkyl with 3 to 8 carbon atoms, or stands for aryl with 6 to 10 carbon atoms, or stands for a 5- to 7-membered, saturated, partially unsaturated or unsaturated, possibly benzocondensated, heterocycle with up to 4 heteroatoms from the series S, N and/or O, where aryl and the heterocyclic ring systems mentioned above are substituted up to 5-fold, identical or different, by cyano, halogen, nitro, carboxyl, hydroxy, trifluoromethyl, trifluoro- methoxy, or by straight-chain or branched alkyl, acyl, hydroxyalkyl, alkylthio, alkoxy, carbonyl, oxyalkoxy, carbonyl or alkoxy each with up to 7 carbon atoms, or by a group of the formula



in which

R_{XI-3} and R_{XI-4} are identical or different and denote hydrogen, phenyl, or straight-chain or branched alkyl with up to 6 carbon atoms

D_{XI} stands for a radical of the formula



in which

R_{XI-5}, R_{XI-6} and R_{XI-9}, independent of each other, denote cycloalkyl with 3 to 6 carbon atoms, or denote aryl with 6 to 10 carbon atoms, or denote a 5- to 7-membered, possibly benzocondensated, saturated or unsaturated, mono-, bi- or tricyclic heterocycle with up to 4 heteroatoms of the series S, N and/or O, where the cycles are possibly substituted in the case of the nitrogen-containing rings


also via the N-functionCup to 5-fold, identical or different, by halogen, trifluoromethyl, nitro, hydroxy, cyano, carboxyl, trifluoromethoxy, straight-chain or branched acyl, alkyl, alkylthio, alkylalkoxy, alkoxy or alkoxycarbonyl with up to 6 carbon atoms each, by aryl or trifluoromethyl substituted aryl with 6 to 10 carbon atoms each, or by a possibly benzocondensated aromatic 5- to 7-membered heterocycle with up to 3 heteroatoms of the series S, N and/or O, and/or are substituted by a group of the formula

5 $-OR_{XI-10}$, $-SR_{XI-11}$, $-SO_2R_{XI-12}$ or $-NR_{XI-13}R_{XI-14}$,
in which

10 R_{XI-10} , R_{XI-11} and R_{XI-12} , independent of each other, denote aryl with 6 to 10 carbon atoms, which itself is substituted up to 2-fold, identical or different, by phenyl, halogen, or by straight-chain or branched alkyl with up to 6 carbon atoms,

15 R_{XI-13} and R_{XI-14} are identical or different and have the meaning given above for R_{XI-3} and R_{XI-4} ,
or

R_{XI-5} and/or R_{XI-6} denote a radical of the formula



20 R_{XI-7} denotes hydrogen, halogen or methyl,
and

R_{XI-8} denotes hydrogen, halogen, azido, trifluoromethyl, hydroxy, trifluoromethoxy, straight-chain or branched alkoxy or alkyl with up to 6 carbon atoms each, or a radical of the formula $-NR_{XI-15}R_{XI-16}$,
in which

25 R_{XI-15} and R_{XI-16} are identical or different and have the meaning given above for R_{XI-3} and R_{XI-4} ,
or

R_{XI-7} and R_{XI-8} together form a radical of the formula $=O$ or $=NR_{XI-17}$, in which

30 R_{XI-17} denotes hydrogen or straight-chain or branched alkyl, alkoxy or acyl with up to 6 carbon atoms each,

L_{Xi} denotes a straight-chain or branched alkylene- or alkenylene chain with up to 8 carbon atoms each, which is possibly substituted up to 2-fold by hydroxy,

T_{Xi} and X_{Xi} are identical or different and denote a straight-chain or
5 branched alkylene chain with up to 8 carbon atoms,
or


T_{X1} and X_{X1} denotes a bond.

V_{X1} stands for an oxygen- or sulfur atom or for an $-NR_{X1-18}$ group,
in which

10 R_{XI-18} denotes hydrogen or straight-chain or branched alkyl with up to 6 carbon atoms, or phenyl.

15 E_{X1} stands for cycloalkyl with 3 to 8 carbon atoms, or stands for straight-chain or branched alkyl with up to 8 carbon atoms, which is possibly substituted by cycloalkyl with 3 to 8 carbon atoms or hydroxy, or stands for phenyl, which is possibly substituted by halogen or trifluoromethyl.

R_{X1-1} and R_{X1-2} together form a straight-chain or branched alkylene chain with up to 7 carbon atoms, which must be substituted by a carbonyl group and/or by a radical of the formula



20 in which

a and b are identical or different and denote a number 1, 2 or 3

25 R_{Xl-19} denotes hydrogen, cycloalkyl with 3 to 7 carbon atoms, straight-chain or branched silylalkyl with up to 8 carbon atoms, or straight-chain or branched alkyl with up to 8 carbon atoms, which is possibly substituted by hydroxy, straight-chain or branched alkoxy with up to 6 carbon atoms, or by phenyl, which itself can be substituted by halogen, nitro, trifluoromethyl, trifluoromethoxy or by phenyl substituted by phenyl or tetrazol, and alkyl is possibly substituted by a group of the formula $-OR_{Xl-22}$,



in which

30 R_{XI-22} denotes straight-chain or branched acyl with up to 4 carbon atoms,
or benzyl,
or

R_{XI-19} denotes straight-chain or branched acyl with up to 20 carbon atoms or benzoyl, which is possibly substituted by halogen, trifluoromethyl, nitro or trifluoromethoxy, or denotes straight-chain or branched fluoroacyl with up to 8 carbon atoms and 9 fluorine atoms,

5 R_{XI-20} and R_{XI-21} are identical or different, denoting hydrogen, phenyl or straight-chain or branched alkyl with up to 6 carbon atoms, or

R_{XI-20} and R_{XI-21} together form a 3- to 6-membered carbocycle, and, possibly also geminally, the alkylene chain formed by R_{XI-1} and R_{XI-2} , is possibly
 10 substituted up to 6-fold, identical or different, by trifluoromethyl, hydroxy, nitrile, halogen, carboxyl, nitro, azido, cyano, cycloalkyl or cycloalkyloxy with 3 to 7 carbon atoms each, by straight-chain or branched alkoxy carbonyl, alkoxy or alkoxythio with up to 6 carbon atoms each, or by straight-chain or branched alkyl with up to 6 carbon atoms, which itself is substituted up to 2-fold,
 15 identical or different, by hydroxyl, benzyloxy, trifluoromethyl, benzoyl, straight-chain or branched alkoxy, oxyacyl or carboxyl with up to 4 carbon atoms each, and/or phenyl- which itself can be substituted by halogen, trifluoromethyl or trifluoromethoxy, and/or the alkylene chain formed by R_{XI-1} and R_{XI-2} is substituted, also geminally, possibly up to 5-fold, identical or different, by
 20 phenyl, benzoyl, thiophenyl or sulfobenzyl -which themselves are possibly substituted by halogen, trifluoromethyl, trifluoromethoxy or nitro, and/or the alkylene chain formed by R_{XI-1} and R_{XI-2} is possibly substituted by a radical of the formula




in which

c denotes a number 1, 2, 3 or 4,

d denotes a number 0 or 1,

30 R_{XI-23} and R_{XI-24} are identical or different and denote hydrogen, cycloalkyl with 3 to 6 carbon atoms, straight-chain or branched alkyl with up to 6 carbon atoms, benzyl or phenyl, which is possibly substituted up to 2-fold, identical or different, by halogen, trifluoromethyl, cyano, phenyl or nitro, and/or the alkylene

chain formed by R_{X1-1} and R_{X1-2} is possibly substituted by a spiro-jointed radical of the formula



5 in which

W_{X1} denotes either an oxygen or a sulfur atom,

Y_{X1} and Y'_{X1} together form a 2- to 6-membered straight-chain or branched alkylene chain,

e is a number 1, 2, 3, 4, 5, 6 or 7,

10 f denotes a number 1 or 2,


R_{X1-25} , R_{X1-26} , R_{X1-27} , R_{X1-28} , R_{X1-29} , R_{X1-30} and R_{X1-31} are identical or different and denote hydrogen, trifluoromethyl, phenyl, halogen, or straight-chain or branched alkyl or alkoxy with up to 6 carbon atoms each,

or

15 R_{X1-25} and R_{X1-26} or R_{X1-27} and R_{X1-28} together form a straight-chain or branched alkyl chain with up to 6 carbon atoms,

or

R_{X1-25} and R_{X1-26} or R_{X1-27} and R_{X1-28} together form a radical of the formula



20 in which

W_{X1} has the meaning given above,

g is a number 1, 2, 3, 4, 5, 6 or 7,


R_{X1-32} and R_{X1-33} together form a 3- to 7-membered heterocycle that contains an oxygen- or sulfur atom or a group of the formula SO , SO_2 or $-NR_{X1-34}$,

25 in which

R_{X1-34} denotes hydrogen, phenyl, benzyl, or straight-chain or branched alkyl with up to 4 carbon atoms.

Compounds of Formula XI and their methods of manufacture are disclosed in PCT Publication No. WO 9914174, which is incorporated herein by reference in its entirety for all purposes.

Another class of CETP inhibitors that finds utility with the present invention
5 consists of 2-aryl-substituted pyridines having the Formula (XII)



Formula XII

or pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds,

10 in which


A_{XII} and E_{XII} are identical or different and stand for aryl with 6 to 10 carbon atoms which is possibly substituted, up to 5-fold identical or different, by halogen, hydroxy, trifluoromethyl, trifluoromethoxy, nitro or by straight-chain or branched alkyl, acyl, hydroxy alkyl or alkoxy with up to 7 carbon atoms each, or
15 by a group of the formula $-NR_{XII-1}R_{XII-2}$,
where

R_{XII-1} and R_{XII-2} are identical or different and are meant to be hydrogen, phenyl or straight-chain or branched alkyl with up to 6 carbon atoms,

D_{XII} stands for straight-chain or branched alkyl with up to 8 carbon
20 atoms, which is substituted by hydroxy,

L_{XII} stands for cycloalkyl with 3 to 8 carbon atoms or for straight-chain or branched alkyl with up to 8 carbon atoms, which is possibly substituted by cycloalkyl with 3 to 8 carbon atoms, or by hydroxy,

T_{XII} stands for a radical of the formula $R_{XII-3}-X_{XII}-$ or



25

where

R_{XII-3} and R_{XII-4} are identical or different and are meant to be cycloalkyl

with 3 to 8 carbon atoms, or aryl with 6 to 10 carbon atoms, or a 5- to 7-membered aromatic, possibly benzocondensated heterocycle with up to 3 heteroatoms from the series S, N and/or O, which are possibly substituted. up to 3-fold identical or different, by trifluoromethyl, trifluoromethoxy, halogen, hydroxy, carboxyl, nitro, by straight-chain or branched alkyl, acyl, alkoxy or alkoxy carbonyl with up to 6 carbon atoms each. or by phenyl, phenoxy or phenylthio which in turn can be substituted by halogen. trifluoromethyl or trifluoromethoxy, and/or where the cycles are possibly substituted by a group of the formula $-NR_{XII-7}R_{XII-8}$,

10 where

R_{XII-7} and R_{XII-8} are identical or different and have the meaning of R_{XII-1} and R_{XII-2} given above,

X_{XII} is a straight-chain or branched alkyl or alkenyl with 2 to 10 carbon atoms each, possibly substituted up to 2-fold by hydroxy or halogen,

15 R_{XII-5} stands for hydrogen.

and

R_{XII-6} means to be hydrogen, halogen, mercapto, azido, trifluoromethyl, hydroxy, trifluoromethoxy, straight-chain or branched alkoxy with up to 5 carbon atoms, or a radical of the formula $BNR_{XII-9}R_{XII-10}$,

20 where

R_{XII-9} and R_{XII-10} are identical or different and have the meaning of R_{XII-1} and R_{XII-2} given above,

or

R_{XII-5} and R_{XII-6} , together with the carbon atom, form a carbonyl group.

25 Compounds of Formula XII and their methods of manufacture are disclosed in EP 796846-A1, United States Patent No. 6,127,383 and United States Patent No. 5,925,645, all of which are incorporated herein by reference in their entireties for all purposes.

30 In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula XII:

4,6-bis-(p-fluorophenyl)-2-isopropyl-3-[(p-trifluoromethylphenyl)-(fluoro)-methyl]-5-(1-hydroxyethyl)pyridine;


2,4-bis-(4-fluorophenyl)-6-isopropyl-5-[4-(trifluoromethylphenyl)-fluoromethyl]-3-hydroxymethylpyridine; and

35 2,4-bis-(4-fluorophenyl)-6-isopropyl-5-[2-(3-trifluoromethylphenyl)vinyl]-

3-hydroxymethyl)pyridine.

Another class of CETP inhibitors that finds utility with the present invention consists of compounds having the Formula (XIII)

5



Formula XIII

or pharmaceutically acceptable salts, enantiomers, stereoisomers, hydrates, or solvates of said compounds, in which

10 R_{XIII} is a straight chain or branched C_{1-10} alkyl; straight chain or branched C_{2-10} alkenyl; halogenated C_{1-4} lower alkyl; C_{3-10} cycloalkyl that may be substituted; C_{5-8} cycloalkenyl that may be substituted; C_{3-10} cycloalkyl C_{1-10} alkyl that may be substituted; aryl that may be substituted; aralkyl that may be substituted; or a 5- or 6-membered heterocyclic group having 1 to 3 nitrogen
15 atoms, oxygen atoms or sulfur atoms that may be substituted,

X_{XIII-1} , X_{XIII-2} , X_{XIII-3} , X_{XIII-4} may be the same or different and are a hydrogen atom; halogen atom; C_{1-4} lower alkyl; halogenated C_{1-4} lower alkyl; C_{1-4} lower alkoxy; cyano group; nitro group; acyl; or aryl, respectively;

Y_{XIII} is $-CO-$; or BSO_2- ; and

20 Z_{XIII} is a hydrogen atom; or mercapto protective group.

Compounds of Formula XIII and their methods of manufacture are disclosed in PCT Publication No. WO 98/35937, which is incorporated herein by reference in its entirety for all purposes.

25 In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula XIII:

N,N'-(dithiodi-2,1-phenylene)bis[2,2-dimethyl-propanamide];

N,N'-(dithiodi-2,1-phenylene)bis[1-methyl-cyclohexanecarboxamide];

N,N'-(dithiodi-2,1-phenylene)bis[1-(3-methylbutyl)-

cyclopentanecarboxamide];

N,N'-(dithiodi-2,1-phenylene)bis[1-(3-methylbutyl)-
cyclohexanecarboxamide];

5 N,N'-(dithiodi-2,1-phenylene)bis[1-(2-ethylbutyl)-
cyclohexanecarboxamide];


N,N'-(dithiodi-2,1-phenylene)bis-tricyclo[3.3.1.^{3,7}]decane-1-
carboxamide;

propanethioic acid, 2-methyl-, S-[2[[[1-(2-
ethylbutyl)cyclohexyl]carbonyl]amino]phenyl] ester;

10 propanethioic acid, 2,2-dimethyl-, S-[2[[[1-(2-
ethylbutyl)cyclohexyl]carbonyl]amino]phenyl] ester; and

ethanethioic acid, S-[2[[[1-(2-
ethylbutyl)cyclohexyl]carbonyl]amino]phenyl] ester.

15 Another class of CETP inhibitors that finds utility with the present
invention consists of polycyclic aryl and heteroaryl tertiary-heteroalkylamines
having the Formula XIV



Formula XIV

and pharmaceutically acceptable forms thereof, wherein:

n_{XIV} is an integer selected from 0 through 5;

R_{XIV-1} is selected from the group consisting of haloalkyl, haloalkenyl, haloalkoxyalkyl, and haloalkenyloxyalkyl;

X_{XIV} is selected from the group consisting of O, H, F, S, S(O), NH, N(OH), N(alkyl), and N(alkoxy);

R_{XIV-16} is selected from the group consisting of hydrido, alkyl, alkenyl, alkynyl, aryl, aralkyl, aryloxyalkyl, alkoxyalkyl, alkenyloxyalkyl, alkylthioalkyl, arylthioalkyl, aralkoxyalkyl, heteroaralkoxyalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl, cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, monocarboalkoxyalkyl, monocarboalkoxy, dicarboalkoxyalkyl, monocarboxamido, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, acyl, aroyl, heteroaroyl, heteroaryloxyalkyl, dialkoxyphosphonoalkyl, trialkylsilyl, and a spacer selected from the group consisting of a covalent single bond and a linear spacer moiety having from 1 through 4 contiguous atoms linked to the point of bonding of an aromatic substituent selected from the group consisting of R_{XIV-4} , R_{XIV-8} , R_{XIV-9} , and R_{XIV-13} to form a heterocyclyl ring having from 5 through 10 contiguous members with the provisos that said spacer moiety is other than a covalent single bond when R_{XIV-2} is alkyl and there is no R_{XIV-16} wherein X is H or F;

D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} are independently selected from the group consisting of C, N, O, S and a covalent bond with the provisos that no more than one of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} is a covalent bond, no more than one of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} is O, no more than one of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} is S, one of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} must be a covalent bond when two of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} are O and S, and no more than four of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} are N;

D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} are independently selected from the group consisting of C, N, O, S and a covalent bond with the provisos that no more than one of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} is a covalent bond, no more than one of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} is O, no more than one of D_{XIV-3} ,

D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} is S, one of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} must be a covalent bond when two of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} are O and S, and no more than four of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} and K_{XIV-2} are N;

R_{XIV-2} is independently selected from the group consisting of hydrido,
 5 hydroxy, hydroxyalkyl, amino, aminoalkyl, alkylamino, dialkylamino, alkyl,
 alkenyl, alkynyl, aryl, aralkyl, aralkoxyalkyl, aryloxyalkyl, alkoxyalkyl,
 heteroaryloxyalkyl, alkenyloxyalkyl, alkylthioalkyl, aralkylthioalkyl, arylthioalkyl,
 cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl, cycloalkenylalkyl,
 haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy,
 10 aloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy,
 halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl,
 perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl,
 heteroaralkylthioalkyl, monocarboalkoxyalkyl, dicarboalkoxyalkyl,
 monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl,
 15 alkylsulfonyl, alkylsulfinylalkyl, alkylsulfonylalkyl, haloalkylsulfinyl,
 haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl,
 aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl,
 cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylsulfonylalkyl,
 heteroarylsulfinyl, heteroarylsulfonyl, heteroarylsulfinylalkyl, aralkylsulfinylalkyl,
 20 aralkylsulfonylalkyl, carboxy, carboxyalkyl, carboalkoxy, carboxamide,
 carboxamidoalkyl, carboaralkoxy, dialkoxyphosphono, diaralkoxyphosphono,
 dialkoxyphosphonoalkyl, and diaralkoxyphosphonoalkyl;

R_{XIV-2} and R_{XIV-3} are taken together to form a linear spacer moiety
 25 selected from the group consisting of a covalent single bond and a moiety
 having from 1 through 6 contiguous atoms to form a ring selected from the
 group consisting of a cycloalkyl having from 3 through 8 contiguous members, a
 cycloalkenyl having from 5 through 8 contiguous members, and a heterocyclyl
 having from 4 through 8 contiguous members;

R_{XIV-3} is selected from the group consisting of hydrido, hydroxy, halo,
 30 cyano, aryloxy, hydroxyalkyl, amino, alkylamino, dialkylamino, acyl, sulfhydryl,
 acylamido, alkoxy, alkylthio, arylthio, alkyl, alkenyl, alkynyl, aryl,
 aralkyl, aryloxyalkyl, alkoxyalkyl, heteroarylthio, aralkylthio, aralkoxyalkyl,
 alkylsulfinylalkyl, alkylsulfonylalkyl, aroyl, heteroaroyl, aralkylthioalkyl,
 heteroaralkylthioalkyl, heteroaryloxyalkyl, alkenyloxyalkyl, alkylthioalkyl,
 35 arylthioalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl,

cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl, 5 monocarboalkoxyalkyl, dicarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl, aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl, cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylulfonylalkyl, 10 heteroarylulfonyl, heteroarylulfonyl, heteroarylulfonylalkyl, aralkylulfonylalkyl, aralkylulfonylalkyl, carboxy, carboxyalkyl, carboalkoxy, carboxamide, carboxamidoalkyl, carboaralkoxy, dialkoxyposphono, diaralkoxyposphono, dialkoxyposphonoalkyl, and diaralkoxyposphonoalkyl;

Y_{XIV} is selected from a group consisting of a covalent single bond, $(C(R_{XIV-14})_2)_{q_{XIV}}$ wherein q_{XIV} is an integer selected from 1 and 2 and $(CH(R_{XIV-14}))_{g_{XIV}}-W_{XIV}-(CH(R_{XIV-14}))_{p_{XIV}}$ wherein g_{XIV} and p_{XIV} are integers 15 independently selected from 0 and 1;

R_{XIV-14} is independently selected from the group consisting of hydrido, hydroxy, halo, cyano, aryloxy, amino, alkylamino, dialkylamino, hydroxyalkyl, 20 acyl, aroyl, heteroaroyl, heteroaryloxyalkyl, sulfhydryl, acylamido, alkoxy, alkylthio, arylthio, alkyl, alkenyl, alkynyl, aryl, aralkyl, aryloxyalkyl, aralkoxyalkylalkoxy, alkylsulfinylalkyl, alkylsulfonylalkyl, aralkylthioalkyl, heteroaralkoxythioalkyl, alkoxyalkyl, heteroaryloxyalkyl, alkenyloxyalkyl, alkylthioalkyl, arylthioalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, 25 cycloalkenyl, cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl, heteroaralkylthioalkyl, monocarboalkoxyalkyl, 30 dicarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl, aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl, cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylulfonylalkyl, heteroarylulfonyl, 35 heteroarylulfonyl, heteroarylulfonylalkyl, aralkylulfonylalkyl,

aralkylsulfonylalkyl, carboxy, carboxyalkyl, carboalkoxy, carboxamide, carboxamidoalkyl, carboaralkoxy, dialkoxyphosphono, diaralkoxyphosphono, dialkoxyphosphonoalkyl, diaralkoxyphosphonoalkyl, a spacer selected from a moiety having a chain length of 3 to 6 atoms connected to the point of bonding
 5 selected from the group consisting of R_{XIV-9} and R_{XIV-13} to form a ring selected from the group consisting of a cycloalkenyl ring having from 5 through 8 contiguous members and a heterocyclyl ring having from 5 through 8 contiguous members and a spacer selected from a moiety having a chain length of 2 to 5 atoms connected to the point of bonding selected from the group
 10 consisting of R_{XIV-4} and R_{XIV-8} to form a heterocyclyl having from 5 through 8 contiguous members with the proviso that, when Y_{XIV} is a covalent bond, an R_{XIV-14} substituent is not attached to Y_{XIV} ;

R_{XIV-14} and R_{XIV-14} , when bonded to the different atoms, are taken together to form a group selected from the group consisting of a covalent bond;
 15 alkylene, haloalkylene, and a spacer selected from a group consisting of a moiety having a chain length of 2 to 5 atoms connected to form a ring selected from the group of a saturated cycloalkyl having from 5 through 8 contiguous members, a cycloalkenyl having from 5 through 8 contiguous members, and a heterocyclyl having from 5 through 8 contiguous members;

20 R_{XIV-14} and R_{XIV-14} , when bonded to the same atom are taken together to form a group selected from the group consisting of oxo, thiono, alkylene, haloalkylene, and a spacer selected from the group consisting of a moiety having a chain length of 3 to 7 atoms connected to form a ring selected from the group consisting of a cycloalkyl having from 4 through 8 contiguous members, a
 25 cycloalkenyl having from 4 through 8 contiguous members, and a heterocyclyl having from 4 through 8 contiguous members;

W_{XIV} is selected from the group consisting of O, C(O), C(S), C(O) $N(R_{XIV-14})$, C(S) $N(R_{XIV-14})$, $(R_{XIV-14})NC(O)$, $(R_{XIV-14})NC(S)$, S, S(O), S(O)₂, S(O)₂ $N(R_{XIV-14})$, $(R_{XIV-14})NS(O)_2$, and $N(R_{XIV-14})$ with the proviso that R_{XIV-14} is selected from other
 30 than halo and cyano;

Z_{XIV} is independently selected from a group consisting of a covalent single bond, $(C(R_{XIV-15})_2)_{q_{XIV-2}}$ wherein q_{XIV-2} is an integer selected from 1 and 2, $(CH(R_{XIV-15}))_{j_{XIV}}-W-(CH(R_{XIV-15}))_{k_{XIV}}$ wherein j_{XIV} and k_{XIV} are integers independently selected from 0 and 1 with the proviso that, when Z_{XIV} is
 35 a covalent single bond, an R_{XIV-15} substituent is not attached to Z_{XIV} ;

R_{XIV-15} is independently selected, when Z_{XIV} is $(C(R_{XIV-15})_2)_{q_{XIV}}$ wherein q_{XIV} is an integer selected from 1 and 2, from the group consisting of hydrido, hydroxy, halo, cyano, aryloxy, amino, alkylamino, dialkylamino, hydroxyalkyl, acyl, aroyl, heteroaroyl, heteroaryloxyalkyl, sulfhydryl, acylamido, alkoxy, alkylthio, arylthio, alkyl, alkenyl, alkynyl, aryl, aralkyl, aryloxyalkyl, aralkoxyalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, aralkylthioalkyl, heteroaralkylthioalkyl, alkoxyalkyl, heteroaryloxyalkyl, alkenyloxyalkyl, alkylthioalkyl, arylthioalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl, cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl, heteroaralkylthioalkyl, monocarboalkoxyalkyl, dicarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl, aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl, cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylulfonylalkyl, heteroarylulfonyl, heteroarylulfonylalkyl, aralkylsulfinylalkyl, aralkylsulfonylalkyl, carboxy, carboxyalkyl, carboalkoxy, carboxamide, carboxamidoalkyl, carboaralkoxy, dialkoxyposphono, diaralkoxyposphono, dialkoxyposphonoalkyl, diaralkoxyposphonoalkyl, a spacer selected from a moiety having a chain length of 3 to 6 atoms connected to the point of bonding selected from the group consisting of R_{XIV-4} and R_{XIV-8} to form a ring selected from the group consisting of a cycloalkenyl ring having from 5 through 8 contiguous members and a heterocyclyl ring having from 5 through 8 contiguous members, and a spacer selected from a moiety having a chain length of 2 to 5 atoms connected to the point of bonding selected from the group consisting of R_{XIV-9} and R_{XIV-13} to form a heterocyclyl having from 5 through 8 contiguous members; R_{XIV-15} and R_{XIV-15} , when bonded to the different atoms, are taken together to form a group selected from the group consisting of a covalent bond, alkylene, haloalkylene, and a spacer selected from a group consisting of a moiety having a chain length of 2 to 5 atoms connected to form a ring selected from the group of a saturated cycloalkyl having from 5 through 8 contiguous

members, a cycloalkenyl having from 5 through 8 contiguous members, and a heterocyclyl having from 5 through 8 contiguous members;

R_{XIV-15} and R_{XIV-15} , when bonded to the same atom are taken together to form a group selected from the group consisting of oxo, thiono, alkylene, haloalkylene, and a spacer selected from the group consisting of a moiety having a chain length of 3 to 7 atoms connected to form a ring selected from the group consisting of a cycloalkyl having from 4 through 8 contiguous members, a cycloalkenyl having from 4 through 8 contiguous members, and a heterocyclyl having from 4 through 8 contiguous members;

R_{XIV-15} is independently selected, when Z_{XIV} is $(CH(R_{XIV-15}))_{j_{XIV}}-W-(CH(R_{XIV-15}))_{k_{XIV}}$ wherein j_{XIV} and k_{XIV} are integers independently selected from 0 and 1, from the group consisting of hydrido, halo, cyano, aryloxy, carboxyl, acyl, aroyl, heteroaroyl, hydroxyalkyl, heteroaryloxyalkyl, acylamido, alkoxy, alkylthio, arylthio, alkyl, alkenyl, alkynyl, aryl, aralkyl, aryloxyalkyl, alkoxyalkyl, heteroaryloxyalkyl, aralkoxyalkyl, heteroaralkoxyalkyl, alkylsulfonylalkyl, alkylsulfinylalkyl, alkenyloxyalkyl, alkylthioalkyl, arylthioalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl, cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl, heteroaralkylthioalkyl, monocarboalkoxyalkyl, dicarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl, aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl, cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylsulfonylalkyl, heteroarylsulfinyl, heteroarylsulfonyl, heteroarylsulfinylalkyl, aralkylsulfinylalkyl, aralkylsulfonylalkyl, carboxyalkyl, carboalkoxy, carboxamide, carboxamidoalkyl, carboaralkoxy, dialkoxyposphonoalkyl, diaralkoxyposphonoalkyl, a spacer selected from a linear moiety having a chain length of 3 to 6 atoms connected to the point of bonding selected from the group consisting of R_{XIV-4} and R_{XIV-8} to form a ring selected from the group consisting of a cycloalkenyl ring having from 5 through 8 contiguous members and a heterocyclyl ring having from 5 through 8 contiguous members, and a spacer

selected from a linear moiety having a chain length of 2 to 5 atoms connected to the point of bonding selected from the group consisting of R_{XIV-9} and R_{XIV-13} to form a heterocyclyl ring having from 5 through 8 contiguous members;

- R_{XIV-4} , R_{XIV-5} , R_{XIV-6} , R_{XIV-7} , R_{XIV-8} , R_{XIV-9} , R_{XIV-10} , R_{XIV-11} , R_{XIV-12} , and R_{XIV-13}
- 5 are independently selected from the group consisting of perhaloaryloxy, alkanoylalkyl, alkanoylalkoxy, alkanoyloxy, N-aryl-N-alkylamino, heterocyclylalkoxy, heterocyclylthio, hydroxyalkoxy, carboxamidoalkoxy, alkoxy carbonylalkoxy, alkoxy carbonylalkenyloxy, aralkanoylalkoxy, aralkenoyl, N-alkylcarboxamido, N-haloalkylcarboxamido, N-cycloalkylcarboxamido,
- 10 N-arylcarboxamidoalkoxy, cycloalkylcarbonyl, cyanoalkoxy, heterocyclylcarbonyl, hydrido, carboxy, heteroaralkylthio, heteroaralkoxy, cycloalkylamino, acylalkyl, acylalkoxy, aroylalkoxy, heterocycliloxy, aralkylaryl, aralkyl, aralkenyl, aralkynyl, heterocyclyl, perhaloaralkyl, aralkylsulfonyl, aralkylsulfonylalkyl, aralkylsulfinyl, aralkylsulfinylalkyl, halocycloalkyl,
- 15 halocycloalkenyl, cycloalkylsulfinyl, cycloalkylsulfinylalkyl, cycloalkylsulfonyl, cycloalkylsulfonylalkyl, heteroaryl amino, N-heteroaryl amino-N-alkyl amino, heteroaryl aminoalkyl, haloalkylthio, alkanoyloxy, alkoxy, alkoxyalkyl, haloalkoxyalkyl, heteroaralkoxy, cycloalkoxy, cycloalkenyloxy, cycloalkoxyalkyl, cycloalkylalkoxy, cycloalkenyloxyalkyl, cycloalkylenedioxy, halocycloalkoxy,
- 20 halocycloalkoxyalkyl, halocycloalkenyloxy, halocycloalkenyloxyalkyl, hydroxy, amino, thio, nitro, lower alkyl amino, alkylthio, alkylthioalkyl, aryl amino, aralkyl amino, arylthio, arylthioalkyl, heteroaralkoxyalkyl, alkylsulfinyl, alkylsulfinylalkyl, arylsulfinylalkyl, arylsulfonylalkyl, heteroarylsulfinylalkyl, heteroarylsulfonylalkyl, alkylsulfonyl, alkylsulfonylalkyl, haloalkylsulfinylalkyl,
- 25 haloalkylsulfonylalkyl, alkylsulfonamido, alkylaminosulfonyl, amidosulfonyl, monoalkylamidossulfonyl, dialkyl amidosulfonyl, monoarylamidosulfonyl, arylsulfonamido, diarylamidosulfonyl, monoalkyl monoaryl amidosulfonyl, arylsulfinyl, arylsulfonyl, heteroarylthio, heteroarylsulfinyl, heteroarylsulfonyl, heterocyclylsulfonyl, heterocyclylthio, alkanoyl, alkenoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, haloalkanoyl, alkyl, alkenyl, alkynyl, alkenyloxy,
- 30 alkenyloxyalkyl, alkylenedioxy, haloalkylenedioxy, cycloalkyl, cycloalkylalkanoyl, cycloalkenyl, lower cycloalkylalkyl, lower cycloalkenylalkyl, halo, haloalkyl; haloalkenyl, haloalkoxy, hydroxyhaloalkyl, hydroxyaralkyl, hydroxyalkyl, hydroxyheteroaralkyl, haloalkoxyalkyl, aryl, heteroaralkynyl, aryloxy, aralkoxy,
- 35 aryloxyalkyl, saturated heterocyclyl, partially saturated heterocyclyl, heteroaryl,

heteroaryloxy, heteroaryloxyalkyl, arylalkenyl, heteroarylalkenyl, carboxyalkyl, carboalkoxy, alkoxycarboxamido, alkylamidocarbonylamido, arylamidocarbonylamido, carboalkoxyalkyl, carboalkoxyalkenyl, carboaralkoxy, carboxamido, carboxamidoalkyl, cyano, carbohaloalkoxy, phosphono, phosphonoalkyl, diaralkoxyphosphono, and diaralkoxyphosphonoalkyl with the proviso that there are one to five non-hydrido ring substituents R_{XIV-4} , R_{XIV-5} , R_{XIV-6} , R_{XIV-7} , and R_{XIV-8} present, that there are one to five non-hydrido ring substituents R_{XIV-9} , R_{XIV-10} , R_{XIV-11} , R_{XIV-12} , and R_{XIV-13} present, and R_{XIV-4} , R_{XIV-5} , R_{XIV-6} , R_{XIV-7} , R_{XIV-8} , R_{XIV-9} , R_{XIV-10} , R_{XIV-11} , R_{XIV-12} , and R_{XIV-13} are each independently selected to maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen;

R_{XIV-4} and R_{XIV-5} , R_{XIV-5} and R_{XIV-6} , R_{XIV-6} and R_{XIV-7} , R_{XIV-7} and R_{XIV-8} , R_{XIV-8} and R_{XIV-9} , R_{XIV-9} and R_{XIV-10} , R_{XIV-10} and R_{XIV-11} , R_{XIV-11} and R_{XIV-12} , and R_{XIV-12} and R_{XIV-13} are independently selected to form spacer pairs wherein a spacer pair is taken together to form a linear moiety having from 3 through 6 contiguous atoms connecting the points of bonding of said spacer pair members to form a ring selected from the group consisting of a cycloalkenyl ring having 5 through 8 contiguous members, a partially saturated heterocyclyl ring having 5 through 8 contiguous members, a heteroaryl ring having 5 through 6 contiguous members, and an aryl with the provisos that no more than one of the group consisting of spacer pairs R_{XIV-4} and R_{XIV-5} , R_{XIV-5} and R_{XIV-6} , R_{XIV-6} and R_{XIV-7} , and R_{XIV-7} and R_{XIV-8} are used at the same time and that no more than one of the group consisting of spacer pairs R_{XIV-9} and R_{XIV-10} , R_{XIV-10} and R_{XIV-11} , R_{XIV-11} and R_{XIV-12} , and R_{XIV-12} and R_{XIV-13} are used at the same time;

R_{XIV-4} and R_{XIV-9} , R_{XIV-4} and R_{XIV-13} , R_{XIV-8} and R_{XIV-9} , and R_{XIV-8} and R_{XIV-13} are independently selected to form a spacer pair wherein said spacer pair is taken together to form a linear moiety wherein said linear moiety forms a ring selected from the group consisting of a partially saturated heterocyclyl ring having from 5 through 8 contiguous members and a heteroaryl ring having from 5 through 6 contiguous members with the proviso that no more than one of the group consisting of spacer pairs R_{XIV-4} and R_{XIV-9} , R_{XIV-4} and R_{XIV-13} , R_{XIV-8} and R_{XIV-9} , and R_{XIV-8} and R_{XIV-13} is used at the same time.

Compounds of Formula XIV and their methods of manufacture are disclosed in PCT Publication No. WO 00/18721, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula XIV:

- 3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-isopropylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-cyclopropylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-(2-furyl)phenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2,3-dichlorophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-methylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3,5-dimethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-ethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-*t*-butylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(3-methylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

5 3-[[3-(phenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

10 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethoxy)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethyl)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3,5-dimethylphenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

15 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethylthio)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3,5-difluorophenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

20 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[cyclohexylmethoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

25 3-[[3-(3-difluoromethoxyphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

30 3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(pentafluoroethyl)methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(3-isopropylphenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[3-(3-cyclopropylphenoxy)phenyl][[3-(pentafluoroethyl) phenyl]methyl]-
amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-(2-furyl)phenoxy)phenyl][[3-(pentafluoroethyl) phenyl]methyl]-
amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[3-(2,3-dichlorophenoxy)phenyl][[3-(pentafluoroethyl) phenyl]methyl]-
amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-methylphenoxy)phenyl][[3-(pentafluoroethyl)
10 phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[3-
(pentafluoroethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[3-(pentafluoroethyl)phenyl]-
methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3,5-dimethylphenoxy)phenyl][[3-(pentafluoroethyl) phenyl]methyl]-
20 amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-ethylphenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-butylphenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[3-(3-methylphenoxy)phenyl][[3-pentafluoroethyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[3-
(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(phenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]-
30 amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[3-
(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[[3-(trifluoromethoxy)phenyl]-
methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[[3-(trifluoromethyl)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[[3-(trifluoromethylthio)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[[3,5-difluorophenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[cyclohexylmethoxy]phenyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-(3-difluoromethoxyphenoxy)phenyl][[3-(pentafluoroethyl) phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[3-(pentafluoroethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-isopropylphenoxy)phenyl][[3-(heptafluoropropyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[[3-(3-cyclopropylphenoxy)phenyl][[3-(heptafluoropropyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-(2-furyl)phenoxy)phenyl][[3-(heptafluoropropyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(2,3-dichlorophenoxy)phenyl][[3-(heptafluoropropyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[[3-(4-fluorophenoxy)phenyl][[3-(heptafluoropropyl) phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(4-methylphenoxy)phenyl][[3-(heptafluoropropyl) phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

MISSING AT THE TIME OF PUBLICATION

- (heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl]][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(3-difluoromethoxyphenoxy)phenyl]][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
5 3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl]][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl]][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl]][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
10 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(3-isopropylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(3-cyclopropylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
15 3-[[3-(3-(2-furyl)phenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(2,3-dichlorophenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
20 3-[[3-(4-fluorophenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(4-methylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
25 3-[[3-(2-fluoro-5-bromophenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(4-chloro-3-ethylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
30 3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(3,5-dimethylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;


- 3-[[3-(3-ethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-t-butylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[3-(3-methylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(phenoxy)phenyl][[2-fluoro-5-(trifluoromethyl) phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethoxy)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethyl)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-(cyclohexylmethoxy)-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-difluoromethoxyphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-isopropylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[3-(3-cyclopropylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-(2-furyl)phenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2,3-dichlorophenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[3-(4-fluorophenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-methylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3,5-dimethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[3-(3-ethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-t-butylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[3-(3-methylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(phenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 35

91

- 3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethoxy)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethyl)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[cyclohexylmethoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-difluoromethoxyphenoxy)phenyl][2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol; and
- 3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][2-fluoro-4-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol.

25 Another class of CETP inhibitors that finds utility with the present invention consists of substituted N-Aliphatic-N-Aromatic *tertiary*-Heteroalkylamines having the Formula XV




Formula XV


and pharmaceutically acceptable forms thereof, wherein:

5 n_{xv} is an integer selected from 1 through 2;

AQ-1



AQ-2



A_{xv} and Q_{xv} are independently selected from the group consisting of
 $-\text{CH}_2(\text{CR}_{xv-37}\text{R}_{xv-38})_{v_{xv}}-(\text{CR}_{xv-33}\text{R}_{xv-34})_{u_{xv}}-\text{T}_{xv}-(\text{CR}_{xv-35}\text{R}_{xv-36})_{w_{xv}}-\text{H}$,

with the provisos that one of A_{XV} and Q_{XV} must be AQ-1 and that one of A_{XV} and Q_{XV} must be selected from the group consisting of AQ-2 and $-\text{CH}_2(\text{CR}_{XV-37}\text{R}_{XV-38})_{vXV}-(\text{CR}_{XV-33}\text{R}_{XV-34})_{uXV}-\text{T}_{XV}-(\text{CR}_{XV-35}\text{R}_{XV-36})_{wXV}-\text{H}$;

T_{XV} is selected from the group consisting of a single covalent bond, O, S, S(O), S(O)₂, C(R_{XV-33})=C(R_{XV-35}), and

$\text{C} \equiv \text{C}$;

vXV is an integer selected from 0 through 1 with the proviso that vXV is 1 when any one of R_{XV-33} , R_{XV-34} , R_{XV-35} , and R_{XV-36} is aryl or heteroaryl;

uXV and wXV are integers independently selected from 0 through 6;

A_{XV-1} is C(R_{XV-30});

D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} are independently selected from the group consisting of C, N, O, S and a covalent bond with the provisos that no more than one of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} is a covalent bond, no more than one of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} is O, no more than one of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} is S, one of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} must be a covalent bond when two of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} are O and S, and no more than four of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} are N;

B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are independently selected from the group consisting of C, C(R_{XV-30}), N, O, S and a covalent bond with the provisos that no more than 5 of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are a covalent bond, no more than two of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are O, no more than two of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are S, no more than two of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are simultaneously O and S, and no more than two of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are N;

B_{XV-1} and D_{XV-3} , D_{XV-3} and J_{XV-3} , J_{XV-3} and K_{XV-2} , K_{XV-2} and J_{XV-4} , J_{XV-4} and D_{XV-4} , and D_{XV-4} and B_{XV-2} are independently selected to form an in-ring spacer pair wherein said

spacer pair is selected from the group consisting of C(R_{XV-33})=C(R_{XV-35}) and N=N with the provisos that AQ-2 must be a ring of at least five contiguous members, that no more than two of the group of said spacer pairs are simultaneously C(R_{XV-33})=C(R_{XV-35}) and that no more than one of the group of said spacer pairs can be N=N unless the other spacer pairs are other than C(R_{XV-33})=C(R_{XV-35}), O, N, and S;

R_{XV-1} is selected from the group consisting of haloalkyl and haloalkoxymethyl;

R_{XV-2} is selected from the group consisting of hydrido, aryl, alkyl, alkenyl, haloalkyl, haloalkoxy, haloalkoxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl and heteroaryl;

R_{XV-3} is selected from the group consisting of hydrido, aryl, alkyl, alkenyl, haloalkyl, and haloalkoxyalkyl;

Y_{XV} is selected from the group consisting of a covalent single bond, $(CH_2)_q$ wherein q is an integer selected from 1 through 2 and $(CH_2)_j-O-(CH_2)_k$ wherein j and k are integers independently selected from 0 through 1;

Z_{XV} is selected from the group consisting of covalent single bond, $(CH_2)_q$ wherein q is an integer selected from 1 through 2, and $(CH_2)_j-O-(CH_2)_k$ wherein j and k are integers independently selected from 0 through 1;

R_{XV-4} , R_{XV-8} , R_{XV-9} and R_{XV-13} are independently selected from the group consisting of hydrido, halo, haloalkyl, and alkyl;

R_{XV-30} is selected from the group consisting of hydrido, alkoxy, alkoxyalkyl, halo, haloalkyl, alkylamino, alkylthio, alkylthioalkyl, alkyl, alkenyl, haloalkoxy, and haloalkoxyalkyl with the proviso that R_{XV-30} is selected to maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen;

R_{XV-30} , when bonded to A_{XV-1} , is taken together to form an intra-ring linear spacer connecting the A_{XV-1} -carbon at the point of attachment of R_{XV-30} to the point of bonding of a group selected from the group consisting of R_{XV-10} , R_{XV-11} , R_{XV-12} , R_{XV-31} , and R_{XV-32} wherein said intra-ring linear spacer is selected from the group consisting of a covalent single bond and a spacer moiety having from 1 through 6 contiguous atoms to form a ring selected from the group consisting of a cycloalkyl having from 3 through 10 contiguous members, a cycloalkenyl having from 5 through 10 contiguous members, and a heterocyclyl having from 5 through 10 contiguous members;

R_{XV-30} , when bonded to A_{XV-1} , is taken together to form an intra-ring branched spacer connecting the A_{XV-1} -carbon at the point of attachment of R_{XV-30} to the points of bonding of each member of any one of substituent pairs selected from the group consisting of substituent pairs R_{XV-10} and R_{XV-11} , R_{XV-10} and R_{XV-31} , R_{XV-10} and R_{XV-32} , R_{XV-10} and R_{XV-12} , R_{XV-11} and R_{XV-31} , R_{XV-11} and R_{XV-32} , R_{XV-11} and R_{XV-12} , R_{XV-31} and R_{XV-32} , R_{XV-31} and R_{XV-12} , and R_{XV-32} and R_{XV-12} and

wherein said intra-ring branched spacer is selected to form two rings selected from the group consisting of cycloalkyl having from 3 through 10 contiguous members, cycloalkenyl having from 5 through 10 contiguous members, and heterocyclyl having from 5 through 10 contiguous members;

- 5 R_{XV-4} , R_{XV-5} , R_{XV-6} , R_{XV-7} , R_{XV-8} , R_{XV-9} , R_{XV-10} , R_{XV-11} , R_{XV-12} , R_{XV-13} , R_{XV-31} , R_{XV-32} , R_{XV-33} , R_{XV-34} , R_{XV-35} , and R_{XV-36} are independently selected from the group consisting of hydrido, carboxy, heteroaralkylthio, heteroaralkoxy, cycloalkylamino, acylalkyl, acylalkoxy, aroylalkoxy, heterocyclyloxy, aralkylaryl, aralkyl, aralkenyl, aralkynyl, heterocyclyl, perhaloaralkyl, aralkylsulfonyl, 10 aralkylsulfonylalkyl, aralkylsulfinyl, aralkylsulfinylalkyl, halocycloalkyl, halocycloalkenyl, cycloalkylsulfinyl, cycloalkylsulfinylalkyl, cycloalkylsulfonyl, cycloalkylsulfonylalkyl, heteroaryl amino, N-heteroaryl amino-N-alkyl amino, heteroaryl aminoalkyl, haloalkylthio, alkanoyloxy, alkoxy, alkoxyalkyl, haloalkoxylalkyl, heteroaralkoxy, cycloalkoxy, cycloalkenyloxy, cycloalkoxyalkyl, 15 cycloalkylalkoxy, cycloalkenyloxyalkyl, cycloalkylenedioxy, halocycloalkoxy, halocycloalkoxyalkyl, halocycloalkenyloxy, halocycloalkenyloxyalkyl, hydroxy, amino, thio, nitro, lower alkyl amino, alkylthio, alkylthioalkyl, aryl amino, aralkyl amino, arylthio, arylthioalkyl, heteroaralkoxyalkyl, alkylsulfinyl, 20 alkylsulfinylalkyl, arylsulfinylalkyl, arylsulfonylalkyl, heteroaryl sulfinylalkyl, heteroaryl sulfonylalkyl, alkylsulfonyl, alkylsulfonylalkyl, haloalkylsulfinylalkyl, haloalkylsulfonylalkyl, alkylsulfonamido, alkylaminosulfonyl, amidosulfonyl, monoalkylamid sulfonyl, dialkyl amid sulfonyl, monoarylamid sulfonyl, arylsulfonamido, diarylamid sulfonyl, monoalkyl monoaryl amid sulfonyl, 25 arylsulfinyl, arylsulfonyl, heteroarylthio, heteroaryl sulfinyl, heteroaryl sulfonyl, heterocyclylsulfonyl, heterocyclylthio, alkanoyl, alkenoyl, aroyl, heteroaroyl, aralkanoyl, heteroaralkanoyl, haloalkanoyl, alkyl, alkenyl, alkynyl, alkenyloxy, alkenyloxyalkyl, alkylenedioxy, haloalkylenedioxy, cycloalkyl, cycloalkylalkanoyl, cycloalkenyl, lower cycloalkylalkyl, lower cycloalkenylalkyl, halo, haloalkyl, 30 haloalkenyl, haloalkoxy, hydroxyhaloalkyl, hydroxyaralkyl, hydroxyalkyl, hydroxyheteroaralkyl, haloalkoxyalkyl, aryl, heteroaralkynyl, aryloxy, aralkoxy, aryloxyalkyl, saturated heterocyclyl, partially saturated heterocyclyl, heteroaryl, heteroaryloxy, heteroaryloxyalkyl, arylalkenyl, heteroarylalkenyl, carboxyalkyl, carboalkoxy, alkoxycarboxamido, 35 alkylamidocarbonylamido, alkylamidocarbonylamido, carboalkoxyalkyl,

carboalkoxyalkenyl, carboaralkoxy, carboxamido, carboxamidoalkyl, cyano, carbohaloalkoxy, phosphono, phosphonoalkyl, diaralkoxyphosphono, and diaralkoxyphosphonoalkyl with the provisos that R_{XV-4} , R_{XV-5} , R_{XV-6} , R_{XV-7} , R_{XV-8} , R_{XV-9} , R_{XV-10} , R_{XV-11} , R_{XV-12} , R_{XV-13} , R_{XV-31} , R_{XV-32} , R_{XV-33} , R_{XV-34} , R_{XV-35} , and R_{XV-36} are each independently selected to maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen, that no more than three of the R_{XV-33} and R_{XV-34} substituents are simultaneously selected from other than the group consisting of hydrido and halo, and that no more than three of the R_{XV-35} and R_{XV-36} substituents are simultaneously selected from other than the group consisting of hydrido and halo;

15 R_{XV-9}, R_{XV-10}, R_{XV-11}, R_{XV-12}, R_{XV-13}, R_{XV-31}, and R_{XV-32} are independently selected to be oxo with the provisos that B_{XV-1}, B_{XV-2}, D_{XV-3}, D_{XV-4}, J_{XV-3}, J_{XV-4}, and K_{XV-2} are independently selected from the group consisting of C and S, no more than two of R_{XV-9}, R_{XV-10}, R_{XV-11}, R_{XV-12}, R_{XV-13}, R_{XV-31}, and R_{XV-32} are simultaneously oxo, and that R_{XV-9}, R_{XV-10}, R_{XV-11}, R_{XV-12}, R_{XV-13}, R_{XV-31}, and R_{XV-32} are each independently selected to maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen;

20 R_{XV-4} and R_{XV-5} , R_{XV-5} and R_{XV-6} , R_{XV-6} and R_{XV-7} , R_{XV-7} and R_{XV-8} , R_{XV-9} and
 R_{XV-10} , R_{XV-10} and R_{XV-11} , R_{XV-11} and R_{XV-31} , R_{XV-31} and R_{XV-32} , R_{XV-32} and R_{XV-12} ,
and R_{XV-12} and R_{XV-13} are independently selected to form spacer pairs wherein a
spacer pair is taken together to form a linear moiety having from 3 through 6
contiguous atoms connecting the points of bonding of said spacer pair members
25 to form a ring selected from the group consisting of a cycloalkenyl ring having 5
through 8 contiguous members, a partially saturated heterocyclyl ring having 5
through 8 contiguous members, a heteroaryl ring having 5 through 6 contiguous
members, and an aryl with the provisos that no more than one of the group
consisting of spacer pairs R_{XV-4} and R_{XV-5} , R_{XV-5} and R_{XV-6} , R_{XV-6} and R_{XV-7} , R_{XV-7}
30 and R_{XV-8} is used at the same time and that no more than one of the group
consisting of spacer pairs R_{XV-9} and R_{XV-10} , R_{XV-10} and R_{XV-11} , R_{XV-11} and R_{XV-31} ,
 R_{XV-31} and R_{XV-32} , R_{XV-32} and R_{XV-12} , and R_{XV-12} and R_{XV-13} are used at the same
time;

35 R_{XV-9} and R_{XV-11} , R_{XV-9} and R_{XV-12} , R_{XV-9} and R_{XV-13} , R_{XV-9} and R_{XV-31} , R_{XV-9} and R_{XV-32} , R_{XV-10} and R_{XV-12} , R_{XV-10} and R_{XV-13} , R_{XV-10} and R_{XV-31} , R_{XV-10} and R_{XV-32} , R_{XV-11} and R_{XV-12} , R_{XV-11} and R_{XV-13} , R_{XV-11} and R_{XV-31} , R_{XV-11} and R_{XV-32} , R_{XV-12} and R_{XV-13} , R_{XV-12} and R_{XV-31} , R_{XV-12} and R_{XV-32} , R_{XV-13} and R_{XV-31} , R_{XV-13} and R_{XV-32} , R_{XV-31} and R_{XV-32} .

32, R_{XV-11} and R_{XV-12} , R_{XV-11} and R_{XV-13} , R_{XV-11} and R_{XV-32} , R_{XV-12} and R_{XV-31} , R_{XV-13} and R_{XV-31} , and R_{XV-13} and R_{XV-32} are independently selected to form a spacer pair wherein said spacer pair is taken together to form a linear spacer moiety selected from the group consisting of a covalent single bond and a moiety having from 1 through 3 contiguous atoms to form a ring selected from the group consisting of a cycloalkyl having from 3 through 8 contiguous members, a cycloalkenyl having from 5 through 8 contiguous members, a saturated heterocyclyl having from 5 through 8 contiguous members and a partially saturated heterocyclyl having from 5 through 8 contiguous members with the provisos that no more than one of said group of spacer pairs is used at the same time;

R_{XV-37} and R_{XV-38} are independently selected from the group consisting of hydrido, alkoxy, alkoxyalkyl, hydroxy, amino, thio, halo, haloalkyl, alkylamino, alkylthio, alkylthioalkyl, cyano, alkyl, alkenyl, haloalkoxy, and haloalkoxyalkyl.

Compounds of Formula XV and their methods of manufacture are disclosed in PCT Publication No. WO 00/18723, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula XV:

- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl] (cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl] (cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl] (cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][(3-trifluoromethyl)cyclohexylmethyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][(3-pentafluoroethyl) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][(3-trifluoromethoxy) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][(3-(1,1,2,2-tetrafluoroethoxy)cyclo-hexylmethyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]

- (cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]
 (cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]
 5 (cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl][(3-trifluoromethyl)cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]](3-pentafluoroethyl)cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 10 3-[[3-(3-trifluoromethoxyphenoxy)phenyl][(3-
 trifluoromethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]](3-(1,1,2,2-
 tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl](cyclohexylmethyl)amino]-1,1,1-
 15 trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl](cyclopentylmethyl)amino]-1,1,1-
 trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl](cyclopropylmethyl)amino]-1,1,1-
 trifluoro-2-propanol;
 20 3-[[3-(3-isopropylphenoxy)phenyl][(3-trifluoromethyl) cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl][(3-pentafluoroethyl) cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl][(3-trifluoromethoxy) cyclohexyl-
 25 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl]](3-(1,1,2,2-
 tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(2,3-dichlorophenoxy)phenyl](cyclohexylmethyl)amino]-1,1,1-
 trifluoro-2-propanol;
 30 3-[[3-(2,3-dichlorophenoxy)phenyl](cyclopentylmethyl)
 amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(2,3-dichlorophenoxy)phenyl](cyclopropylmethy)amino]-1,1,1-
 trifluoro-2-propanol;
 3-[[3-(2,3-dichlorophenoxy)phenyl][(3-trifluoromethyl)
 35 cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[3-(2,3-dichlorophenoxy)phenyl][(3-pentafluoroethyl) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2,3-dichlorophenoxy)phenyl][(3-trifluoromethoxy) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[3-(2,3-dichlorophenoxy)phenyl][3-(1,1,2,2-tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl](cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl](cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 10 2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl](cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][(3-trifluoromethyl) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-(4-fluorophenoxy)phenyl][(3-pentafluoroethyl) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][(3-trifluoromethoxy) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][3-(1,1,2,2-tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl](cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl](cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl](cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl][(3-trifluoromethyl)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl][(3-pentafluoroethyl)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl][(3-trifluoromethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl][3-(1,1,2,2-tetrafluoroethoxy)-cyclohexylmethyl]amino]-1,1,1-trifluoro-2-propanol;
- 35 3-[[3-(3-trifluoromethylbenzyloxy)phenyl]

- (cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethylbenzyloxy)phenyl]
 (cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethylbenzyloxy)phenyl]
 5 (cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethylbenzyloxy)phenyl][(3-trifluoromethyl)cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethylbenzyloxy)phenyl][(3-pentafluoroethyl)cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 10 3-[[3-(3-trifluoromethylbenzyloxy)phenyl][(3-
 trifluoromethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethylbenzyloxy)phenyl][3-(1,1,2,2-
 tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethyl)phenyl]methyl](cyclohexyl)amino]-1,1,1-trifluoro-2-
 15 propanol;
 3-[[[(3-pentafluoroethyl)phenyl]methyl](cyclohexyl)amino]-1,1,1-trifluoro-
 2-propanol;
 3-[[[(3-trifluoromethoxy)phenyl]methyl](cyclohexyl)amino]-1,1,1-trifluoro-
 2-propanol;
 20 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]
 methyl](cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethyl)phenyl]methyl]
 (4-methylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-pentafluoroethyl)phenyl]methyl]
 25 (4-methylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethoxy)phenyl]methyl]
 (4-methylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](4-
 methylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 30 3-[[[(3-trifluoromethyl)phenyl]methyl](3-
 trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-pentafluoroethyl)phenyl]methyl](3-
 trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethoxy)phenyl]methyl](3-
 35 trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;

- 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](3-trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)cyclohexyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[[3-(3-pentafluoroethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)cyclohexyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethoxy)phenyl]methyl][3-(4-chloro-3-methylphenoxy)cyclohexyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)-cyclohexyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethyl)phenyl]methyl](3-phenoxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-pentafluoroethyl)phenyl]methyl](3-phenoxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[[3-(3-trifluoromethoxy)phenyl]methyl](3-phenoxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](3-phenoxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[[3-(3-trifluoromethyl)phenyl]methyl](3-isopropoxycyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-pentafluoroethyl)phenyl]methyl](3-isopropoxycyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethoxy)phenyl]methyl](3-isopropoxycyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](3-isopropoxycyclohexyl)-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethyl)phenyl]methyl](3-cyclopentyloxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[[3-(3-pentafluoroethyl)phenyl]methyl](3-cyclopentyloxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethoxy)phenyl]methyl](3-cyclopentyloxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](3-cyclopentyloxy-cyclohexyl)-amino]-1,1,1-trifluoro-2-propanol;
- 35 3-[[[3-(2-trifluoromethyl)pyrid-6-yl]methyl](3-

- isopropoxycyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-cyclopentyloxycyclohexyl)-
 amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-phenoxy-cyclohexyl)amino]-
 5 1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-
 trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl][3-(4-chloro-3-
 ethylphenoxy)cyclo-hexyl]amino]-1,1,1-trifluoro-2-propanol;
 10 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl][3-(1,1,2,2-
 tetrafluoroethoxy)cyclo-hexyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-pentafluoroethylcyclohexyl)-
 amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-trifluoromethoxycyclohexyl)-
 15 amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)propyl]-
 amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-pentafluoroethyl)phenyl]methyl][3-(4-chloro-3-
 ethylphenoxy)propyl]-amino]-1,1,1-trifluoro-2-propanol;
 20 3-[[[(3-trifluoromethoxy)phenyl]methyl][3-(4-chloro-3-
 ethylphenoxy)propyl]-amino]-1,1,1-trifluoro-2-propanol;
 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-(4-chloro-3-
 ethylphenoxy)-propyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)-2,2,-di-
 25 fluoropropyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-pentafluoroethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)-2,2-
 di-fluoropropyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethoxy)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)-2,2,-
 di-fluoropropyl]amino]-1,1,1-trifluoro-2-propanol;
 30 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-(4-chloro-3-
 ethylphenoxy)-2,2,-difluoropropyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethyl)phenyl]methyl][3-(isopropoxy)propyl]amino]-1,1,1-
 trifluoro-2-propanol;
 3-[[[(3-pentafluoroethyl)phenyl]methyl][3-(isopropoxy)propyl]amino]-
 35 1,1,1-trifluoro-2-propanol;

103


3-[[[(3-trifluoromethoxy)phenyl]methyl][3-(isopropoxy)propyl]amino]-
1,1,1-trifluoro-2-propanol;

3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-
(isopropoxy)propyl]amino]-1,1,1-trifluoro-2-propanol; and

3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-
(phenoxy)propyl]amino]-1,1,1-trifluoro-2-propanol.

Another class of CETP inhibitors that finds utility with the present
invention consists of (*R*)-chiral halogenated 1-substituted amino-(*n*+1)-alkanols
having the Formula XVI

10



Formula XVI

and pharmaceutically acceptable forms thereof, wherein:


15

*n*_{XVI} is an integer selected from 1 through 4;


X_{XVI} is oxy;

R_{XVI-1} is selected from the group consisting of haloalkyl, haloalkenyl,
haloalkoxymethyl, and haloalkenyloxymethyl with the proviso that R_{XVI-1} has a
higher Cahn-Ingold-Prelog stereochemical system ranking than both R_{XVI-2} and
(CHR_{XVI-3})_n-N(A_{XVI})Q_{XVI} wherein A_{XVI} is Formula XVI-(II) and Q is Formula XVI-
(III);

20



XVI-II



XVI-III

R_{XVI-16} is selected from the group consisting of hydrido, alkyl, acyl, aroyl, heteroaroyl, trialkylsilyl, and a spacer selected from the group consisting of a covalent single bond and a linear spacer moiety having a chain length of 1 to 4 atoms linked to the point of bonding of any aromatic substituent selected from the group consisting of R_{XVI-4} , R_{XVI-8} , R_{XVI-9} , and R_{XVI-13} to form a heterocyclyl ring having from 5 through 10 contiguous members;

D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} are independently selected from the group consisting of C, N, O, S and covalent bond with the provisos that no more than one of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} is a covalent bond, no more than one D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} is be O, no more than one of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} is S, one of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} must be a covalent bond when two of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} are O and S, and no more than four of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} is N;

D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} are independently selected from the group consisting of C, N, O, S and covalent bond with the provisos that no more than one is a covalent bond, no more than one of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} is O, no more than one of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} is S, no more than two of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} is O and S, one of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} must be a covalent bond when two of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} are O and S, and no more than four of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} are N;

R_{XVI-2} is selected from the group consisting of hydrido, aryl, aralkyl, alkyl, alkenyl, alkenyloxyalkyl, haloalkyl, haloalkenyl, halocycloalkyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, dicyanoalkyl, and carboalkoxycyanoalkyl, with the proviso that R_{XVI-2} has a lower Cahn-Ingold-Prelog system ranking than both R_{XVI-1} and $(CHR_{XVI-3})_n-N(A_{XVI})Q_{XVI}$;

R_{XVI-3} is selected from the group consisting of hydrido, hydroxy, cyano, aryl, aralkyl, acyl, alkoxy, alkyl, alkenyl, alkoxyalkyl, heteroaryl, alkenyloxyalkyl, haloalkyl, haloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, monocyanoalkyl, dicyanoalkyl, carboxamide, and carboxamidoalkyl, with the provisos that $(CHR_{XVI-3})_n-N(A_{XVI})Q_{XVI}$ has a lower Cahn-Ingold-Prelog stereochemical system ranking than R_{XVI-1} and a higher Cahn-Ingold-Prelog stereochemical system ranking than R_{XVI-2} ;

Y_{XVI} is selected from a group consisting of a covalent single bond, $(C(R_{XVI-14})_2)_q$ wherein q is an integer selected from 1 and 2 and $(CH(R_{XVI-14}))_g-W_{XVI}-(CH(R_{XVI-14}))_p$ wherein g and p are integers independently selected from 0 and 1;

R_{XVI-14} is selected from the group consisting of hydrido, hydroxy, cyano, hydroxyalkyl, acyl, alkoxy, alkyl, alkenyl, alkynyl, alkoxyalkyl, haloalkyl, haloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, monocarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, carboalkoxy, carboxamide, and carboxamidoalkyl;

Z_{XVI} is selected from a group consisting of a covalent single bond, $(C(R_{XVI-15})_2)_q$, wherein q is an integer selected from 1 and 2, and $(CH(R_{XVI-15}))_j-W_{XVI}-(CH(R_{XVI-15}))_k$ wherein j and k are integers independently selected from 0 and 1;

W_{XVI} is selected from the group consisting of O, C(O), C(S), C(O)N(R_{XVI-14}), C(S)N(R_{XVI-14}), (R_{XVI-14})NC(O), (R_{XVI-14})NC(S), S, S(O), S(O)₂, S(O)₂N(R_{XVI-14}), (R_{XVI-14})NS(O)₂, and N(R_{XVI-14}) with the proviso that R_{XVI-14} is other than cyano;

R_{XVI-15} is selected, from the group consisting of hydrido, cyano, hydroxyalkyl, acyl, alkoxy, alkyl, alkenyl, alkynyl, alkoxyalkyl, haloalkyl, haloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, monocarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, carboalkoxy, carboxamide, and carboxamidoalkyl;

R_{XVI-4} , R_{XVI-5} , R_{XVI-6} , R_{XVI-7} , R_{XVI-8} , R_{XVI-9} , R_{XVI-10} , R_{XVI-11} , R_{XVI-12} , and R_{XVI-13}
 are independently selected from the group consisting of hydrido, carboxy,
 heteroaralkylthio, heteroaralkoxy, cycloalkylamino, acylalkyl, acylalkoxy,
 aroylalkoxy, heterocyclyloxy, aralkylaryl, aralkyl, aralkenyl, aralkynyl,
 5 heterocyclyl, perhaloaralkyl, aralkylsulfonyl, aralkylsulfonylalkyl, aralkylsulfinyl,
 aralkylsulfinylalkyl, halocycloalkyl, halocycloalkenyl, cycloalkylsulfinyl,
 cycloalkylsulfinylalkyl, cycloalkylsulfonyl, cycloalkylsulfonylalkyl,
 heteroaryl amino, N-heteroaryl amino-N-alkyl amino, heteroaralkyl,
 heteroaryl aminoalkyl, haloalkylthio, alkanoyloxy, alkoxy, alkoxyalkyl,
 10 haloalkoxylalkyl, heteroaralkoxy, cycloalkoxy, cycloalkenyloxy, cycloalkoxyalkyl,
 cycloalkylalkoxy, cycloalkenyloxyalkyl, cycloalkylenedioxy, halocycloalkoxy,
 halocycloalkoxyalkyl, halocycloalkenyloxy, halocycloalkenyloxyalkyl, hydroxy,
 amino, thio, nitro, lower alkyl amino, alkylthio, alkylthioalkyl, aryl amino,
 aralkyl amino, arylthio, arylthioalkyl, heteroaralkoxyalkyl, alkylsulfinyl,
 15 alkylsulfinylalkyl, arylsulfinylalkyl, arylsulfonylalkyl, heteroaryl sulfinylalkyl,
 heteroaryl sulfonylalkyl, alkylsulfonyl, alkylsulfonylalkyl, haloalkylsulfinylalkyl,
 haloalkylsulfonylalkyl, alkylsulfonamido, alkylaminosulfonyl, amidosulfonyl,
 monoalkyl amidosulfonyl, dialkyl, amidosulfonyl, monoaryl amidosulfonyl,
 arylsulfonamido, diarylamidosulfonyl, monoalkyl monoaryl amidosulfonyl,
 20 arylsulfinyl, arylsulfonyl, heteroarylthio, heteroaryl sulfinyl, heteroaryl sulfonyl,
 heterocyclylsulfonyl, heterocyclylthio, alkanoyl, alkenoyl, aroyl, heteroaroyl,
 aralkanoyl, heteroaralkanoyl, haloalkanoyl, alkyl, alkenyl, alkynyl, alkenyloxy,
 alkenyloxyalkyl, alkylenedioxy, haloalkylenedioxy, cycloalkyl, cycloalkylalkanoyl,
 cycloalkenyl, lower cycloalkylalkyl, lower cycloalkenylalkyl, halo, haloalkyl,
 25 haloalkenyl, haloalkoxy, hydroxyhaloalkyl,
 hydroxyaralkyl, hydroxyalkyl, hydroxyheteroaralkyl, haloalkoxyalkyl, aryl,
 heteroaralkynyl, aryloxy, aralkoxy, aryloxyalkyl, saturated heterocyclyl, partially
 saturated heterocyclyl, heteroaryl, heteroaryl oxy, heteroaryl oxyalkyl, arylalkenyl,
 heteroarylalkenyl, carboxyalkyl, carboalkoxy, alkoxycarboxamido,
 30 alkylamidocarbonylamido, arylamidocarbonylamido, carboalkoxyalkyl,
 carboalkoxyalkenyl, carboaralkoxy, carboxamido, carboxamidoalkyl, cyano,
 carbohaloalkoxy, phosphono, phosphonoalkyl, diaralkoxyphosphono, and
 diaralkoxyphosphonoalkyl with the proviso that R_{XVI-4} , R_{XVI-5} , R_{XVI-6} , R_{XVI-7} , R_{XVI-8} ,
 R_{XVI-9} , R_{XVI-10} , R_{XVI-11} , R_{XVI-12} , and R_{XVI-13} are each independently selected to

maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen;

R_{XVI-4} and R_{XVI-5} , R_{XVI-5} and R_{XVI-6} , R_{XVI-6} and R_{XVI-7} , R_{XVI-7} and R_{XVI-8} , R_{XVI-9} and R_{XVI-10} , R_{XVI-10} and R_{XVI-11} , R_{XVI-11} and R_{XVI-12} , and R_{XVI-12} and R_{XIV-13} are independently selected to form spacer pairs wherein a spacer pair is taken together to form a linear moiety having from 3 through 6 contiguous atoms connecting the points of bonding of said spacer pair members to form a ring selected from the group consisting of a cycloalkenyl ring having 5 through 8 contiguous members, a partially saturated heterocyclyl ring having 5 through 8 contiguous members, a heteroaryl ring having 5 through 6 contiguous members, and an aryl with the provisos that no more than one of the group consisting of spacer pairs R_{XVI-4} and R_{XVI-5} , R_{XVI-5} and R_{XVI-6} , R_{XVI-6} and R_{XVI-7} , and R_{XVI-7} and R_{XVI-8} is used at the same time and that no more than one of the group consisting of spacer pairs R_{XIV-9} and R_{XVI-10} , R_{XVI-10} and R_{XVI-11} , R_{XVI-11} and R_{XVI-12} , and R_{XVI-12} and R_{XVI-13} can be used at the same time;

R_{XVI-4} and R_{XVI-9} , R_{XVI-4} and R_{XVI-13} , R_{XVI-8} and R_{XVI-9} , and R_{XVI-8} and R_{XVI-13} is independently selected to form a spacer pair wherein said spacer pair is taken together to form a linear moiety wherein said linear moiety forms a ring selected from the group consisting of a partially saturated heterocyclyl ring having from 5 through 8 contiguous members and a heteroaryl ring having from 5 through 6 contiguous members with the proviso that no more than one of the group consisting of spacer pairs R_{XVI-4} and R_{XVI-9} , R_{XVI-4} and R_{XVI-13} , R_{XVI-8} and R_{XVI-9} , and R_{XVI-8} and R_{XVI-13} is used at the same time.

Compounds of Formula XVI and their methods of manufacture are disclosed in PCT Publication No. WO 00/18724, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula XVI:

- (2R)-3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-isopropylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-cyclopropylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;

- (2R)-3-[[3-(3-(2-furyl)phenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2,3-dichlorophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[3-(4-fluorophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(4-methylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 (2R)-3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoro-ethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3,5-dimethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-ethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 (2R)-3-[[3-(3-t-butylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-methylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 (2R)-3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[3-(1,1,2,2-tetrafluoro-ethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(phenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoro-ethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethoxy)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoro-methyl)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

- (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
5 (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[cyclohexylmethoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
10 (2R)-3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[3-(3-difluoromethoxyphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
15 (2R)-3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
20 (2R)-3-[[3-(3-isopropylphenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[3-(3-cyclopropylphenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
25 (2R)-3-[[3-(3-(2-furyl)phenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[3-(2,3-dichlorophenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[3-(4-fluorophenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
30 (2R)-3-[[3-(4-methylphenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
35 (2R)-3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-

- (pentafluoroethyl)phenyl)methyl]-amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][3-(pentafluoroethyl)-phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 5 (2R)-3-[[3-(3,5-dimethylphenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]-amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(3-ethylphenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 10 (2R)-3-[[3-(3-t-butylphenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(3-methylphenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 15 (2R)-3-[[3-(phenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 20 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[[3-(trifluoromethoxy)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[[3-(trifluoromethyl)-phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 25 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[[3,5-dimethylphenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[[3-(trifluoromethylthio)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[[3,5-difluorophenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
 30 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[cyclohexylmethoxy]phenyl]-amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;



- (2R)-3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(3-difluoromethoxyphenoxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[3-(pentafluoroethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 (2R)-3-[[[3-(3-isopropylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(3-cyclopropylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[[3-(3-(2-furyl)phenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(2,3-dichlorophenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(4-fluorophenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 (2R)-3-[[[3-(4-methylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 (2R)-3-[[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[3-(heptafluoropropyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 (2R)-3-[[[3-(3,5-dimethylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(3-ethylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

MISSING AT THE TIME OF PUBLICATION

- (2R)-3-[[3-(3-isopropylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-cyclopropylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[3-(3-(2-furyl)phenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2,3-dichlorophenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(4-fluorophenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-3-propanol;
- 10 (2R)-3-[[3-(4-methylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[2-fluoro-5-(trifluoro-methyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 (2R)-3-[[3-[3-(3,5-dimethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-ethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 25 (2R)-3-[[3-(3-t-butylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-methylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 (2R)-3-[[3-(phenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(N,N-dimethylamino,phenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 35 (2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[3-

- (trifluoromethoxy)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-3-propanol;
 (2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethyl)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 5 (2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 10 (2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[cyclohexylmethoxyl-phenyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 15 (2R)-3-[[3-(3-difluoromethoxyphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 20 (2R)-3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[2-fluoro-5-(trifluoro-methyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(3-isopropylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 25 (2R)-3-[[3-(3-cyclopropylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(3-(2-furyl)phenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 30 (2R)-3-[[3-(2,3-dichlorophenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(4-fluorophenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(4-methylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 35 (trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;

- (2R)-3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3,5-dimethylphenoxy)phenyl][[2-fluoro-4-
- 10 (trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-ethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-t-butylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[3-(3-methylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(phenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)
- 20 phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethoxy)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 (3R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethyl)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-
- 30 (trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[cyclohexylmethoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;



wherein

R_{XVII-6} , R_{XVII-7} , $R_{XVII-10}$ denote, independently from one another, a
 cycloalkyl containing 3 to 6 carbon atoms, or an aryl containing 6 to 10 carbon
 atom or a 5- to 7-membered, optionally benzo-condensed, saturated or
 5 unsaturated, mono-, bi- or tricyclic heterocycle containing up to 4 heteroatoms
 from the series of S, N and/or O, wherein the rings are optionally substituted, in
 the case of the nitrogen-containing rings also via the N function, with up to five
 identical or different substituents in the form of a halogen, trifluoromethyl, nitro,
 10 hydroxyl, cyano, carboxyl, trifluoromethoxy, a straight-chain or branched acyl,
 alkyl, alkylthio, alkylalkoxy, alkoxy or alkoxycarbonyl containing up to 6 carbon
 atoms each, an aryl or trifluoromethyl-substituted aryl containing 6 to 10 carbon
 atoms each, or an optionally benzo-condensed, aromatic 5- to 7-membered
 heterocycle containing up to 3 heteroatoms from the series of S, N and/or O,
 15 and/or in the form of a group according to the formula $-OR_{XVII-11}$, $-SR_{XVII-12}$,
 $-SO_2R_{XVII-13}$, or $-NR_{XVII-14}R_{XVII-15}$;


$R_{XVII-11}$, $R_{XVII-12}$, and $R_{XVII-13}$ denote, independently from one another, an
 aryl containing 6 to 10 carbon atoms, which is in turn substituted with up to two
 identical or different substituents in the form of a phenyl, halogen or a straight-
 chain or branched alkyl containing up to 6 carbon atoms,
 20

$R_{XVII-14}$ and $R_{XVII-15}$ are identical or different and have the meaning of
 R_{XVII-4} and R_{XVII-5} given above, or

R_{XVII-6} and/or R_{XVII-7} denote a radical according to the formula

- (2R)-3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 5 (2R)-3-[[3-(3-difluoromethoxyphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol; and
 10 (2R)-3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol.

Another class of CETP inhibitors that finds utility with the present invention consists of quinolines of Formula XVII



Formula XVII

15

and pharmaceutically acceptable forms thereof, wherein:

A_{XVII} denotes an aryl containing 6 to 10 carbon atoms, which is optionally substituted with up to five identical or different substituents in the form of a halogen, nitro, hydroxyl, trifluoromethyl, trifluoromethoxy or a straight-chain or branched alkyl, acyl, hydroxyalkyl or alkoxy containing up to 7 carbon atoms
 20 each, or in the form of a group according to the formula -NR_{XVII-4}R_{XVII-5}, wherein R_{XVII-4} and R_{XVII-5} are identical or different and denote a hydrogen, phenyl or a straight-chain or branched alkyl containing up to 6 carbon atoms,

D_{XVII} denotes an aryl containing 6 to 10 carbon atoms, which is optionally substituted with a phenyl, nitro, halogen, trifluoromethyl or trifluoromethoxy, or a radical according to the formula
 25

R_{XVII-1} and/or R_{XVII-2} are straight-chain or branched alkyl with up to 6 carbon atoms, optionally substituted with halogen, trifluoromethoxy, hydroxy, or a straight-chain or branched alkoxy with up to 4 carbon atoms, aryl containing 6-10 carbon atoms optionally substituted with up to five of the same or different substituents selected from halogen, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, nitro, straight-chain or branched alkyl, acyl, hydroxyalkyl, alkoxy with up to 7 carbon atoms and $NR_{XVII-22}R_{XVII-23}$;

$R_{XVII-22}$ and $R_{XVII-23}$ are identical or different and denote hydrogen, phenyl or a straight-chain or branched alkyl up to 6 carbon atoms; and/or

R_{XVII-1} and R_{XVII-2} taken together form a straight-chain or branched alkene or alkane with up to 6 carbon atoms optionally substituted with halogen, trifluoromethyl, hydroxy or straight-chain or branched alkoxy with up to 5 carbon atoms;


R_{XVII-3} denotes hydrogen, a straight-chain or branched acyl with up to 20 carbon atoms, a benzoyl optionally substituted with halogen, trifluoromethyl, nitro or trifluoromethoxy, a straight-chained or branched fluoroacyl with up to 8 carbon atoms and 7 fluoro atoms, a cycloalkyl with 3 to 7 carbon atoms, a straight chained or branched alkyl with up to 8 carbon atoms optionally substituted with hydroxyl, a straight-chained or branched alkoxy with up to 6 carbon atoms optionally substituted with phenyl which may in turn be substituted with halogen, nitro, trifluoromethyl, trifluoromethoxy, or phenyl or a tetrazol substituted phenyl, and/or an alkyl that is optionally substituted with a group according to the formula $-OR_{XVII-24}$;

$R_{XVII-24}$ is a straight-chained or branched acyl with up to 4 carbon atoms or benzyl.

Compounds of Formula XVII and their methods of manufacture are disclosed in PCT Publication No. WO 98/39299, which is incorporated herein by reference in its entirety for all purposes.

Another class of CETP inhibitors that finds utility with the present invention consists of 4-Phenyltetrahydroquinolines of Formula XVIII

118



R_{XVII-8} denotes a hydrogen or halogen, and

R_{XVII-9} denotes a hydrogen, halogen, azido, trifluoromethyl, hydroxyl, trifluoromethoxy, a straight-chain or branched alkoxy or alkyl containing up to 6 carbon atoms each, or a radical according to the formula $NR_{XVII-16}R_{XVII-17}$;

$R_{XVII-16}$ and $R_{XVII-17}$ are identical or different and have the meaning of R_{XVII-4} and R_{XVII-5} above; or

R_{XVII-8} and R_{XVII-9} together form a radical according to the formula $=O$ or $=NR_{XVII-18}$;

$R_{XVII-18}$ denotes a hydrogen or a straight-chain or branched alkyl, alkoxy or acyl containing up to 6 carbon atoms each;

L_{XVII} denotes a straight-chain or branched alkylene or alkenylene chain containing up to 8 carbon atoms each, which are optionally substituted with up to two hydroxyl groups;

T_{XVII} and X_{XVII} are identical or different and denote a straight-chain or branched alkylene chain containing up to 8 carbon atoms; or

T_{XVII} and X_{XVII} denotes a bond;

V_{XVII} denotes an oxygen or sulfur atom or $-NR_{XVII-19}$;

$R_{XVII-19}$ denotes a hydrogen or a straight-chain or branched alkyl containing up to 6 carbon atoms or a phenyl;

E_{XVII} denotes a cycloalkyl containing 3 to 8 carbon atoms, or a straight-chain or branched alkyl containing up to 8 carbon atoms, which is optionally substituted with a cycloalkyl containing 3 to 8 carbon atoms or a hydroxyl, or a phenyl, which is optionally substituted with a halogen or trifluoromethyl;

R_{XVII-1} and R_{XVII-2} are identical or different and denote a cycloalkyl containing 3 to 8 carbon atoms, hydrogen, nitro, halogen, trifluoromethyl, trifluoromethoxy, carboxy, hydroxy, cyano, a straight-chain or branched acyl, alkoxycarbonyl or alkoxy with up to 6 carbon atoms, or $NR_{XVII-20}R_{XVII-21}$;

$R_{XVII-20}$ and $R_{XVII-21}$ are identical or different and denote hydrogen, phenyl, or a straight-chain or branched alkyl with up to 6 carbon atoms; and or

R_{XVIII-3} and R_{XVIII-4} taken together form an alkenylene made up of between two and four carbon atoms.

Compounds of Formula XVIII and their methods of manufacture are disclosed in PCT Publication No. WO 99/15504 and United States Patent No. 6,291,477, both of which are incorporated herein by reference in their entireties for all purposes.


The invention is particularly advantageous for the class of drugs which are both acid-sensitive and low-solubility. Exemplary acid-sensitive, low-solubility drugs include (+)-N-{3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl}-N-hydroxyurea; omeprazole; etoposide; famotidine; erythromycin; quinapril; lansoprazole; progabide; as well as CCR1 inhibitors such as quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3-fluorobenzyl-2(S),7-dihydroxy-7-methyloctyl]amide and quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide.

The invention is useful for improving the intrinsic dissolution rate of compounds selected from the following. The intrinsic dissolution rate is defined as the rate of dissolution of a pure pharmaceutical active ingredient when conditions such as surface area, agitation-stirring speed, pH and ionic-strength of the dissolution medium are kept constant. Intrinsic dissolution rate is further defined as being measured in water at 37°C using a USP II dissolution apparatus equipped with a Wood's apparatus (Wood, JH; Syarto, JE and Letterman, H: J.Pharm. Sci. 54 (1965), 1068) with a stirring speed of 50 rpm. The intrinsic dissolution rate is defined in terms of mg of drug dissolved per minute from a unit surface area, therefore, the intrinsic dissolution rate is referred to in units of mg/min.cm².

The compositions and methods of the invention are particularly useful for compounds with an intrinsic dissolution rate of preferably less than 0.1 mg/min.cm² and more preferably with less than 0.05 mg/min.cm².


Turning now to the chemical structures of specific CCR1 inhibitors, one class of CCR1 inhibitors that finds utility with the present invention consists of dihydroxyhexanoic acid derivatives having the Formula CCR1-I

120



Formula XVIII

5 , N oxides thereof, and pharmaceutically acceptable forms thereof, wherein:
 A_{XVIII} denotes a phenyl optionally substituted with up to two identical or different substituents in the form of halogen, trifluoromethyl or a straight-chain or branched alkyl or alkoxy containing up to three carbon atoms;
 D_{XVIII} denotes the formula




10 or $R_{XVIII-8}-CH_2-O-CH_2-$;

$R_{XVIII-5}$ and $R_{XVIII-6}$ are taken together to form $=O$; or
 $R_{XVIII-5}$ denotes hydrogen and $R_{XVIII-6}$ denotes halogen or hydrogen; or
 $R_{XVIII-5}$ and $R_{XVIII-6}$ denote hydrogen;
 $R_{XVIII-7}$ and $R_{XVIII-8}$ are identical or different and denote phenyl, naphthyl,
 15 benzothiazolyl, quinolinyl, pyrimidyl or pyridyl with up to four identical or different substituents in the form of halogen, trifluoromethyl, nitro, cyano, trifluoromethoxy, $-SO_2-CH_3$ or $NR_{XVIII-9}R_{XVIII-10}$;
 $R_{XVIII-9}$ and $R_{XVIII-10}$ are identical or different and denote hydrogen or a straight-chained or branched alkyl of up to three carbon atoms;
 E_{XVIII} denotes a cycloalkyl of from three to six carbon atoms or a straight-
 20 chained or branched alkyl of up to eight carbon atoms;
 $R_{XVIII-1}$ denotes hydroxy;
 $R_{XVIII-2}$ denotes hydrogen or methyl;
 $R_{XVIII-3}$ and $R_{XVIII-4}$ are identical or different and denote straight-chained or
 25 branched alkyl of up to three carbon atoms; or

(C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 5 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-
 (S=O)-, (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-
 (C₁-C₆)alkyl, (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl,
 CF₃SO₃-, (C₁-C₆)alkyl-SO₃-, phenyl, phenoxy, benzyloxy, (C₃-C₁₀)cycloalkyl,
 10 (C₂-C₉)heterocycloalkyl, and (C₂-C₉)heteroaryl;
 wherein R³ is hydrogen, (C₁-C₁₀)alkyl, (C₃-C₁₀)cycloalkyl-(CH₂)_n-, (C₂-
 C₉)heterocycloalkyl-(CH₂)_n-, (C₂-C₉)heteroaryl-(CH₂)_n- or aryl-(CH₂)_n-; wherein n
 is an interger from zero to six;

wherein said R₃ (C₁-C₁₀)alkyl group may optionally be substituted with
 15 one or more substituents, (preferably from one to three substituents)
 independently selected from hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-
 (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-
 (C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-
 (C=O)-O-, (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl,
 20 (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 25 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 30 (C₂-C₉)heteroaryl; and wherein any of the carbon-carbon single bonds of said
 (C₁-C₁₀)alkyl may optionally be replaced by a carbon-carbon double bond;
 wherein the (C₃-C₁₀)cycloalkyl moiety of said R₃ (C₃-C₁₀)cycloalkyl-
 (CH₂)_n- group may optionally be substituted by one to three substituents
 independently selected from the group consisting of hydrogen, halo, CN,
 35 (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy,

122



CCR1-I

wherein R_1 is (C_2 - C_9) heteroaryl optionally substituted with one, two or three substituents independently selected from the group consisting of hydrogen, halo, cyano, (C_1 - C_6)alkyl optionally substituted with one, two or three fluorine atoms, hydroxy, hydroxy-(C_1 - C_6)alkyl, (C_1 - C_6)alkoxy, (C_1 - C_6)alkoxy(C_1 - C_6)alkyl, $HO-(C=O)-$, (C_1 - C_6)alkyl- $O-(C=O)-$, $HO-(C=O)-$ (C_1 - C_6)alkyl, (C_1 - C_6)alkyl- $O-(C=O)-$ (C_1 - C_6)alkyl, (C_1 - C_6)alkyl- $(C=O)-O-$, (C_1 - C_6)alkyl- $(C=O)-O-(C_1$ - C_6)alkyl, $H(O=C)-$, $H(O=C)-(C_1$ - C_6)alkyl, (C_1 - C_6)alkyl($O=C)-$, (C_1 - C_6)alkyl($O=C)-(C_1$ - C_6)alkyl, NO_2 , amino, (C_1 - C_6)alkylamino, $[(C_1$ - C_6)alkyl] $_2$ amino, amino(C_1 - C_6)alkyl, (C_1 - C_6)alkylamino(C_1 - C_6)alkyl, $[(C_1$ - C_6)alkyl] $_2$ amino(C_1 - C_6)alkyl, $H_2N-(C=O)-$, (C_1 - C_6)alkyl- $NH-(C=O)-$, $[(C_1$ - C_6)alkyl] $_2N-(C=O)-$, $H_2N(C=O)-(C_1$ - C_6)alkyl, (C_1 - C_6)alkyl- $HN(C=O)-(C_1$ - C_6)alkyl, $[(C_1$ - C_6)alkyl] $_2N-(C=O)-(C_1$ - C_6)alkyl, $H(O=C)-NH-$, (C_1 - C_6)alkyl($C=O)-NH$, (C_1 - C_6)alkyl($C=O)-[NH](C_1$ - C_6)alkyl, (C_1 - C_6)alkyl($C=O)-[N(C_1$ - C_6)alkyl](C_1 - C_6)alkyl, (C_1 - C_6)alkyl- $S-$, (C_1 - C_6)alkyl- $(S=O)-$, (C_1 - C_6)alkyl- SO_2- , (C_1 - C_6)alkyl- SO_2-NH- , H_2N-SO_2- , H_2N-SO_2- (C_1 - C_6)alkyl, (C_1 - C_6)alkyl- $HN-SO_2-(C_1$ - C_6)alkyl, $[(C_1$ - C_6)alkyl] $_2N-SO_2-(C_1$ - C_6)alkyl, CF_3SO_3- , (C_1 - C_6)alkyl- SO_3- , phenyl, (C_3 - C_{10})cycloalkyl, (C_2 - C_9)heterocycloalkyl, and (C_2 - C_9)heteroaryl;

wherein R_2 is phenyl- $(CH_2)_m-$, naphthyl- $(CH_2)_m-$, (C_3 - C_{10})cycloalkyl- $(CH_2)_m-$, (C_1 - C_6)alkyl or (C_2 - C_9)heteroaryl- $(CH_2)_m-$, wherein each of said phenyl, naphthyl, (C_3 - C_{10})cycloalkyl or (C_2 - C_9)heteroaryl moieties of said phenyl- $(CH_2)_m-$, naphthyl- $(CH_2)_m-$, (C_3 - C_{10})cycloalkyl- $(CH_2)_m-$ or (C_2 - C_9)heteroaryl- $(CH_2)_m-$ groups may optionally be substituted with one, two, or three substituents independently selected from the group consisting of hydrogen, halo, cyano, (C_1 - C_6)alkyl, hydroxy, hydroxy-(C_1 - C_6)alkyl, (C_1 - C_6)alkoxy, (C_1 - C_6)alkoxy(C_1 - C_6)alkyl, $HO-(C=O)-$, (C_1 - C_6)alkyl- $O-(C=O)-$, $HO-(C=O)-$ (C_1 - C_6)alkyl, (C_1 - C_6)alkyl- $O-(C=O)-$ (C_1 - C_6)alkyl, (C_1 - C_6)alkyl- $(C=O)-O-$, (C_1 - C_6)alkyl- $(C=O)-O-(C_1$ - C_6)alkyl, $H(O=C)-$, $H(O=C)-(C_1$ - C_6)alkyl, (C_1 - C_6)alkyl($O=C)-$, (C_1 - C_6)alkyl($O=C)-(C_1$ - C_6)alkyl, NO_2 , amino,

(C₁-C₆)alkyl-SO₃⁻, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and (C₂-C₉)heteroaryl;

wherein the (C₂-C₉)heteroaryl moiety of said R³ (C₂-C₉)heteroaryl-(CH₂)_n- group may contain from one to three heteroatoms independently selected from
 5 nitrogen, sulfur or oxygen, wherein said (C₂-C₉)heteroaryl moiety of said (C₂-C₉)heteroaryl-(CH₂)_n- group may optionally be substituted on any of the ring carbon atoms capable of forming an additional bond (preferably one to three substituents per ring) with a substituent selected from the group consisting of
 10 hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-, (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino, (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 15 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-, (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-NH-, (C₁-C₆)alkyl(C=O)-NH-, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl, (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 20 (C₁-C₆)alkyl-SO₂⁻, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂⁻, H₂N-SO₂-(C₁-C₆)alkyl, (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃⁻, (C₁-C₆)alkyl-SO₃⁻, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and (C₂-C₉)heteroaryl; and

wherein said aryl moiety of said R₃ aryl-(CH₂)_n- group is optionally
 25 substituted phenyl or naphthyl, wherein said phenyl and naphthyl may optionally be substituted with from one to three substituents independently selected from the group consisting of hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-, (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino, (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 30 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-, (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 35 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-

(C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 5 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 10 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl HN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl;

15 wherein the (C₂-C₉)heterocycloalkyl moiety of said R₃ (C₂-
 C₉)heterocycloalkyl-(CH₂)_n- group may contain from one to three heteroatoms
 independently selected from nitrogen, sulfur, oxygen, >S(=O), >SO₂ or >NR⁶,
 wherein said (C₂-C₉)heterocycloalkyl moiety of said (C₂-C₉)heterocycloalkyl-
 (CH₂)_n- group may optionally be substituted on any of the ring carbon atoms
 20 capable of forming an additional bond (preferably one to three substituents per
 ring) with a substituent independently selected from the group consisting of
 hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy,
 (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 25 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 30 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,

NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 5 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl;

wherein R₄ is hydrogen, (C₁-C₆)alkyl, hydroxy, (C₁-C₆)alkoxy, hydroxy(C₁-
 C₆)alkyl, (C₁-C₆)alkoxy(C=O)-, (C₃-C₁₀)cycloalkyl-(CH₂)_q-,
 (C₂-C₉)heterocycloalkyl-(CH₂)_q-, (C₂-C₉)heteroaryl-(CH₂)_q-, phenyl-(CH₂)_q-, or
 10 naphthyl-(CH₂)_q-; wherein said (C₂-C₉)heterocycloalkyl, (C₂-C₉)heteroaryl,
 phenyl and naphthyl groups may be optionally substituted with one or two
 substituents from the group consisting of hydrogen, halo, cyano, (C₁-C₆)alkyl,
 hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-
 (C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-
 15 (C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-, (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-,
 H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂,
 amino, (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂ amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino (C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 20 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-
 , (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 25 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl;

wherein R₅ is hydrogen, (C₁-C₆)alkyl or amino; or

R₄ and R₅ together with the nitrogen atom to which they are attached
 form a (C₂-C₉)heterocycloalkyl group optionally substituted with one or two
 30 substituents selected from the group consisting of hydrogen, halo, cyano,
 (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy,
 (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)
 35 alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino, (C₁-C₆)alkylamino,

NH-, (C₁-C₆)alkyl(C=O)-NH-, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl HN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 5 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl;

or R₃ and the carbon to which it is attached form a five to seven
 membered carbocyclic ring, wherein any of the carbon atoms of said five
 membered carbocyclic ring may optionally be substituted with a substituent
 10 selected from the group consisting of hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy,
 hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-,
 (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-, (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-,
 H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂,
 15 amino, (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 NH-, (C₁-C₆)alkyl(C=O)-NH-, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 20 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl; wherein one of the carbon-carbon bonds of said five to seven
 25 membered carbocyclic ring may optionally be fused to an optionally substituted
 phenyl ring, wherein said substituents may be independently selected from
 hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy,
 (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 30 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 35 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-

"(C₂-C₉)heterocycloalkyl" when used herein refers to pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydropyranyl, pyranal, thiopyranal, aziridinyl, oxiranyl, methylenedioxy, chromenyl, isoxazolidinyl, 1,3-oxazolidin-3-yl, isothiazolidinyl, 1,3-thiazolidin-3-yl, 1,2-pyrazolidin-2-yl, 1,3-pyrazolidin-1-yl, 5 piperidinyl, thiomorpholinyl, 1,2-tetrahydrothiazin-2-yl, 1,3-tetrahydrothiazin-3-yl, tetrahydrothiadiazinyl, morpholinyl, 1,2-tetrahydrodiazin-2-yl, 1,3-tetrahydrodiazin-1-yl, tetrahydroazepinyl, piperazinyl, chromanyl, and the like. One of ordinary skill in the art will understand that the connection of said (C₂-C₉)heterocycloalkyl rings is through a carbon or a sp³ hybridized nitrogen 10 heteroatom.

"(C₂-C₉)heteroaryl" when used herein refers to furyl, thienyl, thiazolyl, pyrazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrrolyl, triazolyl, tetrazolyl, imidazolyl, 1,3,5-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,3-oxadiazolyl, 1,3,5-thiadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, pyridyl, pyrimidyl, pyrazinyl, 15 pyridazinyl, 1,2,4-triazinyl, 1,2,3-triazinyl, 1,3,5-triazinyl, pyrazolo[3,4-b]pyridinyl, cinnolinyl, pteridinyl, purinyl, 6,7-dihydro-5H-[1]pyrindinyl, benzo[b]thiophenyl, 5, 6, 7, 8-tetrahydro-quinolin-3-yl, benzoxazolyl, benzothiazolyl, benzisothiazolyl, benzisoxazolyl, benzimidazolyl, thianaphthenyl, isothianaphthenyl, benzofuranyl, isobenzofuranyl, isoindolyl, indolyl, indoliziny, 20 indazolyl, isoquinolyl, quinolyl, phthalazinyl, quinoxaliny, quinazoliny, benzoxazinyl, and the like. One of ordinary skill in the art will understand that the connection of said (C₂-C₉)heterocycloalkyl rings is through a carbon atom or a sp³ hybridized nitrogen heteroatom.

"Aryl" when used herein refers to phenyl or naphthyl.

25 "Protected amine" and "protected amino" refers to an amine group with one of the hydrogen atoms replaced with a protecting group (P). Any suitable protecting group may be used for amine protection. Suitable protecting groups include carbobenzyloxy, t-butoxy carbonyl (BOC) or 9-fluorenyl-methylenoxy carbonyl.

30 Compounds of Formula CCR1-I and their methods of manufacture are disclosed in commonly assigned United States Patent Application Serial No. 09/380,269, filed February 5, 1998, United States Patent Application Serial No. 09/403,218, filed January 18, 1999, PCT Publication No. WO98/38167, and PCT Publication No. WO99/40061, all of which are incorporated herein by 35 reference in their entireties for all purposes.

[(C₁-C₆)alkyl]₂ amino, amino(C₁-C₆)alkyl, (C₁-C₆)alkylamino (C₁-C₆)alkyl,
 [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-, (C₁-C₆)alkyl-NH-(C=O)-,
 [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-HN(C=O)-
 (C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-NH-,
 5 (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl, (C₁-C₆)alkyl(C=O)-
 [N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-, (C₁-C₆)alkyl-
 SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl, (C₁-C₆)alkylHN-
 SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-, (C₁-C₆)alkyl-SO₃-,
 phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and (C₂-C₉)heteroaryl;
 10 wherein R⁶ is hydrogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy-(CH₂)_g-,
 (C₁-C₆)alkoxy(C=O)-(CH₂)_g-, (C₁-C₆)alkyl-(SO₂)-(CH₂)_g-, (C₆-C₁₀)aryloxy-(CH₂)_g-,
 (C₆-C₁₀)aryloxy(C=O)-(CH₂)_g-, or (C₆-C₁₀)aryl-(SO₂)-(CH₂)_g-;

wherein g is an integer from zero to four;

wherein m is an integer from zero to four;

15 wherein n is an integer from zero to six;

with the proviso that when one of R⁴ or R⁵ is hydrogen, and the other of R⁴ or
 R⁵ is (C₁-C₆)alkyl; R² is (C₃-C₁₀)cycloalkyl or isopropyl and R³ is (C₃-C₅)alkyl,
 phenyl, methylvinyl, dimethylvinyl, halovinyl, hydroxy(C₁-C₃)alkyl or amino(C₁-
 C₄)alkyl then R¹ must be other than indol-5-yl, 6-azaindol-2-yl, 2,3-dichloro-
 20 pyrrol-5-yl, 4-hydroxyquinolin-3-yl, 2-hydroxyquinoxalin-3-yl, 6-azaindolin-3-yl,
 or optionally substituted indol-2 or 3-yl;
 and the pharmaceutically acceptable salts of such compounds.

Unless otherwise indicated, the alkyl and alkenyl groups referred to
 herein, as well as the alkyl moieties of other groups referred to herein (e.g.,
 25 alkoxy), may be linear or branched, and they may also be cyclic (e.g.,
 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl) or be linear or
 branched and contain cyclic moieties. Such alkyl and alkoxy groups may be
 substituted with one, two or three halogen and/or hydroxy atoms, preferably
 fluorine atoms.

30 Unless otherwise indicated, "halogen" includes fluorine, chlorine,
 bromine, and iodine.

"(C₃-C₁₀)cycloalkyl" when used herein refers to cycloalkyl groups
 containing zero to two levels of unsaturation such as cyclopropyl, cyclobutyl,
 cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, 1,3-cyclohexadiene,
 35 cycloheptyl, cycloheptenyl, bicyclo[3.2.1]octane, norbornanyl, and the like.

quinoxaline-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-7-fluoro-2(S)-hydroxy-7-methyl-octyl)-amide;

quinoxaline-2-carboxylic acid 1(S)-benzyl-5-cyclohexyl-2(S)-hydroxy-4(R)-methylcarbamoyl-pentyl)-amide;

5 quinoxaline-2-carboxylic acid 1(S)-cyclohexylmethyl-2(S)-hydroxy-7-methyl-4(R)-methylcarbamoyl-octyl)-amide;

quinoxaline-2-carboxylic acid [1(S)-benzyl-2(S)-hydroxy-4(S)-hydroxycarbamoyl-4-(1-hydroxy-4-methyl-cyclohexyl)-butyl]-amide;

10 quinoxaline-2-carboxylic acid [1(S)-benzyl-4(S)-(4,4-difluoro-1-hydroxy-cyclohexyl)-2(S)-hydroxy-4-hydroxycarbamoyl-butyl]-amide;

quinoxaline-2-carboxylic acid [1(S)-benzyl-4(S)-carbamoyl-4(S)-(4,4-difluoro-cyclohexyl)-2(S)-hydroxy-butyl]-amide;

quinoline-3-carboxylic acid (1(S)-benzyl-4(S)-carbamoyl-4-cyclohexyl-2(S)-hydroxy-butyl)-amide;

15 quinoxaline-2-carboxylic acid (4(R)-carbamoyl-2(S)-hydroxy-7-methyl-1(S)-thiophen-2-ylmethyl-octyl)-amide;

quinoxaline-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-7-chloro-2(S)-hydroxy-oct-6-enyl)-amide;

20 quinoxaline-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-2(S)-hydroxy-5-phenyl-pentyl)-amide;

N-1(S)-benzyl-4(R)-carbamoyl-7-fluoro-2(S)-hydroxy-7-methyl-octyl)-5,6-dichloro-nicotinamide;

quinoxaline-2-carboxylic acid (4(R)-carbamoyl-2(S)-hydroxy-7-methyl-1(S)-thiazol-4(R)-ylmethyl-octyl)-amide;

25 benzothiazole-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-7-fluoro-2(S)-hydroxy-7-methyl-octyl)-amide; and

benzofuran-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-7-fluoro-2(S)-hydroxy-7-methyl-octyl)-amide.


30 In another preferred embodiment, the CCR1 compound has a formula Ia-1:

In a preferred embodiment, the CCR1 inhibitor is selected from one of the following compounds of Formula CCR1-I:


- quinoxaline-2-carboxylic acid 4(R)-carbamoyl-1(S)-(3-chloro-benzyl)-
2(S),7-dihydroxy-7-methyl-octyl]-amide;
- 5 7,8-difluoro-quinoline-3-carboxylic acid (1S)-benzyl-4(R)-carbamoyl-
2(S),7-dihydroxy-7-methyl-octyl]-amide;
- 6,7,8-trifluoro-quinoline-3-carboxylic acid (1S)-benzyl-4(R)-carbamoyl-
2(S),7-dihydroxy-7-methyl-octyl]-amide;
- 10 quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-(3-fluoro-benzyl)-
2(S),7-dihydroxy-7-methyl-octyl]-amide;
- quinoxaline-2-carboxylic acid (1S)-benzyl-2(S),7-dihydroxy-4(R)-
hydroxycarbamoyl-7-methyl-octyl]-amide;
- quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-(2-chloro-benzyl)-
2(S),7-dihydroxy-7-methyl-octyl]-amide;
- 15 quinoxaline-2-carboxylic acid [1(S)-(2-fluoro-benzyl)-2(S),7-dihydroxy-
4(R)-hydroxycarbamoyl-7-methyl-octyl]-amide;
- quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-(2-fluoro-benzyl)-
2(S),7-dihydroxy-7-methyl-octyl]-amide;
- quinoxaline-2-carboxylic acid [1(S)-(3,4-difluoro-benzyl)-2(S),7-
20 dihydroxy-4(R)-hydroxycarbamoyl-7-methyl-octyl]-amide;
- quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-(3,4-difluoro-benzyl)-
2(S),7-dihydroxy-7-methyl-octyl]-amide;
- quinoxaline-2-carboxylic acid (4(R)-carbamoyl-2(S),7-dihydroxy-7-
methyl-1(S)-naphthalen-1-ylmethyl-octyl)-amide;
- 25 7,8-difluoro-quinoline-3-carboxylic acid 1(S)-benzyl-2(S)-hydroxy-7-
methyl-4(R)-methylcarbamoyl-octyl]-amide;
- 8-fluoro-quinoline-3-carboxylic acid 1(S)-benzyl-2(S)-hydroxy-7-methyl-
4(R)-methylcarbamoyl-octyl]-amide;
- quinoxaline-2-carboxylic acid [4(R)-carbamoyl-7-fluoro-1-(3(S)-fluoro-
30 benzyl)-2(S)-hydroxy-7-methyl-octyl]-amide;
- quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1-(2(S)-fluoro-benzyl)-
2(S)-hydroxy-7-methyl-octyl]-amide;
- quinoxaline-2-carboxylic acid [1(S)-benzyl-4(S)-carbamoyl-4(S)-(2,6-
dimethyl-tetrahydro-pyran-4-yl)-2(S)-hydroxy-butyl]-amide;

133

Scheme 1



132



wherein the substituents are as defined above.

In a preferred method of making the compound Ia-1, the reaction is started with Scheme 1. In the herein described processes, the substituents are as defined for CCR1-I, and the following:

R_7 is hydroxy, (C_1-C_6) alkyl, or phenyl wherein the phenyl group unsubstituted or substituted with one, two, or three (C_1-C_6) alkyl, hydroxy, or halogen groups;

R_8 is hydroxy or halogen;

R_9 is phenyl, naphthyl, (C_3-C_{10}) cycloalkyl, (C_1-C_6) alkyl or (C_2-C_9) heteroaryl, wherein each of said phenyl, naphthyl, (C_3-C_{10}) cycloalkyl or (C_2-C_9) heteroaryl groups may be unsubstituted or substituted with one, two, or three substituents independently selected from the group consisting of halogen, cyano, and (C_1-C_6) alkyl;

P is a protecting group;

X is hydroxy or halogen; and

q is 0, 1, 2, 3, or 4.

desired lactone in subsequent steps.

Step 5 of scheme 1 includes a) hydrolyzing a compound of the formula (VIg-1) with an aqueous solution in the presence of a base, b) protecting the amine group of the compound so formed, and c) cyclizing the compound so
5 formed with heat and an acid catalyst. In one embodiment, the compound VIg-1 is hydrolyzed with sodium hydroxide. The pH is adjusted to approximately 10 and tetrahydrofuran and BOC dicarbonate are added. This provides the protected hydroxy acid, which may be heated in 10% acetic acid and toluene to provide the protected amine lactone (V-1).

10 The compound of formula (V-1) may also be produced according to scheme 2.

In scheme 1 step 1, a compound of the formula (VI-1) is reduced with a reducing agent under heat to form a compound of the formula (VId-1). In one embodiment, the reducing agent is aluminum triisopropoxide and isopropanol. Preferably, the temperature is maintained above room temperature, more preferably between about 60°C and about 82°C. The product alcohol can be isolated by either cooling the reaction mixture to room temperature, diluting with more isopropanol and collecting the crystalline material or by cooling the reaction to room temperature and adding 1N HCL and water and collecting the crystalline material.

Step 2 of scheme 1 includes reacting a compound of the formula R_7-SO_2-X and a compound of the formula (VId-1) in the presence of a base to form the compound of the formula (VIe-1). Any amine base is suitable, including pyridine, triethylamine, N-methylmorpholine, and diisopropylethylamine. In one embodiment, $R_7-SO_2-R_8$ is p-toluenesulfonic acid, methanesulfonic acid, sulfuric acid, or methanesulfonyl chloride. In another embodiment, the conversion of hydroxy dioxane (VId-1) to dioxane oxazolidinone (VIe-1) can be achieved by treatment of the hydroxy dioxane (VId-1) with methanesulfonyl chloride and triethylamine in tetrahydrofuran solution and heating the mixture to cause the cyclization of the mesylate formed in situ to the oxazolidinone.

In step 3 of scheme 1, a compound of the formula (VIe-1) may be formed by heating the compound of the formula (VIe-1). The reaction may proceed by dissolving compound VIe-1 in a solvent such as pyridine or N-methyl imidazole and heating the mixture for several hours at temperature from about 50°C to about 100°C; preferably at about 80°C. The mesylate (VIe-1) may be recovered by extraction into an organic solvent such as ethyl acetate and removal of the amine solvents by extraction of the solution with aqueous acid.

Step 4 of scheme 1 depicts reacting hydroxylamine hydrochloride, a compound of the formula R_7-SO_2-X , and a compound of the formula (VIe-1) to form a compound of the formula (VIg-1). In one embodiment, R_7-SO_2-X is p-toluenesulfonic acid, methanesulfonic acid, sulfuric acid, or methanesulfonyl chloride. The reaction may occur in a solvent, such as methanol. In one embodiment, the reaction occurs in methanol with tosic acid at reflux for 8 to 24 hours. The resulting nitrile oxazolidinone contains a small amount of the corresponding ethyl ester which is not removed since it also is converted to the

Alternatively, in step 1 of scheme 2, the compound of the formula (VIa-1) may be formed by reacting hypochlorous acid and a compound of the formula (VI-1). Such an oxidation reaction typically produces chlorinated forms of the compound VIa-1 as side products in addition to the compound VIa-1. This
5 oxidation reaction proceeds by mixing the compound VI-1 in solvent, such as acetic acid and/or acetone, and adding sodium hypochlorite, while keeping the mixture at a low temperature, preferably at or below about 0°C.


As a means to convert the side product chlorinated forms of the compound VIa-1 to compounds of the formula V-1, the compounds formed from
10 the hypochlorous acid oxidation reaction may optionally be hydrogenated by reaction with hydrogen in the presence of a catalyst. The hydrogenation may include introducing the products from the hypochlorous acid oxidation reaction into a solvent system of tetrahydrofuran and water, followed by addition of a Pd/C catalyst. The resulting mixture is subjected to hydrogen above
15 atmospheric pressure and temperature. In one embodiment, the pressure is about 80 pounds per square inch and the temperature is maintained from about 60°C to about 70°C until the reaction is substantially complete.

In step 2 of scheme 2, the compound of the formula (VIb-1) may be formed by reacting a silyating agent and a compound of the formula (VIa-1) and
20 reacting the compound so formed with a reducing agent. In one embodiment, the reducing agent is N-selectride. In another embodiment, the silyating agent is 1,1,1,3,3,3-hexamethyl-disilazane. The reduction reaction may occur at temperatures below about 0°C, preferably below about -20°C, more preferably below about -50°C. In addition, the reducing agent may be present in slight
25 excess.

In step 3 of scheme 2, the compound of the formula (V-1) is formed by heating a compound of the formula (VIb-1) in the presence of an acid catalyst, such as acetic acid. In one embodiment, the cyclization reaction occurs by
30 introducing the compound VIb-1 into a solvent mixture, such as toluene and 10% acetic acid, at the solvent reflux temperature for 8 to 16 hours. This provides the desired lactone as a crystalline solid after work up.

One method of making the compound of the formula (VI-1) is by reacting a compound of the formula (VII-1)


Scheme 2



In step 1 of scheme 2, a compound of the formula (VI-1) may be reacted with ozone to form a compound of the formula (VIa-1). The compound VI-1 may be present in a solvent, such as ethyl acetate, and the ozone introduced through sparging at a temperature below room temperature, preferably at about -15°C, until the starting dioxane ketone is substantially reacted. Any excess ozone may be removed by bubbling nitrogen through the solution. The resulting crude ketone ester mixture may be isolated after treatment with aqueous sodium bisulfite to remove any hydroperoxides.

139


Scheme 3



In step 1 of scheme 3, the compound of the formula (IVa1-1) may be formed by reacting 4-halo-2-methyl-2-butene and a compound of the formula (V-1) in the presence of a base. Exemplary bases include lithium dialkyl amides such as lithium N-isopropyl-N-cyclohexylamide, lithium bis(trimethylsilyl)amide, lithium di-isopropylamide, and potassium hydride. Suitable solvents include aprotic polar solvents such as ethers (such as tetrahydrofuran, glyme or dioxane), benzene, or toluene, preferably tetrahydrofuran. The aforesaid reaction is conducted at a temperature from about -78°C to about 0°C, preferably at about -78°C. In one embodiment, alkylation of the lactone (V-1) is accomplished by reacting the lactone (V-1) with lithium bis(trimethylsilyl)amide and dimethylallyl bromide in tetrahydrofuran at a temperature from about -78°C to about -50°C. Reaction times range from several hours or if an additive such as dimethyl imidazolidinone is present, the reaction may be complete in minutes.

Compounds of formula (IVa1-1) may be used to produce compounds of the formula (Ia-1) according to scheme 4:


138



(VII-1)

with a Grignard reagent formed in situ by addition of 2-(2-bromo-ethyl)-[1,3]dioxane to a mixture comprising magnesium and the compound of the formula (VII-1). In one embodiment, the mixture further comprises methyl
 5 magnesium chloride and/or methyl magnesium bromide in a solvent. Any exotherm formed from the reaction may be controlled by the rate of addition of the bromide.

The compound of the formula (VII-1) may be formed by coupling N,O-dimethylhydroxylamine hydrochloride and a compound of the formula (VIII-1)



(VIII-1)

10 This coupling reaction may be performed by mixed anhydride procedure. In one mixed anhydride procedure, compound VIII-1 is combined with methylene chloride and N-methylmorpholine is added followed by isobutyl chloroformate. In a separate mixture, a slurry of N,O-dimethylhydroxylamine hydrochloride is
 15 treated with N-methylmorpholine. The two reaction mixtures are combined and then quenched with a solution of citric acid in water. This procedure preferably operates at a temperature below about 20°C, more preferably below about 0°C.

Compounds of formula (V-1) may be used to produce compounds of the formula (IVa1-1) according to scheme 3:

20

dependent on temperature and batch size, amount other factors, but typically reaction time is from about 2 hours to about 14 hours.


Step 2 of scheme 4 depicts coupling a compound IIIa1-1 with a compound having the formula $R_1\text{-CO-X}$ to form a compound having the formula (IIa1-1). This coupling reaction is generally conducted at a temperature from about -30°C to about 80°C , preferably from about 0°C to about 25°C . The coupling reaction may occur with a coupling reagent that activates the acid functionality. Exemplary coupling reagents include dicyclohexylcarbodiimide/hydroxybenzotriazole (DCC/HBT), N-3-dimethylaminopropyl-N'-ethylcarbodiimide (EDC/HBT), 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), carbonyl diimidazole (CDI), and diethylphosphorylcyanide. The coupling is conducted in an inert solvent, preferably an aprotic solvent, such as tetrahydrofuran, acetonitrile, dichloromethane, chloroform, or N,N-dimethylformamide. One preferred solvent is tetrahydrofuran. In one embodiment, quinoxaline acid is combined with CDI in anhydrous tetrahydrofuran and heated to provide the acyl imidazole. Compound IIIa1-1 is added to the acyl imidazole at room temperature to form the compound IIa1-1.

Step 3 of scheme 4 includes reacting the compound of formula IIa1-1 with an amine having a formula NHR_4R_5 to form a compound of the formula (Ia-1). In one embodiment, the amine is ammonia either anhydrous in an organic solvent or as an aqueous solution of ammonium hydroxide added to a polar solvent at a temperature from about -10°C to about 35°C , preferably at about 30°C . Suitable solvents include, alcohols, such as methanol, ethanol, or butanols; ethers such as tetrahydrofuran, glyme or dioxane; or a mixture thereof, including aqueous mixtures. Preferably the solvent is methanol. In one embodiment, the compound IIa1-1 is dissolved in methanol which has been saturated with ammonia gas. In another embodiment, the compound IIa1-1 in methanol is treated with ammonium hydroxide in tetrahydrofuran at room temperature.

Scheme 5 represents an alternative method to form compounds of formula Ia-1 from compounds of formula IVa1-1.

140

Scheme 4



In step 1 of scheme 4, a compound of the formula (IIIa1-1) is formed by reacting a compound of the formula (IVa1-1) with phosphoric acid. Preferably, this reaction occurs in any suitable solvent, such as non-alcoholic solvents. Two preferred solvents include tetrahydrofuran and dichloroethane. The reaction may take place at any suitable temperature, preferably from about -25°C to about 120°C, more preferably from about 15°C to about 40°C. Reaction time is


In step 1 of scheme 5, a compound of the formula (IVa1-1) is reacted with a compound of the formula $R_9\text{-SO}_2\text{-X}$ to form a compound of the formula (IVa2-1). Any suitable acidic deprotection reaction may be performed. In one example, an excess of p-toluenesulfonic acid hydrate in ethyl acetate is introduced to the compound IVa1-1 at room temperature. Suitable solvents include ethyl acetate, alcohols, tetrahydrofuran, and mixtures thereof. The reaction may proceed at ambient or elevated temperatures. Typically, the reaction is substantially complete within two and twelve hours. The resulting compound IVa2-1 may be crystallized and separated from the reaction mixture, and may be further purified to remove impurities by recrystallization from hot ethyl acetate.

In step 2 of scheme 5, the compound IVa2-1 may be coupled with a compound having the formula $R_1\text{-CO-X}$ to form a compound of the formula (IIIa2-1). This coupling reaction is generally conducted at a temperature from about -30°C to about 80°C, preferably from about 0°C to about 25°C. The coupling reaction may occur with a coupling reagent that activates the acid functionality. Exemplary coupling reagents include dicyclohexylcarbodiimide/hydroxybenzotriazole (DCC/HBT), N-3-dimethylaminopropyl-N'-ethylcarbodiimide (EDC/HBT), 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), carbonyl diimidazole (CDI)/dimethylaminopyridine (DMAP), and diethylphosphorylcyanoide. The coupling is conducted in an inert solvent, preferably an aprotic solvent, such as acetonitrile, dichloromethane, chloroform, or N,N-dimethylformamide. One preferred solvent is methylene chloride. In one embodiment, quinoxaline acid is combined with methylene chloride, oxalyl chloride and a catalytic amount of N,N-dimethylformamide to form an acid chloride complex. The compound IVa2-1 is added to the acid chloride complex followed by triethylamine at a temperature from about 0°C to about 25°C to form the compound IIIa2-1.

Step 3 of scheme 5 includes reacting a compound IIIa2-1 with trifluoroacetic acid to produce a compound of the formula (IIa2-1). In one embodiment, the hydration with trifluoroacetic acid occurs in methylene chloride solution at room temperature. The hydration may take several hours to complete at room temperature. A catalytic amount of sulfuric acid can be added to the reaction solution to increase the rate of reaction.

142

Scheme 5



those that are at least about 10% ionized over at least a portion of the physiologically relevant pH 1 to 8 range and thus such groups have pK_a values of about 0 to 9.

5 The neutral dispersion polymer may be cellulosic or non-cellulosic. A preferred class of neutral cellulosic dispersion polymers are those with at least one ester- and/or ether- linked substituent in which the polymer has a degree of substitution of at least 0.05 for each substituent. It should be noted that in the polymer nomenclature used herein, ether-linked substituents are recited prior to "cellulose" as the moiety attached to the ether group; for example, "methyl
10 cellulose" has a methyl moiety ether-linked to the polymer. In contrast, ester-linked substituents are recited after "cellulose" as the carboxylate; for example, "cellulose acetate" has an acetate moiety ester-linked to the polymer.

It should also be noted that a polymer name such as "cellulose acetate butyrate" refers to any of the family of cellulosic polymers that have acetate and
15 butyrate groups attached via ester linkages to a significant fraction of the cellulosic polymer's hydroxyl groups. Generally, the degree of substitution of each substituent group can range from about 0.05 to 2.9 as long as the other criteria of the polymer are met. "Degree of substitution" refers to the average number of the three hydroxyls per saccharide repeat unit on the cellulose chain that have been substituted. For example, if all of the hydroxyls on the cellulose
20 chain have been butyrate substituted, the butyrate degree of substitution is 3. Also included within each polymer family type are cellulosic polymers that have additional substituents added in relatively small amounts that do not substantially alter the performance of the polymer.

25 Polymer substituents may be either non-ionizable or ionizable; however, the ionizable groups may not be acidic groups. Exemplary ether-linked non-ionizable substituents include: alkyl groups, such as methyl, ethyl, propyl, butyl, etc.; hydroxy alkyl groups such as hydroxymethyl, hydroxyethyl, hydroxypropyl, etc.; and aryl groups such as phenyl. Exemplary ester-linked non-ionizable
30 groups include: alkylate groups, such as acetate, propionate, butyrate, etc.; and arylate groups such as phenylate. However, when ester-linked non-ionizable groups are included, the polymer may need to include a sufficient amount of a hydrophilic substituent so that the polymer has at least some water solubility at any physiologically relevant pH of from 1 to 8. In the case of neutral dispersion
35 polymers, such hydrophilic substituents may consist of non-acidic ionizable

Step 4 of scheme 5 includes reacting the compound of formula IIa2-1 with an amine having a formula NHR_4R_5 to form a compound of the formula (Ia-1). In one embodiment, the amine is ammonia either anhydrous in an organic solvent or as an aqueous solution of ammonium hydroxide added to a polar solvent at a temperature from about -10°C to about 35°C , preferably at about 30°C . Suitable solvents include, alcohols, such as methanol, ethanol, or butanols; ethers such as tetrahydrofuran, glyme or dioxane; or a mixture thereof, including aqueous mixtures. Preferably the solvent is methanol. In one embodiment, the compound IIa2-1 is dissolved in methanol which has been saturated with ammonia gas. In another embodiment, the compound IIa2-1 in methanol is treated with ammonium hydroxide in tetrahydrofuran at room temperature.

NEUTRAL DISPERSION POLYMERS

The neutral polymers suitable for use in the dispersions of the present invention are pharmaceutically acceptable and should be inert, in the sense that they do not chemically react with the drug in an adverse manner when present in the composition. The polymer should also have an aqueous-solubility of at least 0.1 mg/mL over at least a portion of the pH range of 1-8.

The polymer is "neutral," meaning that the dispersion polymer possesses substantially no acidic functional groups. By "substantially no acidic functional groups" is meant that the number of acidic groups covalently attached to the polymer is less than about 0.05 milliequivalents per gram of polymer. Preferably, the number is less than about 0.02 milliequivalents per gram of polymer. By "acidic groups" is meant functional groups that, when attached to the polymer, have pK_a values in a humid or aqueous environment of about 5 or less. Preferably, the pK_a value of the functional groups on the neutral polymer is greater than about 6. Thus, the neutral polymers may contain ionic groups as long as the groups are not acidic.

However, for many drugs, particularly those that are also sensitive to basic conditions, e.g., "base-sensitive drugs," it is preferred that the neutral polymers be substantially nonionizable. By "substantially nonionizable" is meant that the number of "ionizable groups" covalently attached to the polymer is less than about 0.05 milliequivalents per gram of polymer and preferably less than about 0.02 milliequivalents per gram of polymer. "Ionizable groups" are

groups such as methyl, ethyl, propyl, butyl, etc.; or ester-linked alkyl groups such as acetate, propionate, butyrate, etc.; and ether- and/or ester-linked aryl groups such as phenyl, benzoate, or phenylate. As discussed above, hydrophilic substituents may consist of non-acidic ionizable groups such as amino-functionalized groups or phenolate groups. In the case of non-ionizable neutral dispersion polymers, such hydrophilic groups are non-ionizable substituents such as alcohol, ether or ester groups.

Exemplary amphiphilic polymers include non-ionizable cellulosics such as hydroxypropyl methyl cellulose, hydroxyethyl methyl cellulose and hydroxyethyl acetate; non-acidic ionizable cellulosics such as amino ethyl cellulose acetate and hydroxybenzyl cellulose; and non-ionizable non-cellulosics such as polyvinylpyrrolidone ethylene/vinyl alcohol copolymers and polyoxyethylene-polyoxypropylene block copolymers (also referred to as poloxamers); and ionizable non-cellulosics such as amine-functionalized polyacrylates and polymethacrylates. When the amphiphilic polymer is hydroxypropyl methyl cellulose, it is preferred that it is provided in a grade that possesses a dynamic viscosity that is 60 mPa-s or less, more preferably 10 mPa-s or less and most preferably between 1 mPa-s and 6 mPa-s, when measured in a 2% weight/volume solution at 20°C.

A preferred class of neutral non-cellulosic polymers are comprised of vinyl copolymers of a hydrophilic, hydroxyl-containing repeat unit and a hydrophobic, alkyl- or aryl-containing repeat unit. Such neutral vinyl copolymers are termed "amphiphilic hydroxyl-functional vinyl copolymers." Amphiphilic hydroxyl-functional vinyl copolymers are exceptional in that they are both non-ionic and yet, surprisingly, when used as dispersion polymers for low-solubility drugs, yield solid amorphous dispersions that provide high levels of drug concentration enhancement when dosed to an aqueous environment of use. Such polymers may be used with any low-solubility drug, and not simply acid-sensitive drugs.

Dispersions of such copolymers are believed to provide high concentration enhancements due to the amphiphilicity of these copolymers which provide both sufficient hydrophobic groups to interact with the hydrophobic, low-solubility drugs and also sufficient hydrophilic groups to have sufficient aqueous solubility for good dissolution. The copolymeric structure of the amphiphilic hydroxyl-functional vinyl copolymers also allows their

groups such as amino-functionalized groups or phenolate groups. In the case of non-ionizable neutral dispersion polymers, such hydrophilic groups are non-ionizable substituents such as alcohol, ether or ester groups.

Exemplary neutral non-ionizable cellulosic polymers that may be used to form the dispersion include: hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and hydroxyethyl ethyl cellulose.

Exemplary neutral, but ionizable cellulosic polymers, include aminoethyl cellulose, aminoethyl cellulose acetate, hydroxypropyl amino ethyl cellulose and hydroxybenzyl cellulose.

Another class of neutral dispersion polymers is non-cellulosic, neutral polymers. Such polymers may be either non-ionizable or ionizable. Exemplary non-ionizable, neutral polymers include vinyl polymers and copolymers having substituents of hydroxyl, alkylacyloxy, and cyclicamido. Exemplary non-cellulosic, neutral polymers include hydroxyethyl methacrylate, polyvinylhydroxyethyl ether, polyethylene glycol, and polyoxyethylene-polyoxypropylene block copolymers also known as poloxamers.

Exemplary ionizable neutral polymers include amine-functionalized polyacrylates and polymethacrylates, some of which are also sold as EUDRAGITS manufactured by Rohm Tech Inc., and neutral proteins.

A preferred subset of neutral polymers are those that are also concentration-enhancing. Dispersion polymers that are both neutral and also concentration-enhancing are generally amphiphilic in that they possess substituents that are relatively hydrophobic and substituents that are relatively hydrophilic.

Amphiphilic cellulose may be prepared by substituting the cellulose at any or all of the 3 hydroxyl substituents present on each saccharide repeat unit with at least one relatively hydrophobic substituent. Hydrophobic substituents may be essentially any substituent that, if substituted to a high enough level or degree of substitution, can render the cellulosic polymer essentially aqueous insoluble. Hydrophilic regions of the polymer can be either those portions that are relatively unsubstituted, since the unsubstituted hydroxyls are themselves relatively hydrophilic, or those regions that are substituted with hydrophilic substituents. Examples of hydrophobic substituents include ether-linked alkyl

copoly- mers may be synthesized by first forming the homopolymer of the hydrophobic vinyl repeat unit having the substi- tuent B, followed by hydrolysis of a portion of the ester groups to convert a portion of the hydrophobic repeat units to hydrophilic, hydroxyl-containing repeat units having the substituent A.

5 For example, partial hydrolysis of the homopolymer, polyvinylbutyrate, yields the copolymer, vinylalcohol/vinylbutyrate copolymer for which A is hydroxyl (-OH) and B is butyrate (-OOC-CH₂.CH₂.CH₃).

For all types of copolymers, the value of n must be sufficiently large relative to the value of m that the resulting copolymer is at least partially water
10 soluble. Although the value of the ratio, n/m varies depending on the identity of A and B, it is generally at least about 1 and more commonly about 2 or more. The ratio n/m can be as high as 200. When the copolymer is formed by hydrolysis of the hydrophobic homopolymer, the relative values of n and m are typically reported in "percent hydrolysis," which is the fraction (expressed as a
15 percent) of the total repeat units of the copolymer that are in the hydrolyzed or hydroxyl form. The percent hydrolysis, H, is given as

$$\left(\frac{n}{n+m} \right)$$

$$H = 100 *$$


Thus, vinylbutyrate/vinylalcohol copolymer (formed by hydrolysis of a portion of the butyrate groups) having a percent hydrolysis of 75% has an n/m ratio of 3.

20 A particularly preferred family of amphiphilic hydroxyl-functional vinyl copolymers are those where A is hydroxyl and B is acetate. Such copolymers are termed vinylacetate/vinylalcohol copolymers. Some commercial grades are also sometimes referred to simply as polyvinylalcohol. However, the true homopolymer, polyvinylalcohol is not amphiphilic, is almost entirely water
25 insoluble, and is not a part of this invention. Preferred vinylacetate/vinylalcohol copolymers are those where H is between about 67% and 99.5%, or n/m has a value between about 2 and 200. The preferred average molecular weight is between about 2500 and 1,000,000 daltons and more preferably between about 3000 and about 100,000 daltons.

30 In order to form dispersions that have a high level of concentration and bioavailability enhancement, it is important that the identity of A and B, the n/m value, and average molecular weight be chosen such that the polymers are sufficiently hydrophobic that they strongly interact with the hydrophobic, low-solubility drug and yet are still significantly water soluble. Typically good results

hydrophilicity and hydrophobicity to be adjusted to maximize performance with a specific low-solubility drug.

The preferred copolymers have the general structure:



wherein A and B represent "hydrophilic, hydroxyl-containing" and "hydrophobic" substituents, respectively, and n and m represent the average number of hydrophilic vinyl repeat units and average number of hydrophobic vinyl repeat units respectively per polymer molecule. Copolymers may be block copolymers, random copolymers or they may have structures anywhere between these two extremes. The sum of n and m is generally from about 50 to about 20,000 and therefore the polymers have molecular weights from about 2,500 to about 1,000,000 daltons.

The hydrophilic, hydroxyl-containing repeat units, "A," may simply be hydroxyl (-OH) or it may be any short-chain, 1 to 6 carbon, alkyl with one or more hydroxyls attached thereto. The hydroxyl-substituted alkyl may be attached to the vinyl backbone via carbon-carbon or ether linkages. Thus, exemplary "A" structures include, in addition to hydroxyl itself, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxymethoxy, hydroxyethoxy and hydroxypropoxy.

The hydrophobic substituent, "B," may simply be: hydrogen (-H), in which case the hydrophobic repeat unit is ethylene; an alkyl or aryl substituent with up to 12 carbons attached via a carbon-carbon bond such as methyl, ethyl or phenyl; an alkyl or aryl substituent with up to 12 carbons attached via an ether linkage such as methoxy, ethoxy or phenoxy; an alkyl or aryl substituent with up to 12 carbons attached via an ester linkage such as acetate, propionate, butyrate or benzoate. The amphiphilic hydroxyl-functional vinyl copolymers of the present invention may be synthesized by any conventional method used to prepare substituted vinyl copolymers. Some substituted vinyl copolymers such as polyvinyl alcohol/polyvinyl acetate are well known and commercially available.

A particularly convenient subclass of amphiphilic hydroxyl-functional vinyl copolymers to synthesize are those where the hydrophobic substituent "B" comprises the hydrophilic substituent "A" to which an alkylate or arylate group is attached via an ester linkage to one or more of the hydroxyls of A. Such

(MDC) that is at least 1.25-fold the MDC provided by a control composition. In other words, if the MDC provided by the control composition is 100 μ g/mL, then a composition of the present invention containing a concentration-enhancing polymer provides an MDC of at least 125 μ g/mL. More preferably, the MDC of drug achieved with the compositions of the present invention are at least 2-fold, even more preferably at least 3-fold, and most preferably at least 10-fold that of the control composition.

The appropriate control composition depends on whether a dispersion polymer is being tested, or whether a composition comprised of a mixture of a dispersion and concentration-enhancing polymer is being tested. Where the composition is simply the dispersion itself, and the dispersion polymer is being tested to determine whether it is a concentration-enhancing polymer, the control composition is conventionally the undispersed drug alone (e.g., typically, the crystalline drug alone in its most thermodynamically stable crystalline form, or in cases where a crystalline form of the drug is unknown, the control may be the amorphous drug alone) or the drug plus a weight of inert diluent equivalent to the weight of polymer in the test composition. By inert is meant that the diluent is not concentration-enhancing. Where the composition is comprised of a mixture of a dispersion and additional concentration-enhancing polymer, the control composition is the dispersion alone without any additional concentration-enhancing polymer.

Alternatively, the compositions containing concentration-enhancing polymers of the present invention provide in an aqueous use environment a concentration versus time Area Under The Curve (AUC), for any period of at least 90 minutes between the time of introduction into the use environment and about 270 minutes following introduction to the use environment that is at least 1.25-fold that of an appropriate control composition. More preferably, the AUC achieved with the compositions of the present invention are at least 2-fold, more preferably at least 3-fold, and most preferably at least 10-fold that of a control composition.

Alternatively, the compositions of the present invention containing concentration-enhancing polymers, when dosed orally to a human or other animal, provide an AUC in drug concentration in the blood plasma or serum that is at least 1.25-fold that observed when an appropriate control composition is dosed. Preferably, the blood AUC is at least 2-fold, more preferably at least

are obtained when the aqueous solubility of the copolymer at 37°C is between about 0.04 and 10.0 wt% and preferably between about 0.1 and 2.0 wt%.

While specific polymers are discussed as being suitable for use in the compositions of the present invention, blends of such polymers may also be suitable. Thus the term "polymer" is intended to include blends of polymers in addition to a single species of polymer.

CONCENTRATION ENHANCEMENT

As described above, it is often preferred that the neutral dispersion polymer is also concentration-enhancing. The neutral concentration-enhancing dispersion polymers improve the concentration of a low-solubility drug in a use environment, and thereby preferably improve bioavailability of the drug.

The term "concentration-enhancing" means that the polymer is present in a sufficient amount in the composition so as to improve the concentration of the drug in a use environment relative to a control composition free from the concentration-enhancing polymer. As used herein, a "use environment" can be either the *in vivo* environment of the GI tract, subdermal, intranasal, buccal, intrathecal, ocular, intraaural, subcutaneous spaces, vaginal tract, arterial and venous blood vessels, pulmonary tract or intramuscular tissue of an animal, such as a mammal and particularly a human, or the *in vitro* environment of a test solution, such as phosphate buffered saline (PSB) or a Model Fasted Duodenal (MFD) solution. Concentration enhancement may be determined through either *in vitro* dissolution tests or through *in vivo* tests. It has been determined that enhanced drug concentration in *in vitro* dissolution tests in Model Fasted Duodenal (MFD) solution or Phosphate Buffered Saline (PBS) is a good indicator of *in vivo* performance and bioavailability. An appropriate PBS solution is an aqueous solution comprising 20 mM sodium phosphate (Na_2HPO_4), 47 mM potassium phosphate (KH_2PO_4), 87 mM NaCl, and 0.2 mM KCl, adjusted to pH 6.5 with NaOH. An appropriate MFD solution is the same PBS solution wherein additionally is present 7.3 mM sodium taurocholic acid and 1.4 mM of 1-palmitoyl-2-oleyl-sn-glycero-3-phosphocholine. In particular, a composition containing a concentration-enhancing polymer may be dissolution-tested by adding it to MFD or PBS solution and agitating to promote dissolution.

In one aspect, a composition containing a concentration-enhancing polymer of the present invention provides a Maximum Drug Concentration

that have submicron dimensions such as drug aggregates, aggregates of mixtures of polymer and drug, micelles, polymeric micelles, colloidal particles or nanocrystals, polymer/drug complexes, and other such drug-containing species that are present in the filtrate or supernatant in the specified dissolution test.

5 Alternatively, the concentration-enhancing polymer results in improved bioavailability. Relative bioavailability of the drug in the compositions of the present invention can be tested *in vivo* in animals or humans using conventional methods for making such a determination. An *in vivo* test, such as a crossover study, may be used to determine whether a composition provides an enhanced
10 relative bioavailability compared with a control. In an *in vivo* crossover study a "test composition" of drug and concentration-enhancing polymer is dosed to half a group of test subjects and, after an appropriate washout period (e.g., one week) the same subjects are dosed with a "control composition" that comprises an equivalent quantity of drug as the "test composition." The other half of the
15 group is dosed with the control composition first, followed by the test composition. The relative bioavailability is measured as the concentration in the blood (serum or plasma) versus time area under the curve (AUC) determined for the test composition divided by the AUC in the blood provided by the control composition. Preferably, this test/control ratio is determined for each subject, and then the ratios are averaged over all subjects in the study. *In vivo*
20 determinations of AUC can be made by plotting the serum or plasma concentration of drug along the ordinate (y-axis) against time along the abscissa (x-axis). Generally, the values for AUC represent a number of values taken from all of the subjects in a patient test population averaged over the entire test
25 population.

A preferred embodiment of the invention is one in which the relative bioavailability of the test composition is at least 1.25 relative to a control composition as described above. (That is, the AUC in the blood provided by the test composition is at least 1.25-fold the AUC provided by the control
30 composition.) An even more preferred embodiment of the invention is one in which the relative bioavailability of the test composition is at least 2.0, more preferably at least 3, relative to a control composition of the drug but with no polymer present, as described above. The determination of AUCs is a well-known procedure and is described, for example, in Welling, "Pharmacokinetics
35 Processes and Mathematics," ACS Monograph 185 (1986).

3-fold, that of an appropriate control composition. Thus, the compositions that, when evaluated, meet either the *in vitro* or the *in vivo*, or both, performance criteria are a part of this invention.

5 A typical *in vitro* test to evaluate enhanced drug concentration can be conducted by (1) administering with agitation a sufficient quantity of test composition (for example, the dispersion of the low-solubility drug and neutral polymer) in a test medium, such that if all of the drug dissolved, the theoretical concentration of drug would exceed the equilibrium concentration of the drug by a factor of at least 2; (2) adding an appropriate amount of control composition to
10 an equivalent amount of test medium; and (3) determining whether the measured MDC and/or AUC of the test composition in the test medium is at least 1.25-fold that provided by the control composition. The concentration of dissolved drug is typically measured as a function of time by sampling the test medium and plotting drug concentration in the test medium vs. time so that the
15 MDC and/or AUC can be ascertained. In conducting such a dissolution test, the amount of test composition or control composition used is an amount such that if all of the drug dissolved, the drug concentration would be at least 2-fold to 100-fold that of the solubility (that is, the equilibrium concentration), of the drug. For some dispersions of a very low-solubility drug and neutral polymer, it may
20 be necessary to administer an even greater amount of the dispersion to determine the MDC.

To avoid drug particulates which would give an erroneous determination, the test solution is either filtered or centrifuged. "Dissolved drug" is typically taken as that material that either passes a 0.45 μm syringe filter or,
25 alternatively, the material that remains in the supernatant following centrifugation. Filtration can be conducted using a 13 mm, 0.45 μm polyvinylidene difluoride syringe filter sold by Scientific Resources under the trademark TITAN®. Centrifugation is typically carried out in a polypropylene microcentrifuge tube by centrifuging at 13,000 G for 60 seconds. Other similar
30 filtration or centrifugation methods can be employed and useful results obtained. For example, using other types of microfilters may yield values somewhat higher or lower (± 10 -40%) than that obtained with the filter specified above but will still allow identification of preferred dispersions. It is recognized that this definition of "dissolved drug" encompasses not only monomeric solvated drug
35 molecules but also a wide range of species such as polymer/drug assemblies

example, where the degree of degradation of a drug in a test composition comprised of an acid-sensitive drug and a neutral polymer is 1 wt%, and the degree of degradation of a control composition comprised of an equivalent quantity of drug and HPMCAS is 5 wt%, the relative degree of improvement is 5 wt%/1 wt% equals 5.0. For dispersions of acid-sensitive drugs and neutral polymers of the present invention, the relative degree of improvement is at least 1.25. Preferably, the relative degree of improvement is at least 3, more preferably at least 5, and even more preferably at least 10. In fact, some compositions of the present invention may achieve a relative degree of improvement greater than 100.

The particular storage conditions and time of storage may be chosen as convenient depending on the degree of acid-sensitivity of the drug, the particular acidic polymer used in the control dispersion, and the ratio of drug to polymer in the dispersion. Where the drug is particularly acid-sensitive, or where the dispersion has a low ratio of drug to polymer, then shorter storage time periods may be used. Where the rate of drug degradation is linear, the relative degree of improvement will be independent of the storage time. However, where the rate of drug degradation is non-linear under controlled storage conditions, the stability test used to compare the test dispersion with the control dispersion is preferably chosen such that the degree of degradation is sufficiently large that it may be accurately measured. Typically, the time period is chosen so as to observe a degree of degradation in the control composition of at least 0.1 wt% to 0.2 wt%. However, the time period is not so long that the ratio of drug to polymer changes substantially. Typically, the time period is such that the observed degree of degradation for the test composition is less than 50 wt% and preferably less than 20 wt%. When rate of drug degradation in the control composition is relatively slow, the test is preferably conducted over a long enough period of time under controlled storage conditions to allow a meaningful comparison of the stability of the test dispersion with the control dispersion.

A stability test which may be used to test whether a dispersion meets the chemical stability criteria described above is storage of the test dispersion and the control dispersion for six months at 40°C and 75% RH. A relative degree of improvement may become apparent within a shorter time, such as three to five days, and shorter storage times may be used for some very acid-sensitive

Often the enhancement in drug concentration or relative bioavailability that is observed increases as the drug:polymer ratio decreases from a value of about 1 (50 wt% drug) to a value of about 0.11 (10 wt% drug). The drug:polymer ratio that yields optimum results varies from drug to drug and is best determined in *in vitro* dissolution tests and/or *in vivo* bioavailability tests. However, the amount of polymer that can be used in a dosage form is often limited by the total mass requirements of the dosage form. For example, when oral dosing to a human is desired, at low drug-to-polymer ratios the total mass of drug and polymer may be unacceptably large for delivery of the desired dose in a single tablet or capsule. Thus, it is often necessary to use drug-to-polymer ratios that are less than optimum in specific dosage forms to provide a sufficient drug dose in a dosage form that is small enough to be easily delivered to a use environment.

15

IMPROVED CHEMICAL STABILITY

When the compositions of the present invention comprise dispersions of acid-sensitive drug and neutral polymer(s), such compositions provide improved chemical stability of the drug relative to a control composition comprised of an equivalent quantity of drug dispersed in an acidic dispersion polymer. The acidic dispersion polymer used in the control composition may be any polymer suitable for determining whether the drug is acid-sensitive that meets the definition of an "acidic polymer." Convenient acidic dispersion polymers for use in the control composition include HPMCAS, CAP and HPMCP. A composition of the present invention will meet the chemical stability requirements discussed below relative to at least one control dispersion of drug and either HPMCAS, HPMCP, or CAP.

A relative degree of improvement in chemical stability of an acid-sensitive drug in a test composition may be determined by taking the ratio of the degree of degradation of the drug in a control composition and the degree of degradation of the drug in a test composition under the same storage conditions for the same storage time period. The test composition is simply the composition of acid-sensitive drug, neutral polymer and optional additional excipients of the present invention. The control composition is the same as the test composition with the exception that an acidic dispersion polymer (that is, either HPMCAS, HPMCP or CAP) replaces the neutral dispersion polymer. For

increases. It has also been found that the aqueous concentration of the drug in a use environment tends to improve as the fraction of drug present in the amorphous state in the dispersion increases. Accordingly, a "major portion" of the drug in the dispersion is amorphous and preferably the drug in the dispersion is substantially amorphous. As used herein, a "major portion" and "substantially amorphous" mean that the amount of the drug in crystalline form does not exceed about 40 wt% and about 25 wt%, respectively. More preferably, the drug in the dispersion is "almost completely amorphous," meaning that the amount of drug in the crystalline form does not exceed 10 wt%. Amounts of crystalline drug may be measured by powder X-ray diffraction, Scanning Electron Microscope (SEM) analysis, differential scanning calorimetry (DSC), or any other standard quantitative measurement.

The amorphous drug can exist as a pure phase, as a solid solution of drug homogeneously distributed throughout the polymer or any combination of these states or those states that lie intermediate between them. In cases where the drug is a low-solubility drug and concentration or bioavailability enhancement is desired, the dispersion is preferably "substantially homogeneous" so that the amorphous drug is dispersed as homogeneously as possible throughout the polymer. Dispersions of the present invention that are substantially homogeneous generally are more physically stable and have improved concentration-enhancing properties and, in turn improved bioavailability, relative to nonhomogeneous dispersions. As used herein, "substantially homogeneous" means that the drug present in relatively pure amorphous domains within the solid dispersion is relatively small, on the order of less than 20%, and preferably less than 10% of the total amount of drug.

While the dispersion may have some drug-rich domains, it is preferred that the dispersion itself have a single glass transition temperature (T_g) which demonstrates that the dispersion is substantially homogeneous. This contrasts with a simple physical mixture of pure amorphous drug particles and pure amorphous polymer particles which generally display two distinct T_g s, one that of the drug and one that of the polymer. T_g as used herein is the characteristic temperature where a glassy material, upon gradual heating, undergoes a relatively rapid (e.g., 10 to 100 seconds) physical change from a glass state to a rubber state. Almost completely amorphous, substantially homogeneous dispersions may be made by any of the processes discussed above.

drugs. When comparing dispersions under storage conditions which approximate ambient conditions, e.g., 30°C and 60% RH, the storage period may need to be several months up to two years.

In addition, it is preferred that the dispersions comprising acid-sensitive drug and neutral polymer(s) result in drug stability such that the acid-sensitive drug has a degree of degradation of less than about 2 wt%, more preferably less than about 0.5 wt%, and most preferably less than about 0.1 wt% when stored at 40°C and 75% RH for six months, or less than 2 wt%, more preferably 0.5 wt%, and more preferably 0.1 wt%, when stored at 30°C and 60% RH for one year. Nevertheless, the compositions of the present invention may have a degree of degradation that is much greater than the preferred values, so long as the dispersion achieves the degree of improvement relative to a control composition as described above.

PREPARATION OF DISPERSIONS

Dispersions of the drugs and neutral dispersion polymer may be made according to any known process which results in at least a "major portion" (meaning at least 60 wt%) of the drug being in the amorphous state. Such processes include mechanical, thermal and solvent processes. Exemplary mechanical processes include milling and extrusion; melt processes include high temperature fusion, solvent modified fusion and melt-congeal processes; and solvent processes include non-solvent precipitation, spray coating and spray-drying. See, for example, U.S. Patent No. 5,456,923 and U.S. Patent No. 5,939,099 which describe formation of dispersions via extrusion processes; U.S. Patent No. 5,340,591 and U.S. Patent No. 4,673,564 which describe forming dispersions by milling processes; and U.S. Patent No. 5,707,646 and U.S. Patent No. 4,894,235 which describe the formation of dispersions via melt/congeal processes, the disclosures of which are incorporated by reference.

While the drug in its pure state may be crystalline or amorphous, at least a major portion of the drug in the dispersion is amorphous. By "amorphous" is meant simply that the drug is in a non-crystalline state. As used herein, the term "a major portion" of the drug means that at least 60% of the drug in the dispersion is in the amorphous form, rather than the crystalline form. In general, a drug is more reactive in its amorphous state relative to its crystalline state and so the need to use a neutral dispersion polymer to prevent degradation of acid-sensitive drug increases as the fraction of drug in the amorphous state

may vary from about 50°C up to about 200°C depending on the melting point of the drug and polymer, which is a function of the polymer grade chosen and the amount of water, if any, added. Generally, the higher the melting point of the drug and polymer, the higher the processing temperature. Generally, the lowest
5 processing temperature that produces a satisfactory dispersion (almost completely amorphous and substantially homogeneous) is chosen.

A method for forming dispersions that is often preferred is "solvent processing," which consists of dissolution of the drug and one or more neutral polymers in a common solvent. The term "solvent" is used broadly and includes
10 mixtures of solvents. "Common" here means that the solvent, which can be a mixture of compounds, will simultaneously dissolve both the drug and the polymer(s).

After both the drug and polymer(s) have been dissolved, the solvent is removed by evaporation or by mixing with a non-solvent. Exemplary processes
15 are spray-drying, spray-coating (pan-coating, fluidized bed coating, etc.), vacuum evaporation, precipitation by mixing of the polymer and drug solution with a non-solvent such as CO₂, hexane, or heptane, and processes that combine solvents with heat and mechanical energy such as extrusion processes that include the use of a solvent as a processing aid or even wet
20 granulation processes as long as such processes result in at least a major portion of the drug being amorphous. Preferably, removal of the solvent is rapid and results in a solid dispersion which is substantially homogeneous. In substantially homogeneous dispersions, the drug is dispersed as homogeneously as possible throughout the polymer and can be thought of as a
25 solid solution of drug dispersed in the polymer(s). When the resulting dispersion constitutes a solid solution of drug in polymer, the dispersion may be thermodynamically stable, meaning that the concentration of drug in the polymer is at or below its equilibrium value, or it may be considered a supersaturated solid solution where the drug concentration in the dispersion
30 polymer(s) is above its equilibrium value.

The solvent may be removed through the process of spray-drying. The term spray-drying is used conventionally and broadly refers to processes involving breaking up liquid mixtures into small droplets (atomization) and rapidly removing solvent from the mixture in a container (spray-drying
35 apparatus) where there is a strong driving force for evaporation of solvent from

To obtain the maximum level of concentration and bioavailability enhancement, particularly upon storage for long times prior to use, it is preferred that the drug remain, to the extent possible, in the amorphous state. The inventors have found that this is best achieved when the glass-transition temperature, T_g , of the amorphous dispersion is substantially above the storage temperature of the dispersion. In particular, it is preferable that the T_g of the amorphous dispersion is at least 40°C and preferably at least 60°C. Since the T_g is a function of the water content of the dispersion which in turn is a function of the RH to which the dispersion is exposed, these T_g values refer to the T_g of the dispersion containing water in an amount that is in equilibrium with the RH equivalent to that found during storage. For those aspects of the invention in which the dispersion is a solid, substantially amorphous dispersion of drug in the dispersion polymer and in which the drug itself has a relatively low T_g (about 70°C or less) it is preferred that the dispersion polymer have a T_g of at least 40°C, preferably at least 70°C and more preferably greater than 100°C. Exemplary high T_g polymers include hydroxypropyl cellulose and hydroxypropyl methyl cellulose. Since conversion of amorphous drug to the crystalline state is related to the relative values of (1) the T_g of the dispersion (at the storage RH) and (2) the storage temperature, solid amorphous dispersions of the present invention may tend to remain in the amorphous state for longer periods when stored at relatively low temperatures and low relative humidities. In addition, packaging of such dispersions so as to prevent absorption of water or inclusion of a water absorbing material such as a dessicant to also prevent or retard water absorption can lead to a higher T_g during storage, thereby helping to retain the amorphous state. Likewise, storage at lower temperatures can also improve the retention of the amorphous state.

Turning now to particular methods, when either the neutral polymer or the drug has a relatively low melting point, typically less than about 200°C and preferably less than about 160°C, extrusion or melt-congeal processes that provide heat and/or mechanical energy are often suitable for forming almost completely amorphous dispersions. Often, when the drug has significant solubility in the dispersion material, such methods may also make substantially homogeneous dispersions. For example, 10 wt% drug and 90 wt% polyvinyl pyrrolidone (PVP) may be dry blended, with or without the addition of water, and the blend fed to a twin-screw extrusion device. The processing temperature

force for evaporation of solvent leads to actual drying times of a few seconds or less, and more typically less than 0.1 second. This rapid drying is often critical to the particles maintaining a uniform, homogeneous dispersion instead of separating into drug-rich and polymer-rich phases. As above, to get large
5 enhancements in concentration and bioavailability it is often necessary to obtain as homogeneous of a dispersion as possible. Solidification times should be less than 100 seconds, preferably less than a few seconds, and more preferably less than 1 second. In general, to achieve this rapid solidification of the
10 drug/polymer solution, it is preferred that the size of droplets formed during the spray-drying process are less than about 500 μm in diameter, and preferably less than about 300 μm . The resultant solid particles thus formed are generally less than about 300 μm in diameter.

Following solidification, the solid powder typically stays in the spray-drying chamber for about 5 to 60 seconds, further evaporating solvent from the
15 solid powder. The final solvent content of the solid dispersion as it exits the dryer should be low, since this reduces the mobility of drug molecules in the dispersion, thereby improving its stability. Generally, the solvent content of the dispersion as it leaves the spray-drying chamber should be less than about 10 wt% and preferably less than about 3 wt%. In some cases, it may be
20 preferable to spray a solvent or a solution of a polymer or other excipient into the spray-drying chamber to form granules, so long as the dispersion is not adversely affected.

Spray-drying processes and spray-drying equipment are described generally in Perry's *Chemical Engineers' Handbook*, Sixth Edition (R. H. Perry,
25 D. W. Green, J. O. Maloney, eds.) McGraw-Hill Book Co. 1984, pages 20-54 to 20-57. More details on spray-drying processes and equipment are reviewed by Marshall "Atomization and Spray-Drying," 50 *Chem. Eng. Prog. Monogr. Series* 2 (1954).

The solid amorphous dispersions formed from a low-solubility drug and
30 an amphiphilic hydroxyl-functional vinyl copolymer are preferably almost completely amorphous and substantially homogenous. Such almost completely amorphous, substantially homogenous dispersions are generally more physically stable (that is, resist phase separation or crystallization upon storage), and show higher levels of concentration enhancement. As with the
35 dispersions of other embodiments of the invention, such dispersions may in

the droplets. The strong driving force for solvent evaporation is generally provided by maintaining the partial pressure of solvent in the spray-drying apparatus well below the vapor pressure of the solvent at the temperature of the drying droplets. This is accomplished by either (1) maintaining the pressure in the spray-drying apparatus at a partial vacuum (e.g., 0.01 to 0.50 atm);
5 (2) mixing the liquid droplets with a warm drying gas; or (3) both. In addition, at least a portion of the heat required for evaporation of solvent may be provided by heating the spray solution.

Solvents suitable for spray-drying can be any organic compound in which the drug and polymer are mutually soluble. Preferably, the solvent is also volatile with a boiling point of 150°C or less. In addition, the solvent should have relatively low toxicity and be removed from the dispersion to a level that is acceptable according to The International Committee on Harmonization (ICH) guidelines. Removal of solvent to this level may require a processing step such as tray-drying or fluidized-bed drying subsequent to the spray-drying or spray-coating process. Preferred solvents include alcohols such as methanol,
15 ethanol, n-propanol, iso-propanol, and butanol; ketones such as acetone, methyl ethyl ketone and methyl iso-butyl ketone; esters such as ethyl acetate and propylacetate; and various other solvents such as acetonitrile, methylene chloride, toluene, and 1,1,1-trichloroethane. Lower volatility solvents such as dimethyl acetamide or dimethylsulfoxide can also be used. Mixtures of solvents, such as 50% methanol and 50% acetone, can also be used, as can mixtures with water as long as the polymer and drug are sufficiently soluble to make the spray-drying process practicable. In particular, when the neutral
20 polymer is relatively hydrophilic such as hydroxypropyl methyl cellulose (HPMC), or certain amphiphilic hydroxyl-functional vinyl copolymers, it is often preferred to use a mixture of an organic solvent and water to ensure that both drug and polymer are simultaneously soluble.

Generally, the temperature and flow rate of the drying gas is chosen so that the polymer/drug-solution droplets are dry enough by the time they reach the wall of the apparatus that they are essentially solid, and so that they form a fine powder and do not stick to the apparatus wall. The actual length of time to achieve this level of dryness depends on the size of the droplets. Droplet sizes generally range from 1 μ m to 500 μ m in diameter, with 5 to 200 μ m being more
35 typical. The large surface-to-volume ratio of the droplets and the large driving

comprise a low volatility solvent. When the dispersion polymer is a vinylacetate/vinylalcohol copolymer of the preferred type described earlier, preferred solvents include mixtures of methanol and water.

Returning to dispersions generally, the amount of polymer relative to the amount of drug present in the dispersions of the present invention depends on the drug and polymer and may vary widely from a drug-to-polymer weight ratio of from 0.01 to about 4 (e.g., 1 wt% drug to 80 wt% drug). However, in most cases it is preferred that the drug-to-polymer ratio is greater than about 0.05 (4.8 wt% drug) and less than about 2.5 (71 wt% drug).

In addition to the drug and polymer(s), the dispersions of the present invention may include optional additional ingredients. One optional additional ingredient is a buffer. Buffers suitable for use in the dispersions of the present invention are preferably basic. Buffers are particularly preferred for dispersions of neutral polymers and drugs that have a high degree of acid-sensitivity. The buffers may reduce the risk of drug degradation due to the presence of acidic species in the dispersion, elsewhere in the composition, or acidic or basic species formed during storage. The compositions preferably provide improved chemical stability relative to dispersions that are free from the buffer.

Exemplary buffers that may be used in the dispersions of the present invention include sodium acetate, ammonium acetate, sodium carbonate, sodium bicarbonate, disodium hydrogen phosphate and trisodium phosphate. It is generally preferred that the buffer be chosen such that it maintains the pH at the optimum value during storage to minimize degradation reactions. For most acid-sensitive drugs, the preferred pH value is greater than about 6. Since many dispersion polymers such as cellulose are also base-sensitive, it is generally preferred that the buffer be chosen to maintain the dispersion pH between about 6 and 10. In cases where the acid-sensitive drug is also a base-sensitive drug, it is often preferred to maintain the dispersion pH between about 6 and about 8. To accomplish this, it is sometimes desirable for the buffer to be a blend of protonated and deprotonated material, where the pK_a of the protonated material is near the desired dispersion pH. Such buffers may comprise from 0.1 to 20 wt% of the dispersion.

Alternatively, the dispersion may include basic excipients such as mono-, di- or tri-ethanolamine, or glucoseamine, which may comprise from 0.1

principle be formed by any known process, including mechanical, thermal and solvent processes. However, "solvent processing" is preferred for forming dispersions of amphiphilic hydroxyl-functional vinyl copolymers because when properly performed, solvent processing tends to produce dispersions in which the drug is almost completely amorphous and in which the dispersion is substantially homogeneous.

In order to achieve such preferred dispersions, a copolymer type and solvent must be chosen to yield a solution in which the amphiphilic, hydroxyl-functional vinyl copolymer and low-solubility drug are both soluble and preferably highly soluble. Preferably, both the copolymer and low-solubility drug each have solubilities in the solvent at 37°C of at least 0.5 wt% and preferably at least 2.0 wt% and more preferably at least 5.0 wt%.

In order to have the drug and copolymer both have such high solubilities in the solvent while the polymer is also sufficiently soluble in aqueous solution, the polymer molecular weight must generally not be too high and the polymer must have the correct n/m values. For example it has been found that for vinylacetate/vinylalcohol copolymers that n/m values are preferably between about 2 and 100 and molecular weights are preferably less than about 50,000 daltons.

In addition, the solvent is preferably chosen such that the lesser of either (1) the polymer solubility or (2) the product of the drug solubility and the ratio of the polymer to drug in the final dispersion, is high and is preferably maximized. To achieve this, it is often desirable to use blends of two or more solvents. In particular, a blend of water and an organic solvent is often preferred. Exemplary preferred solvents are mixtures of water and alcohols such as methanol, ethanol, n-propanol, iso-propanol and the isomers of butanol.

In addition it is preferable for the solvent processing to be conducted such that solvent is removed rapidly so that the polymer-drug solution solidifies rapidly. Thus solvent processes such as spray-coating and spray-drying are preferred.

In addition, to achieve rapid solvent removal, and to keep the residual solvent level in the dispersion low (preferably less than about 5 wt%), a relatively volatile solvent is chosen. Preferably the boiling points of the solvents are less than about 150°C and more preferably less than about 125°C. When the solvent is a mixture of solvents, up to about 40% of the solvent may

dry or wet granulation, which does not convert the dispersion and the second polymer to another molecular dispersion, may be used.

Alternatively, the dispersion and second polymer may be co-administered, meaning that the dispersion may be administered separately
5 from, but within the same general time frame as, the second polymer. Thus, a dispersion may, for example, be administered in its own dosage form which is taken at approximately the same time as the second polymer which is in a separate dosage form. If administered separately, it is generally preferred to administer both the dispersion and the second polymer within 60 minutes of
10 each other, so that the two are present together in the use environment. When not administered approximately simultaneously (e.g., within a minute or two of each other), the second polymer is preferably administered prior to the dispersion.

The second polymer may be any concentration-enhancing polymer,
15 such as those described above in connection with the neutral dispersion polymers. As described above, the inventors have found that ionizable, cellulosic polymers, particularly those that are acidic cellulosic enteric polymers, provide superior enhancement in aqueous concentration of the drug in a use environment relative to other polymers, and are therefore preferred in the
20 absence of their reactivity with the drug. Many of these ionizable, cellulosic polymers have acidic functional groups and therefore are inappropriate for use as a dispersion polymer with acid-sensitive drugs. However, the concentration-enhancing advantage provided by such ionizable concentration-enhancing polymers may be achieved by simply combining such a polymer as the second
25 polymer with a pre-formed dispersion of the acid-sensitive drug and a neutral polymer in a fashion that does not alter the neutral characteristic of the pre-formed dispersion.

A preferred class of concentration-enhancing polymers suitable for use as the second polymer comprises ionizable cellulosic polymers with at least one
30 ester- and/or ether- linked substituent in which the polymer has a degree of substitution of at least 0.05 for each substituent. Particularly preferred hydrophilic substituents for use in making ionizable polymers are those that are ether- or ester-linked acidic ionizable groups such as carboxylic acids, thiocarboxylic acids and sulfonates, and groups that may be non-acidic such as
35 substituted phenoxy groups, amines, and phosphates. Exemplary ether-linked

to 20 wt% of the dispersion. Such basic excipients may be used to control the dispersion pH as described above.

MIXTURES OF DISPERSIONS AND CONCENTRATION-ENHANCING POLYMER

5 A separate embodiment of the invention comprises a combination of
(1) a dispersion of an acid-sensitive, low-solubility drug and neutral dispersion
polymer(s) and (2) a second polymer that is concentration-enhancing. The
dispersion is free from at least a portion of the second polymer, and is
preferably substantially free from the second polymer. By "substantially free" is
10 meant that the dispersion and the second polymer are not mixed at the
molecular level to form a dispersion containing the second polymer. While a
small amount of the second polymer may be incorporated into the dispersion as
a result of processing, nevertheless the drug in the dispersion meets the
chemical stability requirements discussed above for dispersions of acid-
15 sensitive drugs and neutral polymers notwithstanding the presence of the
second polymer in the composition. The neutral dispersion polymer optionally
may also be concentration-enhancing.

The compositions of this aspect of the present invention are generally
physical combinations comprising the dispersion and the second polymer.
20 "Combination" as used herein means that the dispersion and second polymer
may be in physical contact with each other or in close proximity but without the
necessity of being physically mixed at the molecular level (i.e., a dispersion).
The dispersion and concentration-enhancing polymer may be in different
regions of the composition. For example, the solid composition may be in the
25 form of a multi-layer tablet, as known in the art, wherein one or more layers
comprises the dispersion and one or more different layers comprises the
second polymer. Yet another example may constitute a coated tablet wherein
either the dispersion or the second polymer or both may be present in the tablet
core and the coating may comprise the second polymer. Alternatively, the
30 combination can be in the form of a simple dry physical mixture wherein both
the dispersion and the second polymer are mixed in particulate form and
wherein the particles of each, regardless of size, retain the same individual
physical properties that they exhibit in bulk. Any conventional method used to
mix the dispersion and second polymer together such as physical mixing and

ethyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate, hydroxypropyl methyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate succinate, cellulose propionate trimellitate, cellulose butyrate trimellitate, cellulose acetate terephthalate, cellulose acetate isophthalate, 5 cellulose acetate pyridinedicarboxylate, salicylic acid cellulose acetate, hydroxypropyl salicylic acid cellulose acetate, ethylbenzoic acid cellulose acetate, hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl phthalic acid cellulose acetate, ethyl nicotinic acid cellulose acetate, and ethyl picolinic acid cellulose acetate.

10 Another class of concentration-enhancing polymers suitable for use with the present invention comprises ionizable non-cellulosic polymers. Exemplary polymers of this type that are generally acidic include carboxylic acid-functionalized vinyl polymers, such as the carboxylic acid functionalized polymethacrylates and carboxylic acid functionalized polyacrylates such as the 15 EUDRAGITS® manufactured by Rohm Tech Inc., of Malden, Massachusetts, and acidic proteins.

A preferred class of concentration-enhancing polymers comprises polymers that are "amphiphilic" in nature, meaning that the polymer has hydrophobic and hydrophilic portions. The hydrophobic portion may comprise 20 groups such as aliphatic or aromatic hydrocarbon groups. The hydrophilic portion may comprise either ionizable or non-ionizable groups that are capable of hydrogen bonding such as hydroxyls, carboxylic acids, esters, amines or amides.

Amphiphilic polymers are preferred because it is believed that such 25 polymers may tend to have relatively strong interactions with the drug and may promote the formation of various types of polymer/drug assemblies in the use environment. Ionizable polymers, and particularly those that have a significant level of carboxylic acid functional groups (e.g., acidic polymers) typically show the greatest concentration and bioavailability enhancements and therefore are 30 preferred for use with low-solubility drugs. However, they may not form a part of the dispersion when the drug is acid-sensitive. Thus, a particularly preferred composition is one in which an acid-sensitive drug is dispersed in a neutral or non-ionizable polymer and an ionizable or acidic polymer is added such that it is not part of the dispersion.

ionizable substituents include: carboxylic acids, such as acetic acid, propionic acid, benzoic acid, salicylic acid, alkoxybenzoic acids such as ethoxybenzoic acid or propoxybenzoic acid, the various isomers of alkoxyphthalic acid such as ethoxyphthalic acid and ethoxyisophthalic acid, the various isomers of
5 alkoxy nicotinic acid such as ethoxy nicotinic acid, and the various isomers of picolinic acid such as ethoxypicolinic acid, etc.; thiocarboxylic acids, such as thioacetic acid; substituted phenoxy groups, such as hydroxyphenoxy, etc.; amines, such as aminoethoxy, diethylaminoethoxy, trimethylaminoethoxy, etc.; phosphates, such as phosphate ethoxy; and sulfonates, such as sulphonate
10 ethoxy. Exemplary ester linked ionizable substituents include: carboxylic acids, such as succinate, citrate, phthalate, terephthalate, isophthalate, trimellitate, and the various isomers of pyridinedicarboxylic acid, etc.; thiocarboxylic acids, such as thiosuccinate; substituted phenoxy groups, such as amino salicylic acid; amines, such as natural or synthetic amino acids, such as alanine or
15 phenylalanine; phosphates, such as acetyl phosphate; and sulfonates, such as acetyl sulfonate. For aromatic-substituted polymers to also have the requisite aqueous solubility, it is also desirable that sufficient hydrophilic groups such as hydroxypropyl or carboxylic acid functional groups be attached to the polymer to render the polymer aqueous soluble at least at pH values where any ionizable
20 groups are ionized. In some cases, the aromatic group may itself be ionizable, such as phthalate or trimellitate substituents.

Exemplary ionizable cellulosic concentration-enhancing polymers that are at least partially ionized at physiologically relevant pHs include:
hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl
25 cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, hydroxyethyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, hydroxyethyl methyl cellulose acetate succinate, hydroxyethyl methyl cellulose acetate phthalate, carboxyethyl cellulose, carboxymethyl cellulose, cellulose acetate phthalate, methyl cellulose
30 acetate phthalate, ethyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, hydroxypropyl methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, hydroxypropyl methyl cellulose acetate succinate phthalate, hydroxypropyl methyl cellulose succinate phthalate, cellulose propionate phthalate, hydroxypropyl cellulose butyrate
35 phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate,

acetate phthalate, hydroxypropyl methyl cellulose phthalate, hydroxypropyl methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, cellulose propionate phthalate, hydroxypropyl cellulose butyrate phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate, 5 ethyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate, hydroxypropyl methyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate succinate, cellulose propionate trimellitate, cellulose butyrate trimellitate, cellulose acetate terephthalate, cellulose acetate isophthalate, cellulose acetate pyridinedicarboxylate, salicylic acid cellulose acetate, 10 hydroxypropyl salicylic acid cellulose acetate, ethylbenzoic acid cellulose acetate, hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl phthalic acid cellulose acetate, ethyl nicotinic acid cellulose acetate, and ethyl picolinic acid cellulose acetate.

Another particularly desirable subset of cellulosic ionizable polymers are 15 those that possess a non-aromatic carboxylate substituent. Exemplary polymers include hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, and hydroxyethyl cellulose acetate succinate.

20 Non-cellulosic polymers that are amphiphilic are copolymers of a relatively hydrophilic and a relatively hydrophobic monomer. Examples include acrylate and methacrylate copolymers. Exemplary commercial grades of such copolymers include the EUDRAGITS, which are copolymers of methacrylates and acrylates. Another suitable class of amphiphilic copolymers are the 25 amphiphilic hydroxyl-functional vinyl copolymers described previously.

EXCIPIENTS AND DOSAGE FORMS

Although the key ingredients present in the compositions of the present invention are simply the dispersion and optional non-dispersion concentration-enhancing polymer(s), the inclusion of other excipients in the composition may 30 be useful. These excipients may be utilized with the drug and polymer composition in order to formulate the composition into tablets, capsules, suppositories, suspensions, powders for suspension, creams, transdermal patches, depots, and the like. The composition of drug and polymer can be added to other dosage form ingredients in essentially any manner that does not 35 substantially alter the drug. The excipients may be either physically mixed with

In addition, ionic, and in particular acidic polymers, are thought to show desirable concentration-enhancement properties due to the repulsion of the like charges of the ionized groups of such polymers. This repulsion may serve to limit the size of the polymer/drug assemblies to the nanometer or submicron scale. For example, while not wishing to be bound by a particular theory, such polymer/drug assemblies may comprise hydrophobic drug clusters surrounded by the polymer with the polymer's hydrophobic regions turned inward towards the drug and the hydrophilic regions of the polymer turned outward toward the aqueous environment. Alternatively, depending on the specific chemical nature of the drug, the ionized functional groups of the polymer may associate, for example, via ion pairing or hydrogen bonds, with ionic or polar groups of the drug. In the case of ionizable polymers, the hydrophilic regions of the polymer would include the ionized functional groups. Such polymer/drug assemblies in solution may well resemble charged polymeric micellar-like structures. In any case, regardless of the mechanism of action, the inventors have observed that such amphiphilic polymers, particularly ionizable cellulosic polymers, have been shown to improve the maximum drug concentration (MDC) and/or area under the curve (AUC) of the drug in aqueous solution relative to control compositions free from such polymers.

Surprisingly, such amphiphilic polymers can greatly enhance the maximum concentration of drug obtained when the drug is dosed to a use environment. In addition, such amphiphilic polymers may interact with the drug to prevent the precipitation or crystallization of the drug from solution despite its concentration being substantially above its equilibrium concentration.

Exemplary ionizable cellulosic polymers that meet the definition of amphiphilic, having hydrophilic and hydrophobic regions, include polymers such as cellulose acetate phthalate and cellulose acetate trimellitate where the cellulosic repeat units that have one or more acetate substituents are hydrophobic relative to those that have no acetate substituents or have one or more ionized phthalate or trimellitate substituents.

A particularly desirable subset of cellulosic ionizable concentration-enhancing polymers are those that possess both a carboxylic acid functional aromatic substituent and an alkylate substituent and thus are amphiphilic. Exemplary polymers include cellulose acetate phthalate, methyl cellulose acetate phthalate, ethyl cellulose acetate phthalate, hydroxypropyl cellulose

Examples of matrix materials, fillers, or diluents include lactose, mannitol, xylitol, microcrystalline cellulose, calcium diphosphate, dicalcium phosphate and starch.

5 Examples of disintegrants include sodium starch glycolate, sodium alginate, carboxy methyl cellulose sodium, methyl cellulose, and croscarmellose sodium, and crosslinked forms of polyvinyl pyrrolidone such as those sold under the trade name CROSPOVIDONE (available from BASF Corporation).

Examples of binders include methyl cellulose, microcrystalline cellulose, starch, and gums such as guar gum, and tragacanth.

10 Examples of lubricants include magnesium stearate, calcium stearate, and stearic acid.

Examples of preservatives include sulfites (an antioxidant), benzalkonium chloride, methyl paraben, propyl paraben, benzyl alcohol and sodium benzoate.

15 Examples of suspending agents or thickeners include xanthan gum, starch, guar gum, sodium alginate, carboxymethyl cellulose, sodium carboxymethyl cellulose, methyl cellulose, hydroxypropyl methyl cellulose, polyacrylic acid, silica gel, aluminum silicate, magnesium silicate, and titanium dioxide.

20 Examples of anticaking agents or fillers include silicon oxide and lactose.

Examples of solubilizers include ethanol, propylene glycol or polyethylene glycol.

Other conventional excipients may be employed in the compositions of this invention, including those excipients well-known in the art. Generally,
25 excipients such as pigments, lubricants, flavorants, and so forth may be used for customary purposes and in typical amounts without adversely affecting the properties of the compositions. These excipients may be utilized in order to formulate the composition into tablets, capsules, suspensions, powders for suspension, creams, transdermal patches, and the like.

30 The compositions of the present invention may be delivered by a wide variety of routes, including, but not limited to, oral, nasal, rectal, vaginal, subcutaneous, intravenous, and pulmonary. Generally, the oral route is preferred.

35 Compositions of this invention may also be used in a wide variety of dosage forms for administration of drugs. Exemplary dosage forms are

the dispersion and/or included within the dispersion. However, where the drug is acid-sensitive, acidic excipients should not be added to the dispersion unless neutralized.

One very useful class of excipients is surfactants. Suitable surfactants include fatty acid and alkyl sulfonates; commercial surfactants such as benzalkonium chloride (HYAMINE® 1622, available from Lonza, Inc., Fairlawn, New Jersey); dioctyl sodium sulfosuccinate, DOCUSATE SODIUM™ (available from Mallinckrodt Spec. Chem., St. Louis, Missouri); polyoxyethylene sorbitan fatty acid esters (TWEEN®, available from ICI Americas Inc., Wilmington, Delaware; LIPOSORB® P-20 available from Lipochem Inc., Patterson New Jersey; CAPMUL® POE-0 available from Abitec Corp., Janesville, Wisconsin), and natural surfactants such as sodium taurocholic acid, 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine, lecithin, and other phospholipids and mono- and diglycerides. Preferably, the surfactants are not acidic. In cases where the surfactant is acidic, such as fatty acids, the salts of the acids are preferred. Thus, sodium dodecanate is preferred relative to dodecanoic acid. Such materials can advantageously be employed to increase the rate of dissolution by facilitating wetting, thereby increasing the maximum dissolved concentration, and also to inhibit crystallization or precipitation of drug by interacting with the dissolved drug by mechanisms such as complexation, formation of inclusion complexes, formation of micelles or adsorbing to the surface of solid drug, crystalline or amorphous. These surfactants may comprise up to 5 wt% of the composition.

The addition of pH modifiers such as acids, bases, or buffers may also be beneficial, retarding the dissolution of the composition (e.g., acids such as citric acid or succinic acid when the concentration-enhancing polymer is anionic) or, alternatively, enhancing the rate of dissolution of the composition (e.g., bases such as sodium acetate or amines when the polymer is anionic). Of course, where the drug is acid-sensitive, pH modifiers should not be added to the dispersion except as discussed above in connection with including buffers or basic excipients within the dispersion.

Conventional matrix materials, complexing agents, solubilizers, fillers, disintegrating agents (disintegrants), or binders may also be added as part of the composition itself or added by granulation via wet or mechanical or other means. These materials may comprise up to 90 wt% of the composition.

enteric polymer to prevent or retard dissolution until the dosage form leaves the stomach. Exemplary enteric coating materials include hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, carboxylic acid-functionalized polymethacrylates, and carboxylic acid-functionalized polyacrylate.

Compositions of this invention may be administered in a controlled release dosage form. In one such dosage form, the composition of the drug and polymer is incorporated into an erodible polymeric matrix device. By an erodible matrix is meant aqueous-erodible or water-swellaable or aqueous-soluble in the sense of being either erodible or swellaable or dissolvable in pure water or requiring the presence of an acid or base to ionize the polymeric matrix sufficiently to cause erosion or dissolution. When contacted with the aqueous environment of use, the erodible polymeric matrix imbibes water and forms an aqueous-swollen gel or "matrix" that entraps the dispersion of drug and polymer. The aqueous-swollen matrix gradually erodes, swells, disintegrates or dissolves in the environment of use, thereby controlling the release of the dispersion to the environment of use. Examples of such dosage forms are disclosed more fully in commonly assigned pending U.S. Patent Application Serial No. 09/495,059 filed January 31, 2000 which claimed the benefit of priority of provisional patent application Serial No. 60/119,400 filed February 10, 1999, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the compositions of the present invention may be administered by or incorporated into a non-erodible matrix device.

Alternatively, the compositions of the invention may be delivered using a coated osmotic controlled release dosage form. This dosage form has two components: (a) the core which contains an osmotic agent and the dispersion of drug and concentration-enhancing polymer; and (b) a non-dissolving and non-eroding coating surrounding the core, the coating controlling the influx of water to the core from an aqueous environment of use so as to cause drug release by extrusion of some or all of the core to the environment of use. The osmotic agent contained in the core of this device may be an aqueous-swellaable hydrophilic polymer, osmogen, or osmagent. The coating is preferably polymeric, aqueous-permeable, and has at least one delivery port. Examples of such dosage forms are disclosed more fully in commonly assigned pending U.S. Patent Application Serial No. 09/495,061 filed January 31, 2000 which claimed

powders or granules that may be taken orally either dry or reconstituted by addition of water or other liquids to form a paste, slurry, suspension or solution; tablets; capsules; multiparticulates; and pills. Various additives may be mixed, ground, or granulated with the compositions of this invention to form a material
5 suitable for the above dosage forms.

The compositions of the present invention may be formulated in various forms such that they are delivered as a suspension of particles in a liquid vehicle. Such suspensions may be formulated as a liquid or paste at the time of manufacture, or they may be formulated as a dry powder with a liquid, typically
10 water, added at a later time but prior to oral administration. Such powders that are constituted into a suspension are often termed sachets or oral powder for constitution (OPC) formulations. Such dosage forms can be formulated and reconstituted via any known procedure. The simplest approach is to formulate the dosage form as a dry powder that is reconstituted by simply adding water
15 and agitating. Alternatively, the dosage form may be formulated as a liquid and a dry powder that are combined and agitated to form the oral suspension. In yet another embodiment, the dosage form can be formulated as two powders which are reconstituted by first adding water to one powder to form a solution to which the second powder is combined with agitation to form the suspension.

20 Generally, it is preferred that the dispersion of drug be formulated for long-term storage in the dry state as this promotes the chemical and physical stability of the drug.

A preferred additive to such formulations is additional concentration-enhancing polymer which may act as a thickener or suspending agent as well
25 as to enhance the concentration of drug in the environment of use and may also act to prevent or retard precipitation or crystallization of drug from solution. Such preferred additives are hydroxyethyl cellulose, hydroxypropyl cellulose, and hydroxypropyl methyl cellulose. In particular, the salts of carboxylic acid functional polymers such as cellulose acetate phthalate, hydroxypropyl methyl
30 cellulose acetate succinate, and carboxymethyl cellulose are useful in this regard. Such polymers may be added in their salt forms or the salt form may be formed *in situ* during reconstitution by adding a base such as trisodium phosphate and the acid form of such polymers.

In some cases, the overall dosage form or particles, granules or beads
35 that make up the dosage form may have superior performance if coated with an

compaction, melt-congealing processes or by spray-coating seed cores. For example, in wet- and dry-granulation processes, the composition of drug and concentration-enhancing polymer is prepared as described above. This composition is then granulated to form multiparticulates of the desired size.

5 Other excipients, such as a binder (e.g., microcrystalline cellulose), may be blended with the composition to aid in processing and forming the multiparticulates. In the case of wet granulation, a binder such as microcrystalline cellulose may be included in the granulation fluid to aid in forming a suitable multiparticulate.

10 In any case, the resulting particles may themselves constitute the multiparticulate dosage form or they may be coated by various film-forming materials such as enteric polymers or water-swellaable or water-soluble polymers, or they may be combined with other excipients or vehicles to aid in dosing to patients.

15 Compositions of the present invention may be used to treat any condition which is subject to treatment by administering a drug.

Other features and embodiments of the invention will become apparent from the following examples which are given for illustration of the invention, rather than for limiting its intended scope.

EXAMPLES

Examples 1-2

20 This example discloses solid amorphous dispersions of an acid-sensitive, low-solubility drug and neutral polymer. For Example 1, a dispersion of quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3-fluorobenzyl]-2(S),7-dihydroxy-7-methyl-octyl] amide (Drug 1) and the neutral polymer hydroxypropyl methyl cellulose (HPMC E3 Prem) was made by preparing a solution containing 0.125 wt% Drug 1 and 0.375 wt% HPMC in methanol, and spraying the solution into a drying chamber using an atomizing spray nozzle as described below. For
25 Example 2, a dispersion of Drug 1 with the neutral polymer polyvinyl pyrrolidone (PVP- Plasdone K-29/32 available from ISP Technologies Inc., Wayne, NJ) was made by preparing a solution containing 0.33 wt% Drug 1 and 1.0 wt% PVP in acetone/methanol (9/1, wt/wt), and spray-drying the solution as described below.
30

the benefit of priority of provisional Patent Application Serial No. 60/119,406 filed February 10, 1999, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the compositions may be delivered via a coated hydrogel controlled release form having at least two components: (a) a core comprising the dispersion of the present invention and a hydrogel, and (b) a coating through which the dispersion has passage when the dosage form is exposed to a use environment. Examples of such dosage forms are more fully disclosed in commonly assigned European Patent EP0378404, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the drug mixture of the invention may be delivered via a coated hydrogel controlled release dosage form having at least three components: (a) a composition containing the dispersion, (b) a water-swellaable composition wherein the water-swellaable composition is in a separate region within a core formed by the drug-containing composition and the water-swellaable composition, and (c) a coating around the core that is water-permeable, water-insoluble, and has at least one delivery port therethrough. In use, the core imbibes water through the coating, swelling the water-swellaable composition and increasing the pressure within the core, and fluidizing the dispersion-containing composition. Because the coating remains intact, the dispersion-containing composition is extruded out of the delivery port into an environment of use. Examples of such dosage forms are more fully disclosed in commonly assigned pending U.S. Patent application Serial No. 09/745,095 filed December 20, 2000, which claimed the benefit of priority of provisional patent application 60/171,968 filed December 23, 1999, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the compositions may be administered as multiparticulates. Multiparticulates generally refer to dosage forms that comprise a multiplicity of particles that may range in size from about 10 μ m to about 2 mm, more typically about 100 μ m to 1 mm in diameter. Such multiparticulates may be packaged, for example, in a capsule such as a gelatin capsule or a capsule formed from an aqueous-soluble polymer such as HPMCAS, HPMC or starch or they may be dosed as a suspension or slurry in a liquid.

Such multiparticulates may be made by any known process, such as wet- and dry-granulation processes, extrusion/spheronization, roller-

Table 2

Ex.	Aqueous Soluble Polymer	Drug 1 Conc. in the Dispersion (wt%)	Pot. Before Storage (wt%)	Pot. Day 4 at 70°C (wt%)	Degree of Degradation Day 4 at 70°C (wt%)	Pot. Day 14 at 40°C/75% RH (wt%)	Degree of Degradation Day 14 at 40°C/75% RH (wt%)
1	HPMC	25	100	100	<1	100	<1
2	PVP	25	100	100	<1	99	1
C1	HPMCAS-LF	25	94	<1	>93	<1	>93

5

As the data show, the dispersions of Examples 1 and 2 formed with neutral polymers were much more chemically stable than the control dispersion C1 formed from HPMCAS. In fact, the dispersions of both Example 1 and Example 2 had a relative degree of improvement when compared with the control C1 of greater than 93 after only two weeks at 40°C/75% RH.

Example 4

In this example the dispersions of Examples 1 and 2 were tested to show that the dispersions provided concentration-enhancement of the drug in aqueous solution. For Control C2, the crystalline form of the drug alone was used without further processing. For this test, 7.2 mg of the dispersions of Examples 1 and 2, and 3.6 mg of Control C2, was added to respective microcentrifuge tubes. The tubes were placed in a 37°C temperature-controlled bath, and 1.8 mL phosphate buffered saline (PBS) at pH 6.5 and 290 mOsm/kg was added to each. The samples were quickly mixed using a vortex mixer for about 60 seconds. The samples were centrifuged at 13,000 G at 37°C for 1 minute. The resulting supernatant solutions were then sampled and diluted 1:6 (by volume) with methanol and then analyzed by high-performance liquid chromatography (HPLC). The contents of the tubes were mixed on the vortex mixer and allowed to stand undisturbed at 37°C until the next sample was

For Control C1, a dispersion of Drug 1 with hydroxy propyl methyl cellulose acetate succinate, LF-grade (HPMCAS-LF) (with about 14-18 wt% or 350-450 meq of succinate groups per mol of polymer) was made by preparing a solution containing 0.33 wt% Drug 1 and 1.0 wt% HPMCAS-LF in acetone, and spray-drying the solution as described below.

For Examples 1 and 2, and Control C1, the solutions were spray-dried by pumping the solution into a "mini" spray-dryer apparatus via a Cole Parmer 74900 series rate-controlling syringe pump at a rate of 1.3 mL/min. The drug/polymer solution was atomized through a Spraying Systems Co. two-fluid nozzle, Model No. SU1A using a heated stream of nitrogen (100°C). The spray solution was sprayed into an 11-cm diameter stainless steel chamber. The resulting solid amorphous dispersions containing 25 wt% Drug 1 were collected on filter paper, dried under vacuum, and stored in a dessicator. Table 1 summarizes the spray-drying variables.

Table 1

Ex.	Drug Mass (mg)	Drug No.	Polymer Mass (mg)	Polymer	Solvent Mass (g)	Solvent (wt/wt)
1	50	1	150	HPMC	40	Methanol
2	50	1	150	PVP	15	Acetone/methanol (9/1)
C1	50	1	150	HPMCAS	15	Acetone

Example 3

In this example the chemical stability of the dispersions of Examples 1 and 2 was assessed by monitoring the potency of the drug before and after exposure to increased temperatures and relative humidity (RH) in accelerated-aging studies. Dispersions of Examples 1 and 2, and Control C1, were placed in two controlled atmosphere chambers: one chamber maintained at 70°C (no RH control); the second chamber maintained at 40°C and 75% RH. Potencies of the dispersions before and after storage were determined using HPLC. A Kromasil C₄ HPLC column was used with a mobile phase of 45 vol% of 0.2 vol% H₃PO₄, and 55 vol% acetonitrile. UV detection was measured at 245 nm. Drug 1 potency was the percent of the total HPLC peak area corresponding to the theoretical amount of drug originally present in the dispersion prior to storage based on the amount of drug present in the initial solutions before spray-drying. The results are shown in Table 2 below.

and an AUC_{90} that was 3.0-fold, that of the control C2, while the PVP Drug 1 dispersion of Example 2 provided a C_{max90} that was 2.8-fold, and an AUC_{90} that was 3.1-fold, that of Control C2.

Example 5

The dispersion of Example 2 was stored for 1.5 years at room temperature (sealed), and then analyzed for Drug 1 potency by HPLC using the method described in Example 3. Results are shown in Table 5.

Table 5

Ex.	Polymer	Drug 1 Conc. in the Dispersion (wt%)	Potency Before Storage (wt%)	Potency After 1.5 yrs at room temp./sealed (wt%)
2	PVP	25	100	100

As the data show, the dispersion of Example 2 formed with PVP is chemically stable for extended duration under ambient conditions. This data is consistent with the results of accelerated storage stability tests for the dispersion of Example 2 (stored at 70°C and 40°C/75%RH) shown in Example 3 (Table 2).

In vitro dissolution tests were performed for the dispersion of Example 2 before and after 1.5 years storage, using the procedure described in Example 4, with the following exceptions: 7.2 mg of the dispersion of Example 2 was used in initial dissolution tests, and 5.4 mg of Example 2 was used in dissolution tests after 1.5 years storage. The results are reported as a percentage of the total dose dissolved, to facilitate comparison. The results are shown in Table 6.

taken. Samples were collected at 4, 10, 20, 40, 90, and 1200 minutes. The results are shown in Table 3.

Table 3

Example	Time (min)	[Drug 1] ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
1	0	0	0
	4	863	1,700
	10	895	7,000
	20	885	15,900
	40	925	34,000
	90	907	79,800
	1200	926	1,097,100
2	0	0	0
	4	923	1,800
	10	928	7,400
	20	913	16,600
	40	947	35,200
	90	927	82,100
	1200	929	1,112,100
C2	0	0	0
	4	274	600
	10	266	2,200
	20	338	5,200
	40	289	11,500
	90	300	26,200
	1200	303	360,800

The concentrations of Drug 1 obtained in these samples were used to determine the values of the maximum drug concentration between 0 and 90 minutes ($C_{\text{max}90}$) and the area under the curve from 0 to 90 minutes (AUC_{90}).

The results are shown in Table 4.

Table 4

Example	$C_{\text{max}90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
1	925	79,800
2	947	82,000
C2	338	26,200

As can be seen from the data, the neutral polymers HPMC and PVP are both concentration-enhancing polymers in the dispersions of Examples 1 and 2. The Drug 1 and HPMC dispersion of Example 1 provided a $C_{\text{max}90}$ that was 2.7-fold,

181

Table 7

Ex.	Drug Mass (mg)	Drug No.	Polymer Mass (mg)	Polymer	Solvent Mass (g)	Solvent (wt/wt)
6	50	1	150	HPMC	4	methanol/ water (9/1)
7	50	1	950	HPMC	30	methanol/ water (9/1)
8	125	1	375	Eudragit	10	acetone/ methanol (1/1)
9	250	1	375/375	HPMC/ Eudragit	20	methanol/ water (9/1)
C3	50	1	150	HPMCAS- HF	4	methanol

5

Example 10

In this example the chemical stability of the dispersions of Examples 6-9 was assessed. Dispersions were stored under elevated temperature and humidity conditions to increase the rate of physical changes occurring in the materials in order to simulate a longer storage interval in a typical storage environment. Drug purity was determined using HPLC as in Example 3. Results of purity analysis of the dispersions of Drug 1 after various storage intervals under various storage conditions are shown in Table 8.

10

Table 6

Time (min)	Initial Drug 1 Concentration (% dissolved)	Drug 1 Concentration After 1.5 yrs Storage (% dissolved)
0	0	0
4	91	95
10	92	90
20	90	87
40	94	88
90	92	98
1200	92	89

5 As can be seen from the data, the dissolution performance of the dispersion of Example 2 was maintained after 1.5 years storage, indicating that the dispersion is physically stable and Drug 1 is chemically stable for at least 1.5 years at ambient conditions.

Examples 6-9

10 These examples disclose additional dispersions of Drug 1 prepared with the dispersion polymers HPMC, Eudragit® E100 (available from Rohm America Inc., Piscataway, NJ), and an HPMC/Eudragit® E100 mixture. Eudragit E100 is a polymethacrylate polymer which contains basic groups. Examples 6 through 9 were prepared by spray-drying a solution of Drug 1 and polymer, as described in Example 1. For Example 6, the solution consisted of 1.25 wt% Drug 1 and 3.75 wt% HPMC in methanol/water (9/1, wt/wt). For Example 7, the solution consisted of 0.25 wt% Drug 1 and 4.75 wt% HPMC in methanol/water (9/1, wt/wt). For Example 8, the solution consisted of 1.25 wt% Drug 1 and 3.75 wt% Eudragit® E100 in methanol/acetone (1/1, wt/wt). For Example 9, the solution consisted of 1.25 wt% Drug 1 and 3.75 wt% HPMC/Eudragit (1/1 wt/wt) in methanol/water (9/1, wt/wt). The solution variables are summarized in Table 7.

20 For Control C3, the solution consisted of 1.25 wt% Drug 1 and 3.75 wt% HPMCAS-HF (HF grade) and methanol.

25

Table 9

Ex.	Drug Mass (mg)	Excip-ient	Excip-ient Mass (mg)	Polymer	Polymer Mass (mg)	Solvent (wt/wt)	Solvent Mass (g)
11	125	NaOAc	51.4	HPMC	375	MeOH/ water (9/1)	10
12	125	NaHCO ₃	32	HPMC	375	MeOH/ water (9/1)	10

5

Example 13

Chemical stability of the acid-sensitive Drug 1 in the dispersions of Examples 11 and 12 containing buffers was determined by measuring the drug purity before and after storage for Examples 11 and 12, and Control C3. Dispersions were stored under elevated temperature and humidity conditions to increase the rate of changes occurring in the materials in order to simulate a longer storage interval in a typical storage environment. Drug purity was determined using HPLC. Results of analysis of dispersions of Examples 11 and 12 after five days at 40°C/75% RH are shown in Table 10.

15

Table 10

Ex.	Aqueous-Soluble Polymer/Buffer	Drug 1 Conc. in the Dispersion (wt%)	Potency Day 5 at 40°C/75%RH (%)	Degree of Degradation (%)
11	HPMC/NaOAc	23	97.79	2.2
12	HPMC/NaHCO ₃	24	95.65	4.3
C3	HPMCAS-HF	25	60	40

The stability of the dispersions with neutral polymers containing buffers was significantly improved in comparison to the stability of the dispersion with HPMCAS-HF. The relative degree of improvement after five days compared with the Control C3 was 18.2 for Example 11, and 9.3 for Example 12.

Table 8

Ex.	Aqueous-Soluble Polymer	Drug 1 Conc. in the Dispersion (wt%)	Potency Day 5 at 40°C/75 RH (wt%)	Degree of Degradation Day 5 (wt%)
6	HPMC	25	97.3	2.7
7	HPMC	5	96.3	3.7
8	Eudragit	25	95.6	4.4
9	HPMC/Eudragit	25	97.9	2.1
C3	HPMCAS-HF	25	60	40

5 The chemical stability of the acid-sensitive drug in dispersions of Examples 6-9 containing neutral polymers were all significantly improved in comparison to the stability of dispersions of the control C3 having the acidic polymer HPMCAS-HF. The relative degree of improvement compared with the control C3 was 14.8 for Example 6, 10.8 for Example 7, 9.1 for Example 8, and 19.0 for Example 9.

Examples 11-12

10 These examples disclose dispersions of Drug 1 in a neutral polymer with added buffer. Examples 11 and 12 were prepared with Drug 1, as described in Example 1, with the following exceptions: for Example 11, the solution to be spray-dried consisted of 1.25 wt% Drug 1, 0.513 wt% sodium acetate, and 3.75 wt% HPMC in methanol/water (9/1); and for Example 12, the solution
15 consisted of 1.25 wt% Drug 1, 0.32 wt% sodium bicarbonate, and 3.75 wt% HPMC in methanol/water (9/1). The solution variables are summarized in Table 9.

Table 12

Ex.	Aqueous-Soluble Polymer	Drug 2 Conc. In the Dispersion (wt%)	Potency Day 7 at room temp/0%RH (wt%)	Degree of Degradation (wt%)
14	PVP	10	84	16
C4	HPMCAS-LF	10	45	55
C5	HPMCP	10	53	47

5

As the data show, the dispersion of Example 14 comprised of the acid-sensitive drug and neutral polymer PVP had improved chemical stability relative to dispersions formed using either of the acidic polymers HPMCAS-LF or HPMCP. Example 14 had a relative degree of improvement when compared with control C4 of 3.4, and a relative degree of improvement when compared with control C5 of 2.9.

10

Example 16

This example measures the concentration-enhancing properties of the PVP dispersion of Example 14. Dissolution tests were performed, as described in Example 4 with the following exceptions: the PBS solution contained 7.3 mM sodium taurocholic acid and 1.4 mM 1-palmitoyl-2-oleyl-sn-glycero-3-phosphocholine (model fasted duodenal (MFD) solution). 1.8 mg of dispersion was added to 1.8 mL of test media. For Control C6, 0.18 mg of the crystalline form of the Drug 2 alone was used without further processing and added to 1.8 mL of MFD solution. The results are shown in Table 13 below.

15

20

Table 13

Example	Time (min)	Drug 2 ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
14	0	0	0
	4	57	100
	10	55	500
	20	55	1,000
	40	52	2,100
	90	54	4,700
C6	0	0	0
	4	2	0
	10	4	0
	20	4	100
	40	4	100
	90	7	400

Example 14

This example discloses dispersions of a second acid-sensitive drug, quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-cyclohexyl)-2-hydroxy-4-hydroxycarbonyl-butyl]-amide (Drug 2). For Example 14, a dispersion of Drug 2 and PVP was made by first preparing a solution containing 0.030 wt% drug and 0.27 wt% PVP in methanol/acetone (1/9 wt/wt), and then spraying the solution into a drying chamber using an atomizing spray nozzle in the manner described in Example 1.

For Control C4, a dispersion of Drug 2 and HPMCAS-LF was made by preparing a solution containing 0.030 wt% Drug 2 and 0.27 wt% HPMCAS-LF in acetone, and spray-drying as described above. For Control C5, a dispersion of Drug 2 with the acidic polymer hydroxypropyl methyl cellulose phthalate (HPMCP) was made by preparing a solution containing 0.030 wt% drug and 0.27 wt% HPMCP in acetone, and spray-drying as described above. Table 11 summarizes the solution variables.

Table 11

Ex.	Drug Mass (mg)	Drug No.	Polymer Mass (mg)	Polymer	Solvent Mass (g)	Solvent (wt/wt)
14	3	2	27	PVP	10	methanol/acetone (1/9)
C4	3	2	27	HPMCAS	10	acetone
C5	3	2	27	HPMCP	10	acetone

Example 15

This example assesses the stability of the dispersion of Example 14. The dispersions of Example 14 and Controls C4 and C5 were stored in controlled atmosphere chambers for 7 days at room temperature and 0% RH. Potencies of the dispersions before and after storage were determined using HPLC as in Example 3, with the following exceptions. An HP ODS Hypersil column was used with a mobile phase of 60 vol% 0.02M KH₂PO₄ (pH 3.0) and 40 vol% acetonitrile. UV detection was measured at 248 nm. The results are shown in Table 12 below.

Table 15

Ex.	Drug Mass (mg)	Drug No.	Polymer Mass (mg)	Polymer	Solvent Mass (g)	Solvent (wt/wt)
17	5	3	45	HPMC	10	methanol
18	3	3	27	PVP	10	acetone/ methanol (9/1)
C7	3	3	27	HPMCAS-LF	10	acetone
C8	3	3	27	HPMCP	10	acetone

5

Example 19

This example demonstrates the relative chemical stability of the dispersions of Examples 17 and 18. The dispersions of Examples 17 and 18, and Controls C7 and C8, were placed in controlled atmosphere chambers for 6 days at room temperature and 0% RH (the HPMC dispersion of Example 16 was stored for 4 days). Potencies of the dispersions before and after storage were determined using HPLC, using the method described in Example 15 above. The results are shown in Table 16 below.

10

Table 16

15

Ex.	Aqueous-Soluble Polymer	Drug 3 Conc. in the Dispersion (wt%)	Potency Day 6 at room temp/0%RH (wt%)	Degree of Degradation (wt%)
17	HPMC	10	92 (4 days)	8
18	PVP	10	95	5
C7	HPMCAS-LF	10	46	54
C8	HPMCP	10	56	44

The data show that dispersions of the acid-sensitive drug and neutral polymers were more stable than dispersions formed with the acidic polymers HPMCAS-LF and HPMCP. The dispersion of Example 18 had a relative degree of improvement of 10.8 when compared with the control C7, and 8.8 when compared with the control C8.

20

The concentrations of Drug 2 obtained in these samples were used to determine the values of $C_{\max 90}$ and AUC_{90} . The results are shown in Table 14. As can be seen from the data, dispersions of Example 14 provided an 8-fold improvement in $C_{\max 90}$ and a 12-fold improvement in AUC_{90} .

Table 14

Example	$C_{\max 90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
14	57	4700
C6	7	400

Examples 17-18

These examples disclose dispersions of a third acid-sensitive drug, quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide (Drug 3). For Example 17, a dispersion of Drug 3 and HPMC was made by preparing a solution containing 0.050 wt% Drug 3 and 0.45 wt% HPMC in methanol, and spraying the solution into a drying chamber using an atomizing spray nozzle, as described in Example 1. For Example 18, a dispersion of Drug 3 with PVP was made by preparing a solution containing 0.030 wt% Drug 3 and 0.27 wt% PVP in methanol/acetone (1/9 wt/wt), and spray-drying as described above.

For Control C7, a dispersion of Drug 3 and HPMCAS-LF was made by preparing a solution containing 0.030 wt% drug and 0.27 wt% HPMCAS-LF in acetone, and spray-drying as described above. For Control C8, a dispersion of Drug 3 and HPMCP was made by preparing a solution containing 0.030 wt% drug and 0.27 wt% HPMCP in acetone, and spray-drying as described above. Table 15 summarizes the spray-drying variables.

Example 20

This example demonstrates the concentration enhancement provided by the dispersion of Example 18. Dissolution tests were performed, as described in Example 4 with the following exceptions: samples were tested in MFD Solution, and 1.8 mg of dispersion was added to 1.8 mLs of MFD Solution. Control C9 was simply 0.18 mg of the crystalline Drug 3 alone add to 1.8 mL of MFD solution. The results are shown in Table 17 below.

Table 17

Example	Time (min)	Drug 3 ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
17	0	0	0
	4	88	200
	10	83	700
	20	80	1,500
	40	79	3,100
	90	77	7,000
C9	0	0	0
	4	4	0
	10	6	0
	20	8	100
	40	11	300
	90	13	900

The concentrations of Drug 3 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown in Table 18.

Table 18

Example	$C_{\text{max}90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
18	88	7100
C9	13	900

As can be seen from the data, dispersions of Drug 3, an acid-sensitive drug, and the neutral polymer, PVP, provided concentration-enhancement in an aqueous environment of use. The dispersion of Example 18 provided a $C_{\text{max}90}$ that was 6.8-fold, and an AUC_{90} that was 7.9-fold that of the control C9.

Examples 21-29

These examples disclose dispersions formed from various low-solubility drugs and neutral amphiphilic, hydroxyl-functional vinyl copolymers, particularly

vinyl acetate/vinyl alcohol copolymers (hereinafter "VAVAC"). For Examples 21 to 24, the dispersions were made using Drug 1. For Examples 25 to 27, the dispersions were made using 5-chloro-1H-indole-2-carboxylic acid [(1S) - benzyl-3-((3R, 4S)-dihydroxypyrrolidin-1-yl)-(2R)-hydroxy-3-oxypropyl] amide ("Drug 4"). For Examples 28 to 29, the dispersions were made using (+)-N-{3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl}-N-hydroxyurea ("Drug 5").

Several different grades of VAVAC were used. For Examples 23, 25 and 28, the VAVAC, referred to herein as "VAVAC-20%," was 80% hydrolyzed (meaning that about 20% of the vinyl repeat units were acetylated), and had an average molecular weight of 9,000-10,000 daltons. The VAVAC-20% was supplied by Aldrich Chemical Company of lot #36,062-7.

For Examples 21, 26 and 29, the VAVAC, referred to herein as "VAVAC-12%," was approximately 88% hydrolyzed (approximately 11-13% of the repeat units were acetylated), and had an average molecular weight of 13,000-23,000 daltons. VAVAC-12% was supplied by Aldrich Chemical Company, lot #36,317-0.

For Examples 22, 24 and 27, the VAVAC, referred to herein as "VAVAC-2%," was 98% hydrolyzed (approximately 2% of the repeat units were acetylated) and had an average molecular weight of 13,000-23,000 daltons. VAVAC-2% was supplied by Aldrich Chemical Company, lot #34,840-6.

To form the dispersions of Examples 21-29, solutions containing drug and polymer in a solvent were spray-dried by pumping each solution into a "mini" spray-drier apparatus as described in Example 1.

Table 19 summarizes the variables for the solutions of Examples 21-29.

190

Table 19

Ex.	Drug	Drug Mass (mg)	Drug Conc.* (%)	Polymer	Polymer Mass (mg)	Spray Solvent (wt/wt)	Solvent Mass (g)
21	1	62	25	VAVAC-12%	186	MeOH/H ₂ O (4/1)	18
22	1	692	75	VAVAC-2%	231	MeOH/H ₂ O (1/1)	50
23	1	664	50	VAVAC-20%	664	MeOH/H ₂ O (4/1)	105
24	1	522	25	VAVAC-2%	1565	MeOH/H ₂ O (4/1)	110
25	4	110	25	VAVAC-20%	330	MeOH/H ₂ O (4/1)	33.8
26	4	126	25	VAVAC-12%	376	MeOH/H ₂ O (1.8/1)	50
27	4	78	25	VAVAC-2%	233	MeOH/H ₂ O (1/1)	20.6
28	5	37	10	VAVAC-20%	330	MeOH/H ₂ O (4/1)	33.8
29	5	42	10	VAVAC-12%	376	MeOH/H ₂ O (1.8/1)	50

* Drug concentration in the final dispersion.

5

Example 30

In this example the chemical stability of the dispersions of Examples 23 and 24 was assessed. The dispersions were stored for three weeks at 40°C and 75% RH, then analyzed for Drug 1 potency by HPLC using the method described in Example 3. Results are shown in Table 20, as are the results for Control C1, a dispersion of Drug 1 and the acidic polymer HPMCAS-LF (prepared in Example 1).

10

15

Table 20

Ex.	Polymer	Drug 1 Conc. in the Disper- sion (wt%)	Pot. Before Storage (%)	Potency After 3 weeks @ 40°C/75 RH (%)	Degree of Degrad- ation @ 3 wks (wt%)	Relative Degree of Improve- ment
23	VAVAC- 20%	50	98	97	3	>33
24	VAVAC- 2%	25	98	96	4	>25
C1	HPMCAS- LF	25	94	<1*	>99*	-

*Values were obtained after 2 weeks.

5 As the data show, the dispersions of Examples 23 and 24 formed with VAVAC have improved chemical stability when compared with Control C1, showing a relative degree of improvement of >33 for Example 23 and >25 for Example 24.

Example 31

10 *In vitro* dissolution tests were performed for Examples 21-24, using the procedure described in Example 4, with the following exceptions: 14.4 mg of the dispersions of Example 21, 4.8 mg of the dispersion Example 22, 7.2 mg of the dispersion of Example 23, and 14.4 mg of the dispersion of Example 24 were added to respective tubes. 1.8 mL of PBS was added to each tube to achieve a theoretical total Drug 1 concentration of approximately 2000 µg/mL if all of the drug were to dissolve completely. For comparison, a Control C10 of 1.8 mg of crystalline Drug 1 was added to 1.8 mL of PBS. The results are shown in Table 21.

Table 21

Example	Time (min)	Drug 1 Concentration ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
21	0	0	0
	4	1554	3,100
	10	1860	13,300
	20	1474	30,000
	40	744	52,200
	90	685	87,900
	180	632	147,200
	1200	580	790,000
22	0	0	0
	4	809	1,600
	10	1019	7,100
	20	1162	18,000
	40	1211	41,700
	90	1273	103,835
	1200	1117	1,430,292
23	0	0	0
	4	1698	3,400
	10	1685	13,500
	20	1076	27,400
	40	608	44,200
	90	552	73,200
	1200	499	656,500
24	0	0	0
	4	1081	2,200
	10	1279	9,200
	20	1334	22,300
	40	1342	49,100
	90	662	99,200
	1200	478	731,900
C10	0	0	0
	4	241	500
	10	256	2,000
	20	299	4,700
	40	260	10,300
	90	295	25,500
	180	298	49,400
	1200	313	361,000

- 5 The concentrations of Drug 1 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown below in Table 22.

Table 22

Example	C _{max90} (µg/mL)	AUC ₉₀ (min*µg/mL)
21	1860	87,900
22	1273	103,800
23	1698	73,200
24	1342	99,200
C10	299	25,500

5 As can be seen from the data, the dispersions of Examples 21-24 provided concentration-enhancement over that of Control C10 (crystalline Drug 1 alone). Example 21 provided a C_{max90} that was 6.2-fold and an AUC₉₀ that was 3.4-fold that of Control C10; Example 22 provided a C_{max90} that was 4.3-fold, and an AUC₉₀ that was 4.1-fold that of the Control C10; Example 23 provided a C_{max90} that was 5.7-fold and an AUC₉₀ that was 2.9-fold that of Control C10; and
10 Example 24 provided a C_{max90} that was 4.5-fold and an AUC₉₀ that was 3.9-fold that of Control C10.

Example 32

This example demonstrates a composition comprising a dispersion of an acid-sensitive drug and a neutral polymer mixed with an acidic concentration-enhancing polymer, namely HPMCAS-HF. Example 32 was prepared by mixing
15 the dispersion of Example 21 with HPMCAS-HF. An *in vitro* dissolution test was performed using the procedure described in Example 4, with the following exceptions: 14.4 mg of the dispersion of Example 21 and 14.4 mg of HPMCAS-HF were added to 1.8 mL of PBS solution for a total theoretical Drug 1
20 concentration of approximately 2000 µg/mL if all of the drug were to dissolve completely. The results are shown in Table 23.

Table 23

Example	Time (min)	Drug 1 Concentration ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
32	0	0	0
	4	1365	2,700
	10	1441	11,100
	20	1399	25,300
	40	1577	55,100
	90	1541	133,100
	180	1648	276,600
	1200	1863	2,021,700

5

The concentrations of Drug 1 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown below in Table 24.

10

Table 24

Example	$C_{\text{max}90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
32	1577	133,100
21	1860	87,900

15

As can be seen from the data, the composition of Example 32 (*i.e.*, the dispersion of Example 21 and HPMCAS) had an improved AUC_{90} relative to the dispersion of Example 21 alone. Results of dissolution tests for Example 21 were shown previously in Table 22, however, they are shown again in Table 24 for comparison.

Example 33

20

In vitro dissolution tests were performed for Examples 25-27 using the procedure described in Example 4, with the following exceptions: 14.4 mg each of the dispersions of Examples 25-27 were added to separate, respective centrifuge tubes. 1.8 mL PBS was added to each tube. For comparison, Control C11 was prepared consisting of 3.6 mg of crystalline Drug 4 added to 1.8 mL PBS. Drug 4 concentration in the samples was determined using HPLC. A Zorbax SB C_{18} HPLC column was used with a mobile phase of 35 vol.% water

25

and 65 vol.% methanol. UV detection was measured at 297 nm. The results are shown in Table 25.

5

Table 25

Example	Time (min)	[Drug 4] ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
25	0	0	0
	4	1580	3,200
	10	1581	12,600
	20	1570	28,400
	40	1594	60,000
	90	1476	136,000
	1200	248	1,093,600
26	0	0	0
	4	1957	3,900
	10	1993	15,800
	20	1948	35,500
	40	1980	74,700
	90	1959	173,200
	1200	163	1,350,900
27	0	0	0
	4	735	1,500
	10	733	5,900
	20	707	13,100
	40	691	27,100
	90	676	61,200
	1200	147	518,000
C11	0	0	0
	4	131	300
	10	114	1,000
	20	124	2,200
	40	107	4,500
	90	126	10,300
	1200	72	120,200

The concentrations of Drug 4 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown below in Table 26.

10

15

Table 26

Example	C _{max90} (µg/mL)	AUC ₉₀ (min*µg/mL)
25	1594	136,800
26	1993	173,200
27	735	61,200
C11	131	10,300

5

As can be seen from the data, the dispersions of Examples 25-27 provided concentration-enhancement over that of Control C11, the crystalline Drug 4 alone. The dispersion of Example 25 provided a C_{max90} that was 12-fold and an AUC₉₀ that was 13-fold that of Control C11; Example 26 provided a C_{max90} that was 15-fold and an AUC₉₀ that was 17-fold that provided by Control C11; and Example 27 provided a C_{max90} that was 6-fold, and an AUC₉₀ that was 6-fold that of Control C11.

10

Example 34

15

In vitro dissolution tests were performed for Examples 28-29, using the procedure described in Example 4, with the following exceptions: 9 mg each of the dispersions of Examples 28-29 were added to separate respective centrifuge tubes. 1.8 mL of MFD solution was then added to each tube. For comparison, Control C12 was prepared consisting of 0.36 mg of crystalline Drug 5 added to 1.8 mL of MFD solution for a total theoretical concentration of 200 µg/mL if all of the drug dissolved. Drug 5 concentration in the samples was determined using HPLC. A Waters Symmetry C₁₈ HPLC column was used with a mobile phase of 50 vol.% (0.3 vol.% glacial acetic acid, 0.2 vol.% TEA in water) and 50 vol.% acetonitrile. UV detection was measured at 260 nm. The results are shown in Table 27.

20

25

30

Table 27

Example	Time (min)	Drug 5 Concentration ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
28	0	0	0
	4	211	400
	10	180	1,600
	20	168	3,300
	40	155	6,600
	90	144	14,000
	1200	139	171,100
29	0	0	0
	4	210	400
	10	192	1,600
	20	186	3,500
	40	175	7,100
	90	169	15,700
	1200	155	195,500
C12	0	0	0
	4	9	0
	10	15	100
	20	21	300
	40	27	800
	90	32	2,200
	1200	42	43,300

5 The concentrations of Drug 5 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown below in Table 28. While the amount of Drug 5 used in the Control C12 was less than the amount of Drug 5 in the test compositions of Examples 28 and 29, nevertheless a sufficient amount of Drug 5 in excess of the solubility of Drug 5 was used in Control C12 in order to provide an accurate measure of the performance of Examples 28 and 29 relative to crystalline drug.

Table 28

Example	$C_{\text{max}90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
28	211	14,000
29	210	15,700
C12	32	2,200

15 As can be seen from the data, the dispersions of Examples 28 - 29 provided concentration-enhancement over that of crystalline Drug 5 alone. Example 28

provided a C_{max90} that was 6.6-fold and an AUC_{90} that was 6.4-fold that of Control C12; and Example 29 provided a C_{max90} that was 6.6-fold and an AUC_{90} that was 7.1-fold that provided by Control C12.

Examples 35-38

- 5 Examples 35 and 36 were dispersions of Drug 4 prepared with the dispersion polymers PVP and HPMC respectively. For Examples 37 and 38, dispersions of Drug 5 and the polymers PVP and HPMC were prepared. To form the dispersions of Examples 35-38, solutions containing drug and polymer in a solvent were spray-dried by pumping each solution into a "mini" spray-drier apparatus as described in Example 1. Table 29 summarizes the variables for the solutions used to prepare Examples 35-38.

Table 29

Ex.	Drug	Drug Mass (mg)	Drug Conc.* (%)	Polymer	Polymer Mass (mg)	Spray Solvent (wt/wt)	Solvent Mass (g)
35	4	50	25	PVP	150	acetone	15
36	4	50	25	HPMC	150	MeOH/ water (9/1)	15
37	5	50	10	PVP	450	Acetone	80
38	5	50	10	HPMC	450	MeOH/ acetone (1/1)	80

* Drug concentration in the final dispersion

15

Example 39

- In vitro* dissolution tests were performed for Examples 35-38, using the procedures described in Example 4, with the following exceptions: 14.4 mg each of Example 35 and 36 were added to separate respective test tubes to which 1.8 mL of PBS solution was added, so that the total theoretical Drug 4 concentration was approximately 2000 $\mu\text{g/mL}$ if all of Drug 4 were to dissolve completely; 3.6 mg each of Examples 37 and 38 were added in separate respective test tubes to 1.8 mL of MFD solution, to provide a total theoretical Drug 5 concentration of approximately 200 $\mu\text{g/mL}$ if all of Drug 5 were to dissolve completely. Drug 4 concentration in the samples was determined as in Example 33. Drug 5 concentration in the samples was determined as in Example 34. The results are shown in Table 30.

Table 30

Example	Time (min)	[Drug 4] ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
35	0	0	0
	4	771	1,500
	10	773	6,200
	20	781	13,900
	40	753	29,300
	90	703	65,700
	1200	540	755,500
36	0	0	0
	4	420	1,000
	10	422	3,400
	20	429	7,600
	40	447	16,400
	90	447	38,700
	1200	319	463,900
37	0	0	0
	4	121	200
	10	113	900
	20	111	7,100
	40	111	4,300
	90	120	10,100
	1200	106	135,500
38	0	0	0
	4	429	900
	10	347	3,200
	20	280	6,300
	40	229	11,400
	90	257	23,600
	1200	164	237,200

5 The concentrations of Drug 4 and Drug 5 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results for dispersions containing Drug 4 are shown below in Table 31, and for dispersions containing Drug 5 in Table 32.

Table 31

Example	C_{max90}	AUC_{90}
35	781	65,700
36	447	38,700
C11	131	10,300

5

Table 32

Example	C_{max90}	AUC_{90}
37	121	10,100
38	429	23,600
C12	32	2,200

As can be seen from the data, the dispersions of Examples 35 and 36 provided superior concentration-enhancement relative to the Control C11 and Examples 37 and 38 provided superior concentration-enhancement relative to the Control C12.

The terms and expressions which have been employed in the foregoing specification are used therein as terms of description and not of limitation, and there is no intention, in the use of such terms and expressions, of excluding equivalents of the features shown and described or portions thereof, it being recognized that the scope of the invention is defined and limited only by the claims which follow.

201

What is claimed is:

1. A pharmaceutical composition comprising a solid amorphous dispersion of an acid-sensitive drug and a neutral dispersion polymer, wherein said
5 composition provides improved chemical stability relative to a control acidic dispersion comprised of an equivalent quantity of said drug and an acidic polymer.
2. A pharmaceutical composition comprising a solid amorphous dispersion
10 comprising a low-solubility drug, a neutral dispersion polymer, and an excipient selected from the group consisting of a base and a buffer, wherein said composition provides improved chemical stability relative to a control composition, and wherein said control composition comprises an equivalent quantity of a dispersion of said drug and said neutral dispersion polymer but
15 free from said base and said buffer.
3. The composition of any one of claims 1-2 wherein said drug when present in a control acidic dispersion and stored for a period of six months at 40°C and 75% relative humidity has a degree of degradation of at least 0.01%.
20
4. The composition of any one of claims 1-2 wherein said drug has one or more functional groups selected from the group consisting of sulfonyl ureas, hydroxamic acids, hydroxy amides, carbamates, acetals, hydroxy ureas, esters, and amides.
25
5. The composition of any one of claims 1-2 wherein said drug is selected from the group consisting of quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3- fluorobenzyl-2(S),7-dihydroxy-7-methyl-octyl]amide, quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-
30 amide, quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide, (+)-N-{3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl}-N-hydroxyurea, omeprazole, etoposide, famotidine, erythromycin, quinapril, lansoprazole, and progabide.
- 35 6. The composition of any one of claims 1-2 wherein said dispersion polymer is selected from the group consisting of hydroxypropyl methyl cellulose

acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and hydroxyethyl ethyl cellulose.

- 5 7. The composition of any one of claims 1-2 wherein said dispersion polymer is selected from the group consisting of vinyl polymers and copolymers having one or more substituents selected from the group consisting of hydroxyl-containing repeat units, alkylacyloxy-containing repeat units, or cyclicamido-containing repeat units, polyvinyl alcohols that have at least a portion of their
- 10 repeat units in the unhydrolyzed form, polyvinyl alcohol polyvinyl acetate copolymers, polyethylene glycol, polyethylene glycol polypropylene glycol copolymers, polyvinyl pyrrolidone, polyethylene polyvinyl alcohol copolymers, and polyoxyethylene-polyoxypropylene block copolymers.
- 15 8. The composition of any one of claims 1-2 wherein said dispersion polymer is concentration-enhancing and said drug has a minimum solubility in aqueous solution in the absence of said dispersion polymer of less than 1 mg/mL at any pH of from about 1 to about 8.
- 20 9. The composition of any one of claims 1-2 wherein said dispersion polymer is present in an amount sufficient so that said dispersion provides to a use environment at least one of:
- 25 (a) a maximum concentration of said drug in said use environment that is at least 1.25-fold that provided by a second control composition comprising an equivalent quantity of said drug and free from said dispersion polymer;
- 30 (b) an area under the concentration versus time curve for any period of at least 90 minutes between the time of introduction into said use environment and about 270 minutes following introduction to said use environment that is at least 1.25-fold that provided by said second control composition comprising an equivalent quantity of said drug and free from said dispersion polymer; and
- 35 (c) a relative bioavailability that is at least 1.25 relative to said second control composition comprising an equivalent quantity of said drug and free from said dispersion polymer.

10. A pharmaceutical composition comprising a solid amorphous dispersion of a low-solubility drug and a neutral dispersion polymer, wherein said neutral dispersion polymer comprises a vinyl copolymer having hydrophilic hydroxyl-containing repeat units and hydrophobic repeat units.
- 5
11. The composition of any one of claims 1-2 and 10 wherein said dispersion has a pH from about 6 to about 10.
12. The composition of any one of claims 1-2 and 10 wherein said
10 composition further comprises a second polymer, said dispersion is free from at least a portion of said second polymer, and said second polymer is selected from the group consisting of hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, hydroxyethyl cellulose
15 acetate succinate, hydroxypropyl methyl cellulose phthalate, hydroxyethyl methyl cellulose acetate succinate, hydroxyethyl methyl cellulose acetate phthalate, carboxyethyl cellulose, carboxymethyl cellulose, cellulose acetate phthalate, methyl cellulose acetate phthalate, ethyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, hydroxypropyl methyl cellulose
20 acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, hydroxypropyl methyl cellulose acetate succinate phthalate, hydroxypropyl methyl cellulose succinate phthalate, cellulose propionate phthalate, hydroxypropyl cellulose butyrate phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate, ethyl cellulose acetate trimellitate, hydroxypropyl
25 cellulose acetate trimellitate, hydroxypropyl methyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate succinate, cellulose propionate trimellitate, cellulose butyrate trimellitate, cellulose acetate terephthalate, cellulose acetate isophthalate, cellulose acetate pyridinedicarboxylate, salicylic acid cellulose acetate, hydroxypropyl salicylic acid cellulose acetate,
30 ethylbenzoic acid cellulose acetate, hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl phthalic acid cellulose acetate, ethyl nicotinic acid cellulose acetate, ethyl picolinic acid cellulose acetate, carboxylic acid functionalized polymethacrylates, carboxylic acid functionalized polyacrylates, amine-
35 functionalized polyacrylates, amine-fuctinoalized polymethacrylates, proteins, and carboxylic acid functionalized starches.

13. The composition of any one of claims 1-2 and 10 wherein said composition further comprises a second polymer, said dispersion is free from at least a portion of said second polymer, and said second polymer is selected from the group consisting of hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, hydroxyethyl ethyl cellulose, and vinyl polymers and copolymers having one or more substituents selected from the group consisting of hydroxyl-containing repeat units, alkylacyloxy-containing repeat units, or cyclicamido-containing repeat units, polyvinyl alcohols that have at least a portion of their repeat units in the unhydrolyzed form, polyvinyl alcohol polyvinyl acetate copolymers, polyethylene glycol, polyethylene glycol polypropylene glycol copolymers, polyvinyl pyrrolidone, polyethylene polyvinyl alcohol copolymers, and polyoxyethylene-polyoxypropylene block copolymers.
14. The composition of any one of claims 1, 2 and 10-13 wherein said drug in said composition has a relative degree of improvement in chemical stability of at least 1.25.
15. The composition of any of claims 1, 2 and 10-13 wherein the drug comprises [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester; [2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester; or [2R, 4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester.

INTERNATIONAL SEARCH REPORT

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K9/14 A61K9/16		Inte Application No PC1/1B 02/01783
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SHAH J C ET AL: "PREFORMULATION STUDY OF ETOPOSIDE: II. INCREASED SOLUBILITY AND DISSOLUTION RATE BY SOLID-SOLID DISPERSIONS" INTERNATIONAL JOURNAL OF PHARMACEUTICS, AMSTERDAM, NL, vol. 113, no. 1, 2 January 1995 (1995-01-02), pages 103-111, XP001040972 ISSN: 0378-5173 page 104-105, paragraph entitled "2.2. Preparation of coprecipitates"; page 109-110, paragraph entitled "4. Discussion" paragraph bridging page 103 and 104	1-9, 11, 14
A	--- -/--	1, 3-9, 11-14
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.		
<input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents :		
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 1 August 2002		Date of mailing of the international search report 08/08/2002
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Borst, M

Form PCT/ISA/210 (second sheet) (July 1992)

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 64414 A (BREITENBACH JOERG ; BASF AG (DE); BINDER RUDOLF (DE); LIEPOLD BERND) 2 November 2000 (2000-11-02) page 2, line 31 - page 3, line 21; example 2, 3	1-9, 11, 14
A	page 1, line 30-37;	1, 3-9, 11-14
X	----- SARISUTA N ET AL: "PHYSICO-CHEMICAL CHARACTERIZATION OF INTERACTIONS BETWEEN ERYTHROMYCIN AND VARIOUS FILM POLYMERS" INTERNATIONAL JOURNAL OF PHARMACEUTICS, AMSTERDAM, NL, vol. 186, no. 2, 20 September 1999 (1999-09-20), pages 109-118, XP000974319 ISSN: 0378-5173 page 110-111, paragraph entitled "2.3. Preparation of the drug-polymer mixtures"	1-6, 8, 9, 11, 14
X	----- US 4 673 564 A (KAWATA HIROITSU ET AL) 16 June 1987 (1987-06-16) cited in the application column 1, line 47-51; example 1-3, 9	1-4, 6-9, 11-14
X	----- EP 1 027 886 A (PFIZER PROD INC) 16 August 2000 (2000-08-16) '0027!', '0082!', '0083!	1, 2, 6, 7, 10
X	----- WO 97 08950 A (FUISZ TECHNOLOGIES LTD) 13 March 1997 (1997-03-13) page 6, line 4-9; examples 1-7	1-4, 7-9, 11, 14
P, X	----- WO 02 11710 A (CURATOLO WILLIAM JOHN ; FRIESEN DWAYNE THOMAS (US); GUMKOWSKI MICHA) 14 February 2002 (2002-02-14) example 14, 16	1-4, 6-9, 11, 14, 15
A	----- WO 99 14204 A (SEARLE & CO ; SIKORSKI JAMES A (US)) 25 March 1999 (1999-03-25) page 1, line 1-15; page 84, line 27-29	1-15
X	----- WO 00 17166 A (WESTER RONALD THURE ; PFIZER PROD INC (US); DENINNO MICHAEL PAUL (U) 30 March 2000 (2000-03-30) page 2, line 22 - page 6, line 2	15
A	----- WO 99 40061 A (KATH JOHN C ; BROWN MATTHEW F (US); PFIZER PROD INC (US); POSS CHRI) 12 August 1999 (1999-08-12) page 2, line 21 - page 3, line 19	5
	----- -/-	

INTERNATIONAL SEARCH REPORT

Int. Application No
PCT/18 02/01783

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>LEUNER C ET AL: "Improving drug solubility for oral delivery using solid dispersions"</p> <p>EUROPEAN JOURNAL OF PHARMACEUTICS AND BIOPHARMACEUTICS, ELSEVIER SCIENCE PUBLISHERS B.V., AMSTERDAM, NL, vol. 50, no. 1, 3 July 2000 (2000-07-03), pages 47-60, XP004257179</p> <p>ISSN: 0939-6411</p> <p>whole document</p> <p style="text-align: center;">-----</p>	1-15

1

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

page 3 of 3

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/IB 02/01783

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0064414	A	02-11-2000	DE	19918434 A1	26-10-2000
			WO	0064414 A2	02-11-2000
			EP	1173152 A2	23-01-2002
US 4673564	A	16-06-1987	JP	1272484 C	11-07-1985
			JP	56133217 A	19-10-1981
			JP	59048810 B	29-11-1984
			JP	1591530 C	30-11-1990
			JP	2009007 B	28-02-1990
			JP	56049314 A	02-05-1981
			CA	1146866 A1	24-05-1983
			CH	648484 A5	29-03-1985
			DE	3024858 A1	22-01-1981
			ES	493152 D0	16-11-1981
			ES	8200557 A1	01-02-1982
			ES	499854 D0	01-06-1982
			ES	8205125 A1	16-09-1982
			FR	2460667 A1	30-01-1981
			FR	2489146 A1	05-03-1982
			GB	2053681 A , B	11-02-1981
			IT	1132169 B	25-06-1986
			SE	448342 B	16-02-1987
			SE	8004938 A	06-01-1981
			US	4343789 A	10-08-1982
			US	4404183 A	13-09-1983
EP 1027886	A	16-08-2000	BR	0000346 A	21-08-2001
			EP	1027886 A2	16-08-2000
			JP	2000229887 A	22-08-2000
WO 9708950	A	13-03-1997	AU	7106996 A	27-03-1997
			CA	2231050 A1	13-03-1997
			EP	0952770 A1	03-11-1999
			JP	11514979 T	21-12-1999
			WO	9708950 A1	13-03-1997
			US	6391338 B1	21-05-2002
WO 0211710	A	14-02-2002	AU	2914202 A	18-02-2002
			WO	0211710 A2	14-02-2002
WO 9914204	A	25-03-1999	AU	9297798 A	05-04-1999
			WO	9914204 A1	25-03-1999
WO 0017166	A	30-03-2000	AU	5439899 A	10-04-2000
			BR	9913842 A	12-06-2001
			EP	1114033 A1	11-07-2001
			WO	0017166 A1	30-03-2000
			US	6147090 A	14-11-2000
			US	6395751 B1	28-05-2002
WO 9940061	A	12-08-1999	AP	992 A	06-08-2001
			AU	1778999 A	23-08-1999
			BG	104726 A	30-04-2001
			BR	9907655 A	24-10-2000
			CA	2320388 A1	12-08-1999
			CN	1290255 T	04-04-2001
			EP	1051405 A2	15-11-2000
			HR	20000529 A1	31-08-2001

Form PCT/ISA/210 (patent family annex) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/IB 02/01783

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9940061	A	WO 9940061 A2	12-08-1999
		JP 2002502839 T	29-01-2002
		NO 20003965 A	03-10-2000
		PL 342817 A1	02-07-2001
		SK 11262000 A3	18-01-2001
		TR 200002248 T2	21-12-2000
		ZA 9900873 A	04-08-2000

Form PCT/ISA/210 (patent family annex) (July 1992)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2003 (03.01.2003)

PCT

(10) International Publication Number
WO 03/000235 A1

(51) International Patent Classification⁷: **A61K 9/14, 9/16**

[US/US]; 1511 N.W. Jacksonville Avenue, Bend, OR 97701 (US).

(21) International Application Number: PCT/TB02/01783

(74) Agents: **LUMB, J., Trevor** et al.; Pfizer Inc., 201 Tabor Road, Morris Plains, NJ 07950 (US).

(22) International Filing Date: 10 May 2002 (10.05.2002)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
60/300,255 22 June 2001 (22.06.2001) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/300,255 (CON)
Filed on 22 June 2001 (22.06.2001)

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **PFIZER PRODUCTS INC.** [US/US]; Eastern Point Road, Groton, CT 06340 (US).

Published:

— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **FRIESEN, Dwayne, Thomas** [US/US]; 60779 Currant Way, Bend, OR 97702 (US). **GUMKOWSKI, Michael, Jon** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **KETNER, Rodney, James** [US/US]; 1615 N.W. Trenton Avenue, Bend, OR 97701 (US). **LORENZ, Douglas, Alan** [US/US]; 61332 S.E. King Jehu Way, Bend, OR 97702 (US). **NIGHTINGALE, James, Alan, Schriver** [US/US]; 62900 Santa Cruz, Bend, OR 97701 (US). **SHANKER, Ravi, Mysore** [US/US]; Pfizer Global Research and Development, Eastern Point Road, Groton, CT 06340 (US). **WEST, James, Blair**

(48) Date of publication of this corrected version:

24 December 2003

(15) Information about Correction:

see PCT Gazette No. 52/2003 of 24 December 2003, Section II

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/000235 A1

(54) Title: PHARMACEUTICAL COMPOSITIONS OF DISPERSIONS OF DRUGS AND NEUTRAL POLYMERS

(57) Abstract: In one aspect, pharmaceutical compositions comprising dispersions of an acid-sensitive drug and a neutral dispersion polymer are disclosed. The acid-sensitive drug has improved chemical stability relative to dispersions of the drug and acidic polymers. In another aspect, pharmaceutical compositions of low solubility drugs and amphiphilic, hydroxy-functional vinyl copolymers are disclosed.

PHARMACEUTICAL COMPOSITIONS OF DISPERSIONS OF DRUGS AND NEUTRAL POLYMERS

This application claims the benefit of priority of provisional Patent
5 Application Serial No. 60/300,255 filed June 22, 2001, which is incorporated
herein by reference in its entirety for all purposes.

BACKGROUND OF THE INVENTION

This invention relates to pharmaceutical compositions comprised of
10 amorphous dispersions of drugs and neutral polymers that provide either
improved chemical stability, concentration-enhancement, or both improved
chemical stability and concentration-enhancement.

It is sometimes desired to form a solid amorphous dispersion of a drug
and a polymer. One reason for forming dispersions is that the aqueous
15 concentration of a poorly soluble drug may be improved by forming an
amorphous dispersion of the drug and a polymer. For example, Curatolo, et al.,
EP 0 901 786 A2 disclose forming pharmaceutical spray dried dispersions of
sparingly soluble drugs and the polymer hydroxypropyl methyl cellulose acetate
succinate. The spray dried dispersions disclosed in Curatolo et al. provide
20 superior aqueous concentration relative to dispersions formed from other
methods and relative to the crystalline drug alone.

Similarly, others have recognized the enhancement in aqueous
concentration afforded by dispersing a drug in a polymer. Nakamichi, et al.,
U.S. Patent No. 5,456,923 disclose solid dispersions formed by twin-screw
25 extrusion of low solubility drugs and various polymers.

Another reason for forming an amorphous dispersion is that it may be
desired to use a particular process for forming a pharmaceutical composition,
such as a spray-coating or wet granulation process which results in the
formation of amorphous drug, in whole or in part, rather than pure crystalline
30 drug. Thus, amorphous dispersions may be formed of drugs which are not low-
solubility drugs.

However, regardless of whether the drug is poorly soluble, the inventors
have determined that for some drug and polymer dispersions, the drug is not
chemically stable in the dispersion. In particular, the inventors have observed
35 that for dispersions containing certain drugs and polymers, the drug degrades in

the dispersion over time, resulting in a loss of potency for the composition. The inventors have found this problem to arise especially for acid-sensitive drugs.

Drug degradation within the dispersion is a particular problem for low-solubility, acid-sensitive drugs, since the increase in aqueous concentration of the drug provided by the dispersion is offset by decreasing drug purity. In general, the greatest concentration-enhancement is often observed through the use of acidic dispersion polymers, especially acidic, cellulosic enteric polymers. However, the use of such acidic polymers within the dispersion is precluded due to the acid-sensitive nature of the drug.

Accordingly, there is still a need for pharmaceutical compositions of dispersions containing acid-sensitive drugs that are chemically stable over time. Likewise, there is also a continuing need to provide concentration-enhancement for low-solubility drugs.

BRIEF SUMMARY OF INVENTION

The present invention relates to, in a first aspect, pharmaceutical compositions comprising a solid amorphous dispersion of an acid-sensitive drug and a neutral dispersion polymer, wherein said composition provides improved chemical stability relative to a control acidic dispersion comprising an equivalent quantity of said drug and an acidic polymer.

In a preferred embodiment, the acid-sensitive drug has one or more functional groups selected from the group consisting of sulfonyl ureas, hydroxamic acids, hydroxy amides, carbamates, acetals, hydroxy ureas, esters, and amides.

In another preferred embodiment, the acid-sensitive drug when present in a control acidic dispersion and stored for a period of six months at 40°C and 75% relative humidity has a degree of degradation of at least 0.01%, preferably at least 0.1%.

In another preferred embodiment, the drug is quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3- fluorobenzyl-2(S),7-dihydroxy-7-methyloctyl]amide; quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide; quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide; (+)-N-{3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl}-N-

hydroxyurea; omeprazole; etoposide; famotidine; erythromycin; quinapril; lansoprazole; or progabide.

In another preferred embodiment, the drug in said composition has a relative degree of improvement in chemical stability of at least 1.25, preferably at least 3, more preferably at least 10. Preferably, drug in said composition has a relative degree of improvement in chemical stability of at least 1.25 when stored at 40°C and 75% relative humidity for a period of six months.

In another preferred embodiment, the dispersion polymer is ionizable, non-ionizable, cellulosic, or non-cellulosic. Preferred cellulosic dispersion polymers include hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and hydroxyethyl ethyl cellulose. Preferred non-cellulosic dispersion polymers include vinyl polymers and copolymers having one or more substituents comprising hydroxyl-containing repeat units, alkylacyloxy-containing repeat units, or cyclicamido-containing repeat units; polyvinyl alcohols that have at least a portion of their repeat units in the unhydrolyzed form; polyvinyl alcohol polyvinyl acetate copolymers; polyethylene glycol, polyethylene glycol polypropylene glycol copolymers, polyvinyl pyrrolidone; polyethylene polyvinyl alcohol copolymers, and polyoxyethylene-polyoxypropylene block copolymers. Preferably, the non-cellulosic dispersion polymer comprising a vinyl copolymer having: (1) hydroxyl-containing repeat units; and (2) hydrophobic repeat units.

In another preferred embodiment, the acid-sensitive drug is also a low-solubility drug and said dispersion polymer is concentration-enhancing. Preferably, the drug has a minimum solubility in aqueous solution in the absence of said dispersion polymer of less than 1 mg/mL at any pH of from about 1 to about 8, more preferably the minimum aqueous solubility is less than 0.01 mg/mL.

In another preferred embodiment, the drug has a dose-to-aqueous-solubility ratio of at least 10 mL.

In yet another preferred embodiment, the dispersion polymer is present in an amount sufficient to provide a maximum concentration of said acid-sensitive drug in a use environment that is at least 1.25-fold, preferably at least 2-fold, that provided by a second control composition comprising an equivalent quantity of said acid-sensitive drug and free from said dispersion polymer.

In another preferred embodiment, the dispersion polymer is present in a sufficient amount so that said dispersion provides in a use environment an area under the concentration versus time curve for any period of at least 90 minutes between the time of introduction into the use environment and about 270
5 minutes following introduction to the use environment that is at least 1.25-fold, preferably at least 2-fold, that provided by a second control composition comprising an equivalent quantity of said acid-sensitive drug and free from said dispersion polymer.

In another preferred embodiment, the dispersion polymer is present in a
10 sufficient amount so that said dispersion provides a relative bioavailability that is at least 1.25, preferably at least 2, relative to a second control composition comprising an equivalent quantity of said acid-sensitive drug and free from said polymer.

In yet another preferred embodiment, the drug is base-sensitive and said
15 dispersion polymer is non-ionizable.

In another preferred embodiment, the dispersion comprising a buffer. Preferred buffers include sodium acetate, ammonium acetate, sodium carbonate, sodium bicarbonate, disodium hydrogen phosphate and trisodium phosphate. Preferably, the buffer comprises at least 10 wt% of said dispersion.

20 In another preferred embodiment, the dispersion has a pH from about 6 to about 10. Preferably, the composition comprising a base and the dispersion has a pH of from about 6 to about 10.

In still another preferred embodiment, the acid-sensitive drug is a low-solubility drug and the composition comprising a second polymer, said
25 dispersion is free from at least a portion of said second polymer, and said second polymer is concentration-enhancing. Preferably, the second polymer has at least one hydrophobic portion and at least one hydrophilic portion. More preferably, the second polymer is a cellulosic, ionizable polymer, comprising hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl
30 cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, hydroxyethyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, hydroxyethyl methyl cellulose acetate succinate, hydroxyethyl methyl cellulose acetate phthalate, carboxyethyl cellulose, carboxymethyl cellulose, cellulose acetate phthalate, methyl cellulose
35 acetate phthalate, ethyl cellulose acetate phthalate, hydroxypropyl cellulose

acetate phthalate, hydroxypropyl methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, hydroxypropyl methyl cellulose acetate succinate phthalate, hydroxypropyl methyl cellulose succinate phthalate, cellulose propionate phthalate, hydroxypropyl cellulose butyrate phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate, ethyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate, hydroxypropyl methyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate succinate, cellulose propionate trimellitate, cellulose butyrate trimellitate, cellulose acetate terephthalate, cellulose acetate isophthalate, cellulose acetate pyridinedicarboxylate, salicylic acid cellulose acetate, hydroxypropyl salicylic acid cellulose acetate, ethylbenzoic acid cellulose acetate, hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl phthalic acid cellulose acetate, ethyl nicotinic acid cellulose acetate, and ethyl picolinic acid cellulose acetate. Other more preferable second polymers are non-ionizable cellulosic polymers comprising hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and hydroxyethyl ethyl cellulose. Still other more preferably second polymers are ionizable, non-cellulosic polymers comprising carboxylic acid functionalized polymethacrylates, carboxylic acid functionalized polyacrylates, amine-functionalized polyacrylates, amine-functionalized polymethacrylates, proteins, and carboxylic acid functionalized starches.

Yet other more preferable second polymers are non-ionizable, non-cellulosic polymers comprising vinyl polymers and copolymers having one or more substituents such as hydroxyl-containing repeat units, alkylacyloxy-containing repeat units, or cyclicamido-containing repeat units; polyvinyl alcohols that have at least a portion of their repeat units in the unhydrolyzed form; polyvinyl alcohol polyvinyl acetate copolymers; polyethylene glycol, polyethylene glycol polypropylene glycol copolymers, polyvinyl pyrrolidone polyethylene polyvinyl alcohol copolymers, and polyoxyethylene-polyoxypropylene block copolymers. Within these vinyl copolymers, preferably the second polymer has (1) hydroxyl-containing repeat units; and (2) hydrophobic repeat units.

In another preferred embodiment, the drug in said composition has a relative degree of improvement in chemical stability of at least 1.25, preferably at least 3.

In another preferred embodiment, the dispersion is mixed with said second polymer. Another preferred embodiment, the dispersion and said second polymer occupy separate regions of said composition.

5 In still another preferred embodiment, the second polymer is present in an amount sufficient to provide a maximum concentration of said acid-sensitive drug in a use environment that is at least 1.25-fold, preferably at least 2-fold, that of a control composition comprising an equivalent quantity of said dispersion and free from said second polymer.

10 In another preferred embodiment, the second polymer is present in a sufficient amount so that said composition provides in a use environment an area under the concentration versus time curve for any period of at least 90 minutes between the time of introduction into the use environment and about 270 minutes following introduction to the use environment that is at least 1.25-fold, preferably at least 2-fold, that of a control composition comprising an
15 equivalent quantity of said dispersion and free from said second polymer.

In another preferred embodiment, the second polymer is present in a sufficient amount so that said composition provides a relative bioavailability that is at least 1.25, preferably at least 2, relative to a control composition comprising an equivalent quantity of said dispersion and free from said second
20 polymer.

A second aspect of the present invention relates to methods for treating a condition in an animal comprising by administering to an animal in need of such treatment a therapeutic amount of the above-described composition.

25 A third aspect of the present invention relates to methods of administering a pharmaceutical composition comprising co-administering to a patient: (a) a solid amorphous dispersion comprising an acid-sensitive drug and a neutral polymer; and (b) a second polymer, wherein said dispersion is substantially free from said second polymer and said second polymer is concentration-enhancing.

30 In one preferred embodiment, the second polymer is cellulosic, such as hydroxypropyl methyl cellulose acetate succinate, cellulose acetate phthalate, hydroxypropyl methyl cellulose phthalate, methyl cellulose acetate phthalate, cellulose acetate trimellitate, hydroxypropyl cellulose acetate phthalate, cellulose acetate terephthalate and cellulose acetate isophthalate.

In another preferred embodiment, the dispersion is administered separately from said second polymer. Preferably, the dispersion and said second polymer are administered at about the same time.

5 A fourth aspect of the present invention relates to pharmaceutical compositions that a solid amorphous dispersion of a low-solubility drug and a neutral dispersion polymer, wherein said neutral dispersion polymer comprising a vinyl copolymer having hydrophilic hydroxyl-containing repeat units and hydrophobic repeat units.

10 In one preferred embodiment, the hydrophobic repeat units include ester-linked alkylate or arylate substituents. More preferably, the hydrophobic repeat unit is an alkylate such as acetate, propionate, and butyrate. More preferably, the hydrophobic repeat units comprise the acetylated form of the hydroxyl-containing repeat units. Preferably, the acetylated form of the hydroxyl-containing repeat units comprise 0.5 to 30% of the repeat units of the polymer.

15 In another preferred embodiment, the hydroxyl-containing repeat unit is vinyl alcohol.

In another preferred embodiment, the dispersion polymer is a vinyl alcohol/vinyl acetate copolymer. Preferably, from about 0.5% to about 30% of the repeat units of said polymer are vinyl acetate.

20 A fifth aspect of the present invention relates to pharmaceutical compositions that include a solid amorphous dispersion comprising a low-solubility drug, a neutral dispersion polymer, and an excipient such as a base or a buffer.

25 In a preferred embodiment, the neutral dispersion polymer is concentration enhancing. Preferred neutral dispersion polymers are present in an amount sufficient to provide a maximum concentration of said low-solubility drug in a use environment that is at least 1.25-fold, preferably at least 2-fold, that provided by a second control composition comprising an equivalent quantity of said low-solubility drug and free from a concentration-enhancing polymer.

30 In another preferred embodiment, the neutral dispersion polymer is present in a sufficient amount so that said composition, when introduced to a use environment, provides an area under the concentration-versus time curve for any period of at least 90 minutes between the time of introduction to the use environment and about 270 minutes following introduction to the use
35 environment that is at least 1.25-fold, preferably at least 2-fold, that of a second

control composition comprising an equivalent quantity of said low-solubility drug and free from a concentration-enhancing polymer.

In another preferred embodiment, the neutral dispersion polymer is present in a sufficient amount so that said composition provides a relative
5 bioavailability that is at least 1.25, preferably at least 2, relative to a second control composition comprising an equivalent quantity of said low-solubility drug and free from a concentration-enhancing polymer.

In another preferred embodiment, the drug in said composition has a relative degree of improvement in chemical stability of at least 1.25.

10 In yet another preferred embodiment, the composition provides improved chemical stability relative to a control composition, wherein said control comprising an equivalent quantity of a dispersion of said drug and said neutral dispersion polymer but free from said base and said buffer.

As described more fully below, the term "use environment" may refer to
15 the *in vivo* environment of the gastrointestinal (GI) tract of an animal, particularly a human, or the *in vitro* environment of a test solution, such as phosphate buffered saline (PBS) or model fasted duodenal (MFD) solution.

The composition may be dosed in a variety of dosage forms, including both immediate release and controlled release dosage forms, the latter
20 including both delayed and sustained release forms. The composition may include blends of polymers.

The various aspects of the present invention provide one or more of the following advantages. For those embodiments comprising an acid-sensitive drug, the inventors have recognized that a problem with forming dispersions of
25 acid-sensitive drugs is that for some dispersions, the drug does not remain chemically stable in the dispersion over time. The inventors have found that acid-sensitive drugs dispersed in an acidic polymer, such as hydroxypropyl methyl cellulose acetate succinate (which has carboxylic acid functional groups), have a tendency to chemically degrade over time. It is believed that
30 the presence of acidic ionic groups on the acidic polymer may either catalyze degradation of the drug or react directly with the drug. In any event, regardless of the particular degradation mechanism, the inventors have substantially reduced, if not eliminated the problem of drug degradation, by forming dispersions using neutral polymers, i.e. polymers that do not include acidic
35 functional groups. Thus, the present invention is able to realize the advantages

of forming dispersions of acid-sensitive drugs by improving the chemical stability of the acid-sensitive drug in the dispersion.

In addition, the invention in some embodiments further provides enhanced aqueous concentration and bioavailability for low-solubility drugs while at the same time minimizing the loss in potency and generation of impurities in the composition resulting from reaction or degradation of the drug when in the presence of an acidic species, such as an acidic dispersion polymer.

For those aspects comprising neutral vinyl copolymers of the present invention having hydroxyl-containing repeat units and hydrophobic repeat units, the compositions provide surprisingly effective concentration-enhancement. The polymers may be used with any low-solubility drug to improve the concentration of the drug in a use environment.

The foregoing and other objectives, features, and advantages of the invention will be more readily understood upon consideration of the following detailed description of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The pharmaceutical compositions of the present invention comprise solid amorphous dispersions of a drug and a neutral dispersion polymer. The present invention finds utility anytime it is desired to improve either the chemical stability of an acid-sensitive drug, to improve the concentration or bioavailability of a low-solubility drug, or both. In one embodiment of the invention, the dispersions of the present invention improve the chemical stability of acid-sensitive drugs. Thus, the dispersions may be used to prevent degradation of the drug due to interactions with dispersion polymers, acidic dispersion species, or other acidic excipients present in the composition. In another embodiment, the dispersions improve the concentration or bioavailability of a low-solubility drug. Suitable acid-sensitive drugs, low-solubility drugs, neutral polymers and methods for making the dispersions are discussed in more detail below.

ACID-SENSITIVE DRUGS

The term "drug" is conventional, denoting a compound having beneficial prophylactic and/or therapeutic properties when administered to an animal, especially humans. In one embodiment of the invention, the drug is an acid-sensitive drug, meaning that the drug either chemically reacts with or otherwise

degrades in the presence of acidic species. Acid-sensitive drugs often include functional groups which are reactive under acidic conditions, such as sulfonyl ureas, hydroxamic acids, hydroxy amides, carbamates, acetals, hydroxy ureas, esters, and amides. Drugs which include such functional groups may be prone to reactions such as hydrolysis, lactonization, or transesterification in the presence of acidic species.

Acid-sensitive drugs may be identified experimentally by determining whether the drug chemically reacts or degrades when dispersed in an acidic polymer. In particular, as used herein, the term "acid-sensitive drug" refers to a drug which, when dispersed in a "control acidic dispersion," degrades when stored under controlled aging conditions either for long storage times at ambient storage conditions or for short storage times under elevated temperature and relative humidity conditions.

The "control acidic dispersion" used to determine whether a drug is acid-sensitive is a dispersion of the drug and a pharmaceutically acceptable acidic polymer. A convenient pharmaceutically acceptable acidic polymer for use as the acidic dispersion polymer is the HF grade of hydroxypropyl methyl cellulose acetate succinate (HPMCAS), which is an acidic, or so-called enteric polymer having carboxylic acid functional groups, and a pK_a of about 5. The HPMCAS in the control acidic dispersion should have a minimum degree of substitution of succinate groups ($O(CO)CH_2CH_2(CO)OH$) of at least 4 wt% (or at least about 100 milliequivalents of carboxylic acid functional groups per mole of polymer). Alternatively, other acidic polymers which are at least as acidic as the HF grade of HPMCAS may be used, such as hydroxypropyl methyl cellulose acetate phthalate (HPMCP) or cellulose acetate phthalate (CAP). As with HPMCAS, such polymers should have at least about 100 milliequivalents of carboxylic acid functional groups per mole of polymer. The amount of acidic polymer present in the control acidic dispersion may vary, but should comprise at least 50 wt% or more of the control acidic dispersion. A drug is an acid-sensitive drug if it meets the drug degradation criteria described below in at least one control acidic dispersion comprised of 25 wt% drug and 75 wt% of the dispersion polymer HPMCAS, HPMCP or CAP.

In general, drug degradation may be measured using any conventional method for measuring the purity or potency of drug in a pharmaceutical composition. For example, the amount of active drug present in a dispersion

may be measured initially using high-performance liquid chromatography (HPLC) or any other analytical technique well known in the art. Alternatively, the amount of drug initially present may be calculated from the amount of drug present in the dispersion formulation. The potency of the dispersion may then
5 be measured after storage at controlled temperature and humidity conditions for an appropriate period of time. A decrease in potency indicates that a chemical reaction has occurred, leading to a decrease in the amount of active drug present in the dispersion, and is an indication of poor chemical stability.

An alternative method used to evaluate chemical stability is to analyze
10 the rate of increase in the amount of drug degradant(s) in the dispersion, which would indicate reaction of the drug to form the degradant(s). An HPLC or other analytical technique may be used to determine the concentration of drug degradant(s) in a dispersion. The amount of the degradant(s) is measured before and after storage under controlled aging conditions. The amount of
15 increase in the drug degradant(s) may be used to determine the amount of decrease in "percent drug purity." The "percent drug purity" is defined as 100 times the total amount of drug present divided by the total amount of drug initially present. Thus, a percent drug purity at a given time may be calculated by the formula

$$\begin{array}{l} 20 \qquad \qquad \qquad \text{total amt. of} \\ \qquad \qquad \qquad \text{drug present} \\ \text{wt\% drug purity} = \left(\frac{\text{total amt. of}}{\text{drug init. present}} \right) * 100 \\ 25 \qquad \qquad \qquad \text{drug init. present} \end{array}$$

When the drug purity is calculated from the total amount of impurities, "percent drug purity" may be calculated by assuming that the "total amount of drug initially present," given in wt%, is equal to 100 wt% minus the wt% of total initial impurities, and that "total amount of drug present" is equal to 100 wt%
30 minus the wt% of total impurities after storage, that is, at some later time. This method is equivalent to calculating "percent drug purity" by the formula

$$\begin{array}{l} \text{total amt. of impurities} \\ 35 \text{ wt\% drug purity} = [1 - \left(\frac{\text{total amt. of drug}}{\text{total amt. of drug}} \right)] * 100 \\ \qquad \qquad \qquad \text{init. present} \end{array}$$

The rate at which drug degradation occurs is generally dependent on the storage conditions. The drug, when formulated as a composition of the present invention, should be stable at ambient temperature and humidity conditions (e.g., relative humidities of 20% to 60%) for long periods of time, such as
5 months or years. However, to expedite testing, the storage conditions may employ elevated temperature and/or humidity to simulate longer storage times at ambient conditions. The storage time may vary from a few days to weeks or months, depending on the reactivity of the drug and the storage conditions.

A "degree of degradation" of drug following storage may be determined
10 by subtracting the final percent drug purity (either determined by measuring the decrease in drug present or an increase in the amount of drug degradants present) from the initial percent drug purity. For example, for a dispersion initially containing 100 mg drug, and no measurable impurities, the initial percent drug purity is 100 wt%. If, after storage, the amount of drug in the
15 dispersion decreases to 95 mg, the final percent drug purity would be 95 wt% and the "degree of degradation" is 5 wt% (100 wt% - 95 wt%). Alternatively, if 100 mg of drug substance were found to initially have 1 mg of impurities present, it would have an initial "percent drug purity" of 99 wt%. If, after storage, the total impurities present had increased to 6 wt%, the final percent
20 drug purity would be 94 wt% and the "degree of degradation" would be 5 wt% (99 wt% - 94 wt%).

Alternatively, "degree of degradation" can be determined by subtracting from the amount of one or more specific degradants initially present from the amount of that specific drug degradant present after storage. Such a measure
25 is useful where there are several drug degradants, of which only one (or a few) is of concern. The degree of degradation may be calculated on the basis of only those degradants that are of concern, rather than all of the degradant. For example, if a drug initially contained a specific degradant at a concentration of 1 wt% and after storage the concentration of that degradant was 6 wt%, the
30 degree of degradation would be 5 wt% (6 wt%-1 wt%).

Returning now to the determination of an acid-sensitive drug, as used herein an "acid-sensitive" drug is one in which the degree of degradation of the drug in a control acidic dispersion (as defined above) is at least 0.01 wt% in a six month period at 40°C and 75% relative humidity (RH), or alternatively, in
35 which the degree of degradation is at least 0.01 wt% in a one year period at

30°C at 60% RH, or as defined below with respect to degradation of drug in an acidic aqueous solution. These storage conditions are conventional and well-known in the art, and are the ICH recommendations for storage conditions to be used to evaluate the stability of a drug after storage for two years at ambient
5 temperature and humidity. A relative degree of improvement may become apparent within a shorter time, such as three to five days, and shorter storage times may be used for some very acid-sensitive drugs.

The need for the present invention will generally be greater when the drug's reactivity with or sensitivity to acidic species increases. Dispersions of
10 the present invention are preferred for acid-sensitive drugs having a degree of degradation that is greater than the minimum level when in the presence of an acidic dispersion polymer. Thus, the dispersions of the present invention are preferred for "very acid-sensitive drugs." Very acid-sensitive drugs are those which, when dispersed in a control acidic dispersion, have a degree of
15 degradation of at least 0.1 wt% when stored at 40°C/75% RH for six months. The dispersions are even more preferred for drugs which have a degree of degradation of at least 1 wt%, and are most preferred for drugs which have a degree of degradation of at least 5.0 wt% when stored at 40°C/75% RH for six months.

20 In addition, the acid-sensitive drug when present in the control acidic dispersion degrades at a rate that is greater than the rate at which pure drug in the amorphous form, or if the amorphous form is physically unstable, the crystalline form, degrades. A relative degree of degradation of the drug in a control acidic dispersion to the pure form of the drug may be determined by
25 taking the ratio of the degree of degradation of the drug in the control acidic dispersion and the degree of degradation of the pure form of the drug under the same storage conditions for the same storage time period. For example, where the degree of degradation of the drug in the control acidic dispersion is 5 wt% and the degree of degradation of the pure amorphous form of the drug is
30 0.1 wt%, the relative degree of degradation of the drug in the control acidic dispersion to the pure form of the drug is 50 (5 wt%/0.1 wt%). An acid-sensitive drug is one in which the relative degree of degradation of the drug in the control composition to the pure form of the drug is at least 2. The invention finds increasing utility as the relative degree of degradation increases above this
35 minimum level. Thus, the invention is preferred for drugs having a relative

degree of degradation of about 3 or more, and more preferred for drugs having a relative degree of degradation of about 5 or more.

Alternatively, another test to determine whether a drug is an acid sensitive drug as used herein is to administer the drug to an acidic aqueous solution and plot drug concentration versus time. The acidic solution should have a pH of from 1-4. Drugs which are acid sensitive are those for which the drug concentration decreases by at least 1% within 24 hours of administration of the drug to the acidic solution. If the drug concentration changes by 1% in the 6-24 hour time period, then the drug is "slightly acid-sensitive." If the drug concentration changes by 1% in the 1-6 hour time period, then the drug is "moderately acid-sensitive." If the drug concentration changes by 1% in less than 1 hour, then the drug is "highly acid-sensitive." The present invention finds increasing utility for drugs which are slightly acid-sensitive, moderately acid-sensitive and highly acid-sensitive.

Specific examples of acid-sensitive drugs deliverable by the invention are set forth below, by way of example only. Each named drug should be understood to include the neutral form of the drug, pharmaceutically acceptable salts, and prodrugs. Examples of acid-sensitive drugs include quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3-fluorobenzyl-2(S),7-dihydroxy-7-methyloctyl]amide; quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide; quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide; (+)-N-{3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl}-N-hydroxyurea; omeprazole; etoposide; famotidine; erythromycin; quinapril; lansoprazole; and progabide.

LOW-SOLUBILITY DRUGS

In another embodiment of the invention, the drug is a "low-solubility drug," meaning that the drug may be either "substantially water-insoluble," which means that the drug has a minimum aqueous solubility at physiologically relevant pH (e.g., pH 1-8) of less than 0.01 mg/mL, "sparingly water-soluble," that is, has an aqueous solubility up to about 1 to 2 mg/mL, or even low to moderate aqueous-solubility, having an aqueous-solubility from about 1 mg/mL to as high as about 20 to 40 mg/mL. In general, it may be said that the drug has a dose-to-aqueous solubility ratio greater than 10 mL, and more typically greater than 100 mL, where the drug solubility (mg/mL) is the minimum value observed

in any physiologically relevant aqueous solution (e.g., those with pH values between 1 and 8) including USP simulated gastric and intestinal buffers, and dose is in mg. The dose-to-aqueous-solubility-ratio may be determined by simply dividing the dose (in mg) by the aqueous solubility (in mg/mL).

5 Preferred classes of drugs include, but are not limited to, antihypertensives, antianxiety agents, anticlotting agents, anticonvulsants, blood glucose-lowering agents, decongestants, antihistamines, antitussives, antineoplastics, beta blockers, anti-inflammatories, antipsychotic agents, cognitive enhancers, anti-atherosclerotic agents, cholesterol-reducing agents, 10 antiobesity agents, autoimmune disorder agents, anti-impotence agents, antibacterial and antifungal agents, hypnotic agents, anti-Parkinsonism agents, anti-Alzheimer's disease agents, antibiotics, anti-depressants, and antiviral agents, glycogen phosphorylase inhibitors, and cholesterol ester transfer protein inhibitors.

15 Each named drug should be understood to include the neutral form of the drug, pharmaceutically acceptable salts, as well as prodrugs. Specific examples of antihypertensives include prazosin, nifedipine, amlodipine besylate, trimazosin and doxazosin; specific examples of a blood glucose-lowering agent are glipizide and chlorpropamide; a specific example of an anti-impotence agent 20 is sildenafil and sildenafil citrate; specific examples of antineoplastics include chlorambucil, lomustine and echinomycin; a specific example of an imidazole-type antineoplastic is tubulazole; a specific example of an anti-hypercholesterolemic is atorvastatin calcium; specific examples of anxiolytics include hydroxyzine hydrochloride and doxepin hydrochloride; specific 25 examples of anti-inflammatory agents include betamethasone, prednisolone, aspirin, piroxicam, valdecoxib, carprofen, celecoxib, flurbiprofen and (+)-N-(4-[3-(4-fluorophenoxy)phenoxy]-2-cyclopenten-1-yl)-N-hydroxyurea; a specific example of a barbiturate is phenobarbital; specific examples of antivirals include acyclovir, nelfinavir, and virazole; specific examples of vitamins/nutritional 30 agents include retinol and vitamin E; specific examples of beta blockers include timolol and nadolol; a specific example of an emetic is apomorphine; specific examples of a diuretic include chlorthalidone and spironolactone; a specific example of an anticoagulant is dicumarol; specific examples of cardiotonics include digoxin and digitoxin; specific examples of androgens include 17- 35 methyltestosterone and testosterone; a specific example of a mineral corticoid is

desoxycorticosterone; a specific example of a steroidal hypnotic/anesthetic is alfaxalone; specific examples of anabolic agents include fluoxymesterone and methanstenolone; specific examples of antidepressant agents include sulpiride, [3,6-dimethyl-2-(2,4,6-trimethyl-phenoxy)-pyridin-4-yl]-(1-ethylpropyl)-amine, 5 3,5-dimethyl-4-(3'-pentoxy)-2-(2',4',6'-trimethylphenoxy)pyridine, pyroxidine, fluoxetine, paroxetine, venlafaxine and sertraline; specific examples of antibiotics include carbenicillin indanyl sodium, bacampicillin hydrochloride, troleandomycin, doxycycline hyclate, ampicillin and penicillin G; specific examples of anti-infectives include benzalkonium chloride and chlorhexidine; 10 specific examples of coronary vasodilators include nitroglycerin and miflozine; a specific example of a hypnotic is etomidate; specific examples of carbonic anhydrase inhibitors include acetazolamide and chlorzotamide; specific examples of antifungals include econazole, terconazole, fluconazole, voriconazole, and griseofulvin; a specific example of an antiprotozoal is 15 metronidazole; specific examples of anthelmintic agents include thiabendazole and oxfendazole and morantel; specific examples of antihistamines include astemizole, levocabastine, cetirizine, decarboethoxy loratadine and cinnarizine; specific examples of antipsychotics include ziprasidone, olanzapine, thiothixene hydrochloride, fluspirilene, risperidone and penfluridol; specific examples of 20 gastrointestinal agents include loperamide and cisapride; specific examples of serotonin antagonists include ketanserin and mianserin; a specific example of an anesthetic is lidocaine; a specific example of a hypoglycemic agent is acetohexamide; a specific example of an anti-emetic is dimenhydrinate; a specific example of an antibacterial is cotrimoxazole; a specific example of a 25 dopaminergic agent is L-DOPA; specific examples of anti-Alzheimer's Disease agents are THA and donepezil; a specific example of an anti-ulcer agent/H2 antagonist is famotidine; specific examples of sedative/hypnotic agents include chlordiazepoxide and triazolam; a specific example of a vasodilator is alprostadil; a specific example of a platelet inhibitor is prostacyclin; specific 30 examples of ACE inhibitor/antihypertensive agents include enalaprilic acid and lisinopril; specific examples of tetracycline antibiotics include oxytetracycline and minocycline; specific examples of macrolide antibiotics include erythromycin, clarithromycin, and spiramycin; a specific example of an azalide antibiotic is azithromycin; specific examples of glycogen phosphorylase 35 inhibitors include [R-(R'S')]-5-chloro-N-[2-hydroxy-3-{methoxymethylamino}-3-

oxo-1-(phenylmethyl)propyl-1H-indole-2-carboxamide and 5-chloro-1H-indole-2-carboxylic acid [(1S)-benzyl-(2R)-hydroxy-3-((3R,4S)-dihydroxy-pyrrolidin-1-yl)-3-oxypropyl]amide; and specific examples of cholesterol ester transfer protein (CETP) inhibitors include [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester, [2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester, [2R, 4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester.

The invention is not limited by any particular structure or group of CETP inhibitors. Rather, the invention has general applicability to CETP inhibitors as a class, the class tending to be composed of compounds having low solubility. Compounds which may be the subject of the invention may be found in a number of patents and published applications, including DE 19741400 A1; DE 19741399 A1; WO 9914215 A1; WO 9914174; DE 19709125 A1; DE 19704244 A1; DE 19704243 A1; EP 818448 A1; WO 9804528 A2; DE 19627431 A1; DE 19627430 A1; DE 19627419 A1; EP 796846 A1; DE 19832159; DE 818197; DE 19741051; WO 9941237 A1; WO 9914204 A1; WO 9835937 A1; JP 11049743; WO 200018721; WO 200018723; WO 200018724; WO 200017164; WO 200017165; WO 200017166; EP 992496; and EP 987251, all of which are hereby incorporated by reference in their entireties for all purposes.

The invention is useful for CETP inhibitors that have sufficiently low aqueous solubility, low bioavailability or slow rate of absorption such that it is desirable to increase their concentration in an aqueous environment of use. Therefore, anytime one finds it desirable to raise the aqueous concentration of the CETP inhibitor in a use environment, the invention will find utility. The CETP inhibitor is "substantially water-insoluble" which means that the CETP inhibitor has a minimum aqueous solubility of less than about 0.01 mg/mL (or 10 μ g/mL) at any physiologically relevant pH (e.g., pH 1-8) and at about 22°C. (Unless otherwise specified, reference to aqueous solubility herein and in the claims is determined at about 22°C.) Compositions of the present invention find greater utility as the solubility of the CETP inhibitors decreases, and thus are preferred for CETP inhibitors with solubilities less than about 2 μ g/mL, and even more preferred for CETP inhibitors with solubilities less than about 0.5 μ g/mL.

Many CETP inhibitors have even lower solubilities (some even less than 0.1 $\mu\text{g/mL}$), and require dramatic concentration enhancement to be sufficiently bioavailable upon oral dosing for effective plasma concentrations to be reached at practical doses.

5 In general, it may be said that the CETP inhibitor has a dose-to-aqueous solubility ratio greater than about 100 mL, where the solubility (mg/mL) is the minimum value observed in any physiologically relevant aqueous solution (e.g., those with pH values from 1 to 8) including USP simulated gastric and intestinal buffers, and dose is in mg. Compositions of the present invention, as
10 mentioned above, find greater utility as the solubility of the CETP inhibitor decreases and the dose increases. Thus, the compositions are preferred as the dose-to-solubility ratio increases, and thus are preferred for dose-to-solubility ratios greater than 1000 mL, and more preferred for dose-to-solubility ratios greater than about 5000 mL. The dose-to-solubility ratio may be determined by
15 dividing the dose (in mg) by the aqueous solubility (in mg/mL).

 Oral delivery of many CETP inhibitors is particularly difficult because their aqueous solubility is usually extremely low, typically being less than 2 $\mu\text{g/mL}$, often being less than 0.1 $\mu\text{g/mL}$. Such low solubilities are a direct consequence of the particular structural characteristics of species that bind to
20 CETP and thus act as CETP inhibitors. This low solubility is primarily due to the hydrophobic nature of CETP inhibitors. Clog P, defined as the base 10 logarithm of the ratio of the drug solubility in octanol to the drug solubility in water, is a widely accepted measure of hydrophobicity. In general, Clog P values for CETP inhibitors are greater than 4 and are often greater than 5 to 7.
25 Thus, the hydrophobic and insoluble nature of CETP inhibitors as a class pose a particular challenge for oral delivery. Achieving therapeutic drug levels in the blood by oral dosing of practical quantities of drug generally requires a large enhancement in drug concentrations in the gastrointestinal fluid and a resulting large enhancement in bioavailability. Such enhancements in drug concentration
30 in gastrointestinal fluid typically need to be at least about 10-fold and often at least about 50-fold or even at least about 200-fold to achieve desired blood levels. Surprisingly, the dispersions of the present invention have proven to have the required large enhancements in drug concentration and bioavailability.

 In contrast to conventional wisdom, the relative degree of enhancement
35 in aqueous concentration and bioavailability generally improves for CETP

inhibitors as solubility decreases and hydrophobicity increases. In fact, the inventors have recognized a subclass of these CETP inhibitors that are essentially aqueous insoluble, highly hydrophobic, and are characterized by a set of physical properties. This subclass exhibits dramatic enhancements in aqueous concentration and bioavailability when formulated using the compositions of the present invention.

The first property of this subclass of essentially insoluble, hydrophobic CETP inhibitors is extremely low aqueous solubility. By extremely low aqueous solubility is meant that the minimum aqueous solubility at physiologically relevant pH (pH of 1 to 8) is less than about 10 $\mu\text{g/ml}$ and preferably less than about 1 $\mu\text{g/ml}$.

A second property is a very high dose-to-solubility ratio. Extremely low solubility often leads to poor or slow absorption of the drug from the fluid of the gastrointestinal tract, when the drug is dosed orally in a conventional manner. For extremely low solubility drugs, poor absorption generally becomes progressively more difficult as the dose (mass of drug given orally) increases. Thus, a second property of this subclass of essentially insoluble, hydrophobic CETP inhibitors is a very high dose (in mg) to solubility (in mg/ml) ratio (ml). By "very high dose-to-solubility ratio" is meant that the dose-to-solubility ratio has a value of at least 1000 ml, and preferably at least 5,000 ml, and more preferably at least 10,000 ml.


A third property of this subclass of essentially insoluble, hydrophobic CETP inhibitors is that they are extremely hydrophobic. By extremely hydrophobic is meant that the Clog P value of the drug, has a value of at least 4.0, preferably a value of at least 5.0, and more preferably a value of at least 5.5.

A fourth property of this subclass of essentially insoluble CETP inhibitors is that they have a low melting point. Generally, drugs of this subclass will have a melting point of about 150°C or less, and preferably about 140°C or less.

Primarily, as a consequence of some or all of these four properties, CETP inhibitors of this subclass typically have very low absolute bioavailabilities. Specifically, the absolute bioavailability of drugs in this subclass when dosed orally in their undispersed state is less than about 10% and more often less than about 5%.

Turning now to the chemical structures of specific CETP inhibitors, one class of CETP inhibitors that finds utility with the present invention consists of oxy substituted 4-carboxyamino-2-methyl-1,2,3,4-tetrahydroquinolines having the Formula I

5



Formula I

and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

wherein R_{1-1} is hydrogen, Y_i , W_i-X_i , W_i-Y_i ;

wherein W_i is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_i is $-O-Y_i$, $-S-Y_i$, $-N(H)-Y_i$ or $-N-(Y_i)_2$;

wherein Y_i for each occurrence is independently Z_i or a fully saturated, partially unsaturated or fully unsaturated one to ten membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with Z_i ;

wherein Z_i is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six

membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said Z_i substituent is optionally mono-, di- or tri-substituted independently with halo, (C₂-C₆)alkenyl, (C₁-C₆) alkyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxyl, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxyl, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines; R_{1-3} is hydrogen or Q_i ;

wherein Q_i is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_i ;

wherein V_i is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_i substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carbamoyl, mono-N- or di-N,N-(C₁-C₆) alkylcarbamoyl, carboxyl, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl or (C₂-C₆)alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxyl, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl or

(C₂-C₆)alkenyl substituents are also optionally substituted with from one to nine fluorines;

R₁₋₄ is Q₁₋₁ or V₁₋₁

5 wherein Q₁₋₁ is a fully saturated, partially unsaturated or fully unsaturated
one to six membered straight or branched carbon chain wherein the carbons,
other than the connecting carbon, may optionally be replaced with one
heteroatom selected from oxygen, sulfur and nitrogen and said carbon is
optionally mono-, di- or tri-substituted independently with halo, said carbon is
optionally mono-substituted with hydroxy, said carbon is optionally mono-
10 substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo,
said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon
chain is optionally mono-substituted with

V₁₋₁;

15 wherein V₁₋₁ is a partially saturated, fully saturated or fully unsaturated
three to six membered ring optionally having one to two heteroatoms selected
independently from oxygen, sulfur and nitrogen;

20 wherein said V₁₋₁ substituent is optionally mono-, di-, tri-, or tetra-
substituted independently with halo, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, amino, nitro,
cyano, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein
said (C₁-C₆)alkyl substituent is optionally mono-substituted with oxo, said (C₁-
C₆)alkyl substituent is also optionally substituted with from one to nine fluorines;

25 wherein either R₁₋₃ must contain V₁ or R₁₋₄ must contain V₁₋₁; and R₁₋₅, R₁₋₆,
R₁₋₇ and R₁₋₈ are each independently hydrogen, hydroxy or oxy wherein said
oxy is substituted with T₁ or a partially saturated, fully saturated or fully
unsaturated one to twelve membered straight or branched carbon chain wherein
the carbons, other than the connecting carbon, may optionally be replaced with
one or two heteroatoms selected independently from oxygen, sulfur and
nitrogen and said carbon is optionally mono-, di- or tri-substituted independently
with halo, said carbon is optionally mono-substituted with hydroxy, said carbon
30 is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-
substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo,
and said carbon chain is optionally mono-substituted with T₁;

35 wherein T₁ is a partially saturated, fully saturated or fully unsaturated
three to eight membered ring optionally having one to four heteroatoms selected
independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of

two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said T₁ substituent is optionally mono-, di- or tri-substituted independently with halo, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines.

Compounds of Formula I and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,140,342, United States Patent No. 6,362,198, and European Patent publication 987251, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula I:

- [2R,4S] 4-[(3,5-dichloro-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- [2R,4S] 4-[(3,5-dinitro-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- [2R,4S] 4-[(2,6-dichloro-pyridin-4-ylmethyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-methoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-methoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
- [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
- [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-ethoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid 2,2,2-trifluoroethylester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-dimethoxy-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid tert-butyl ester;


[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-6-trifluoromethoxy-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester,

[2R,4S] (3,5-bis-trifluoromethyl-benzyl)-(1-butyryl-6,7-dimethoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-carbamic acid methyl ester;

[2R,4S] (3,5-bis-trifluoromethyl-benzyl)-(1-butyl-6,7-dimethoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl)-carbamic acid methyl ester; and

[2R,4S] (3,5-bis-trifluoromethyl-benzyl)-[1-(2-ethyl-butyl)-6,7-dimethoxy-2-methyl-1,2,3,4-tetrahydro-quinolin-4-yl]-carbamic acid methyl ester, hydrochloride.

Another class of CETP inhibitors that finds utility with the present invention consists of 4-carboxyamino-2-methyl-1,2,3,4,-tetrahydroquinolines, having the Formula II



Formula II

and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

wherein R_{II-1} is hydrogen, Y_{II} , $W_{II}-X_{II}$, $W_{II}-Y_{II}$;

wherein W_{II} is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_{II} is $-O-Y_{II}$, $-S-Y_{II}$, $-N(H)-Y_{II}$ or $-N-(Y_{II})_2$;

wherein Y_{ii} for each occurrence is independently Z_{ii} or a fully saturated, partially unsaturated or fully unsaturated one to ten membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with Z_{ii} ;

Z_{ii} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said Z_{ii} substituent is optionally mono-, di- or tri-substituted independently with halo, (C_2-C_6) alkenyl, (C_1-C_6) alkyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl is also optionally substituted with from one to nine fluorines;

R_{ii-3} is hydrogen or Q_{ii} ;

wherein Q_{ii} is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{ii} ;

wherein V_{II} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_{II} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxamoyl, mono-N- or di-N,N- (C_1-C_6) alkylcarboxamoyl, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino or said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituents are optionally substituted with from one to nine fluorines;

R_{II-4} is Q_{II-1} or V_{II-1}

wherein Q_{II-1} a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{II-1} ;

wherein V_{II-1} is a partially saturated, fully saturated or fully unsaturated three to six membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen;

wherein said V_{II-1} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, amino, nitro, cyano, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-substituted with oxo, said (C_1-C_6) alkyl substituent is optionally substituted with from one to nine fluorines;

wherein either R_{II-3} must contain V_{II} or R_{II-4} must contain V_{II-1} ; and

R_{II-5} , R_{II-6} , R_{II-7} and R_{II-8} are each independently hydrogen, a bond, nitro or halo wherein said bond is substituted with T_{II} or a partially saturated, fully saturated or fully unsaturated (C_1 - C_{12}) straight or branched carbon chain wherein carbon may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon is optionally mono-substituted with T_{II} ;

wherein T_{II} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said T_{II} substituent is optionally mono-, di- or tri-substituted independently with halo, (C_1 - C_6)alkyl, (C_2 - C_6)alkenyl, hydroxy, (C_1 - C_6)alkoxy, (C_1 - C_4)alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1 - C_6)alkyloxycarbonyl, mono-N- or di-N,N-(C_1 - C_6)alkylamino wherein said (C_1 - C_6)alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1 - C_6)alkoxy, (C_1 - C_4)alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1 - C_6)alkyloxycarbonyl, mono-N- or di-N,N-(C_1 - C_6)alkylamino, said (C_1 - C_6)alkyl substituent is also optionally substituted with from one to nine fluorines; provided that at least one of substituents R_{II-5} , R_{II-6} , R_{II-7} and R_{II-8} is not hydrogen and is not linked to the quinoline moiety through oxy.

Compounds of Formula II and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,147,090, United States Patent Application No. 09/671,400 filed September 27, 2000, and PCT Publication No. WO00/17166, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula II:

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-7-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-chloro-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-chloro-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

5 [2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2,6,7-trimethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6,7-diethyl-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;


10 [2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-ethyl-2-methyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester; and

15 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester.

Another class of CETP inhibitors that finds utility with the present invention consists of annulated 4-carboxyamino-2-methyl-1,2,3,4,-tetrahydroquinolines, having the Formula III

20



Formula III

25 and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

wherein R_{III-1} is hydrogen, Y_{III} , $W_{III}-X_{III}$, $W_{III}-Y_{III}$;

wherein W_{III} is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_{III} is $-O-Y_{III}$, $-S-Y_{III}$, $-N(H)-Y_{III}$ or $-N-(Y_{III})_2$;

Y_{III} for each occurrence is independently Z_{III} or a fully saturated, partially unsaturated or fully unsaturated one to ten membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with Z_{III};

wherein Z_{III} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said Z_{III} substituent is optionally mono-, di- or tri-substituted independently with halo, (C₂-C₆)alkenyl, (C₁-C₆) alkyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl optionally substituted with from one to nine fluorines;

R_{III-3} is hydrogen or Q_{III};

wherein Q_{III} is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{III};

wherein V_{III} is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

5 wherein said V_{III} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxamoyl, mono-N- or di-N,N- (C_1-C_6) alkylcarboxamoyl, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino or said (C_1-C_6) alkyl or (C_2-C_6) alkenyl are optionally substituted with from one to nine fluorines;
15 R_{III-4} is Q_{III-1} or V_{III-1} ;

wherein Q_{III-1} a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with
25 V_{III-1} ;

wherein V_{III-1} is a partially saturated, fully saturated or fully unsaturated three to six membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen;

30 wherein said V_{III-1} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, amino, nitro, cyano, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-substituted with oxo, said (C_1-C_6) alkyl substituent optionally having from one to nine fluorines;

35 wherein either R_{III-3} must contain V_{III} or R_{III-4} must contain V_{III-1} ; and

R_{III-5} and R_{III-6}, or R_{III-6} and R_{III-7}, and/or R_{III-7} and R_{III-8} are taken together and form at least one four to eight membered ring that is partially saturated or fully unsaturated optionally having one to three heteroatoms independently selected from nitrogen, sulfur and oxygen;

5 wherein said ring or rings formed by R_{III-5} and R_{III-6}, or R_{III-6} and R_{III-7}, and/or R_{III-7} and R_{III-8} are optionally mono-, di- or tri-substituted independently with halo, (C₁-C₆)alkyl, (C₁-C₄)alkylsulfonyl, (C₂-C₆)alkenyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with
10 hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent optionally having from one to nine fluorines;

 provided that the R_{III-5}, R_{III-6}, R_{III-7} and/or R_{III-8}, as the case may be, that
15 do not form at least one ring are each independently hydrogen, halo, (C₁-C₆)alkoxy or (C₁-C₆)alkyl, said (C₁-C₆)alkyl optionally having from one to nine fluorines.

 Compounds of Formula III and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,147,089, United
20 States Patent No. 6,310,075, and European Patent Application No. 99307240.4 filed September 14, 1999, all of which are incorporated herein by reference in their entireties for all purposes.

 In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula III:

25 [2R, 4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-2,3,4,6,7,8-hexahydro-cyclopenta[g]quinoline-1-carboxylic acid ethyl ester;

 [6R, 8S] 8-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-methyl-3,6,7,8-tetrahydro-1H-2-thia-5-aza-cyclopenta[b]naphthalene-5-
30 carboxylic acid ethyl ester;

 [6R, 8S] 8-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-methyl-3,6,7,8-tetrahydro-2H-furo[2,3-g]quinoline-5-carboxylic acid ethyl ester;


 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-3,4,6,8-tetrahydro-2H-furo[3,4-g]quinoline-1-carboxylic acid ethyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methyl-3,4,6,7,8,9-hexahydro-2H-benzo[g]quinoline-1-carboxylic acid propyl ester;

5 [7R,9S] 9-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-7-methyl-1,2,3,7,8,9-hexahydro-6-aza-cyclopenta[a]naphthalene-6-carboxylic acid ethyl ester; and

[6S,8R] 6-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-8-methyl-1,2,3,6,7,8-hexahydro-9-aza-cyclopenta[a]naphthalene-9-carboxylic acid ethyl ester.

10 Another class of CETP inhibitors that finds utility with the present invention consists of 4-carboxyamino-2-substituted-1,2,3,4,-tetrahydroquinolines, having the Formula IV



15

Formula IV

and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

wherein R_{IV-1} is hydrogen, Y_{IV} , $W_{IV}-X_{IV}$ or $W_{IV}-Y_{IV}$;

20 wherein W_{IV} is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_{IV} is $-O-Y_{IV}$, $-S-Y_{IV}$, $-N(H)-Y_{IV}$ or $-N-(Y_{IV})_2$;

wherein Y_{IV} for each occurrence is independently Z_{IV} or a fully saturated, partially unsaturated or fully unsaturated one to ten membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is

25

optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with Z_{IV} ;

5 wherein Z_{IV} is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

10 wherein said Z_{IV} substituent is optionally mono-, di- or tri-substituted independently with halo, (C_2-C_6) alkenyl, (C_1-C_6) alkyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl substituent is also optionally substituted with from one to nine fluorines;
15 R_{IV-2} is a partially saturated, fully saturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with oxo, said carbon is optionally mono-substituted with hydroxy, said sulfur is optionally mono- or di-substituted with
20 oxo, said nitrogen is optionally mono- or di-substituted with oxo; or said R_{IV-2} is a partially saturated, fully saturated or fully unsaturated three to seven membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen, wherein said R_{IV-2} ring is optionally attached through (C_1-C_4) alkyl;

30 wherein said R_{IV-2} ring is optionally mono-, di- or tri-substituted independently with halo, (C_2-C_6) alkenyl, (C_1-C_6) alkyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, oxo or (C_1-C_6) alkyloxycarbonyl;
35

with the proviso that R_{IV-2} is not methyl;

R_{IV-3} is hydrogen or Q_{IV} ;

wherein Q_{IV} is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons
5 other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo,
10 said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{IV} ;

wherein V_{IV} is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of
15 two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_{IV} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxamoyl, mono-N- or
20 di-N,N- (C_1-C_6) alkylcarboxamoyl, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl or
25 (C_2-C_6) alkenyl substituents are also optionally substituted with from one to nine fluorines;

R_{IV-4} is Q_{IV-1} or V_{IV-1} ;

wherein Q_{IV-1} a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons,
30 other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo,
35

said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with

V_{IV-1} ;

5 wherein V_{IV-1} is a partially saturated, fully saturated or fully unsaturated three to six membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen;

wherein said V_{IV-1} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, amino, nitro, cyano, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein
10 said (C_1-C_6) alkyl substituent is optionally mono-substituted with oxo, said (C_1-C_6) alkyl substituent is also optionally substituted with from one to nine fluorines;

wherein either R_{IV-3} must contain V_{IV} or R_{IV-4} must contain V_{IV-1} ;
 R_{IV-5} , R_{IV-6} , R_{IV-7} and R_{IV-8} are each independently hydrogen, a bond, nitro or halo wherein said bond is substituted with T_{IV} or a partially saturated, fully
15 saturated or fully unsaturated (C_1-C_{12}) straight or branched carbon chain wherein carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally
20 mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon is optionally mono-substituted with T_{IV} ;

wherein T_{IV} is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected
25 independently from oxygen, sulfur and nitrogen, or, a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said T_{IV} substituent is optionally mono-, di- or tri-substituted
30 independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, $(C_1-$

C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines; and

wherein R_{IV-5} and R_{IV-6}, or R_{IV-6} and R_{IV-7}, and/or R_{IV-7} and R_{IV-8} may also be taken together and can form at least one four to eight membered ring that is partially saturated or fully unsaturated optionally having one to three heteroatoms independently selected from nitrogen, sulfur and oxygen;

wherein said ring or rings formed by R_{IV-5} and R_{IV-6}, or R_{IV-6} and R_{IV-7}, and/or R_{IV-7} and R_{IV-8} are optionally mono-, di- or tri-substituted independently with halo, (C₁-C₆)alkyl, (C₁-C₄)alkylsulfonyl, (C₂-C₆)alkenyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines; with the proviso that when R_{IV-2} is carboxyl or (C₁-C₄)alkylcarboxyl, then R_{IV-1} is not hydrogen.

Compounds of Formula IV and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,197,786, United States Application Serial No. 09/685,3000 filed 10/10/00, and PCT Publication No. WO 00/17164, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula IV:

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-isopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-6-chloro-2-cyclopropyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 2-cyclopropyl-4-[(3,5-dichloro-benzyl)-methoxycarbonyl-amino]-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid tert-butyl ester;

[2R,4R] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

5 [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclobutyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

10 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-methoxymethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

15 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid 2-hydroxyethyl ester;

20 [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;


[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

25 [2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester; and

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester.

30 Another class of CETP inhibitors that finds utility with the present invention consists of 4-amino substituted-2-substituted-1,2,3,4,-tetrahydroquinolines, having the Formula V

38



Formula V

and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said
5 compounds;

wherein R_{V-1} is Y_V , W_V-X_V or W_V-Y_V ;

wherein W_V is a carbonyl, thiocarbonyl, sulfinyl or sulfonyl;

X_V is $-O-Y_V$, $-S-Y_V$, $-N(H)-Y_V$ or $-N-(Y_V)_2$;

wherein Y_V for each occurrence is independently Z_V or a fully saturated,
10 partially unsaturated or fully unsaturated one to ten membered straight or
branched carbon chain wherein the carbons, other than the connecting carbon,
may optionally be replaced with one or two heteroatoms selected independently
from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-
substituted independently with halo, said carbon is optionally mono-substituted
15 with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is
optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-,
or di-substituted with oxo, and said carbon chain is optionally mono-substituted
with Z_V ;

wherein Z_V is a partially saturated, fully saturated or fully unsaturated
20 three to eight membered ring optionally having one to four heteroatoms selected
independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of
two fused partially saturated, fully saturated or fully unsaturated three to six
membered rings, taken independently, optionally having one to four
heteroatoms selected independently from nitrogen, sulfur and oxygen;

25 wherein said Z_V substituent is optionally mono-, di- or tri-substituted
independently with halo, (C_2-C_6) alkenyl, (C_1-C_6) alkyl, hydroxy, (C_1-C_6) alkoxy,
 (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl,
mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is
optionally mono-, di- or tri-substituted independently with halo, hydroxy, $(C_1-$

C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent is also optionally substituted with from one to nine fluorines;

5 R_{V,2} is a partially saturated, fully saturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons, other than the connecting carbon, may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with oxo, said carbon is optionally mono-
10 substituted with hydroxy, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo; or said R_{V,2} is a partially saturated, fully saturated or fully unsaturated three to seven membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen, wherein said R_{V,2} ring is optionally attached through
15 (C₁-C₄)alkyl;

wherein said R_{V,2} ring is optionally mono-, di- or tri-substituted independently with halo, (C₂-C₆)alkenyl, (C₁-C₆) alkyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is
20 optionally mono-, di- or tri-substituted independently with halo, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, oxo or (C₁-C₆)alkyloxycarbonyl;
R_{V,3} is hydrogen or Q_V;

wherein Q_V is a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons,
25 other than the connecting carbon, may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo,
30 said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_V;

wherein V_V is a partially saturated, fully saturated or fully unsaturated three to eight membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of
35 two fused partially saturated, fully saturated or fully unsaturated three to six

membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_v substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxamoyl, mono-N- or di-N,N- (C_1-C_6) alkylcarboxamoyl, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl or (C_2-C_6) alkenyl substituents are also optionally substituted with from one to nine fluorines;

R_{V-4} is cyano, formyl, $W_{V-1}Q_{V-1}$, $W_{V-1}V_{V-1}$, (C_1-C_4) alkylene V_{V-1} or V_{V-2} ;

wherein W_{V-1} is carbonyl, thiocarbonyl, SO or SO_2 ,

wherein Q_{V-1} a fully saturated, partially unsaturated or fully unsaturated one to six membered straight or branched carbon chain wherein the carbons may optionally be replaced with one heteroatom selected from oxygen, sulfur and nitrogen and said carbon is optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono-, or di-substituted with oxo, and said carbon chain is optionally mono-substituted with V_{V-1} ;

wherein V_{V-1} is a partially saturated, fully saturated or fully unsaturated three to six membered ring optionally having one to two heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said V_{V-1} substituent is optionally mono-, di-, tri-, or tetra-substituted independently with halo, (C_1-C_6) alkyl, (C_1-C_6) alkoxy, hydroxy, oxo, amino, nitro, cyano, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-substituted with oxo, said (C_1-C_6) alkyl substituent is also optionally substituted with from one to nine fluorines;

wherein $V_{V.2}$ is a partially saturated, fully saturated or fully unsaturated five to seven membered ring containing one to four heteroatoms selected independently from oxygen, sulfur and nitrogen;

wherein said $V_{V.2}$ substituent is optionally mono-, di- or tri-substituted independently with halo, (C_1-C_2) alkyl, (C_1-C_2) alkoxy, hydroxy, or oxo wherein said (C_1-C_2) alkyl optionally has from one to five fluorines; and

wherein $R_{V.4}$ does not include oxycarbonyl linked directly to the C^4 nitrogen;

wherein either $R_{V.3}$ must contain V_V or $R_{V.4}$ must contain $V_{V.1}$;

$R_{V.5}$, $R_{V.6}$, $R_{V.7}$ and $R_{V.8}$ are independently hydrogen, a bond, nitro or halo wherein said bond is substituted with T_V or a partially saturated, fully saturated or fully unsaturated (C_1-C_{12}) straight or branched carbon chain wherein carbon may optionally be replaced with one or two heteroatoms selected independently from oxygen, sulfur and nitrogen, wherein said carbon atoms are optionally mono-, di- or tri-substituted independently with halo, said carbon is optionally mono-substituted with hydroxy, said carbon is optionally mono-substituted with oxo, said sulfur is optionally mono- or di-substituted with oxo, said nitrogen is optionally mono- or di-substituted with oxo, and said carbon chain is optionally mono-substituted with T_V ;

wherein T_V is a partially saturated, fully saturated or fully unsaturated three to twelve membered ring optionally having one to four heteroatoms selected independently from oxygen, sulfur and nitrogen, or a bicyclic ring consisting of two fused partially saturated, fully saturated or fully unsaturated three to six membered rings, taken independently, optionally having one to four heteroatoms selected independently from nitrogen, sulfur and oxygen;

wherein said T_V substituent is optionally mono-, di- or tri-substituted independently with halo, (C_1-C_6) alkyl, (C_2-C_6) alkenyl, hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino wherein said (C_1-C_6) alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C_1-C_6) alkoxy, (C_1-C_4) alkylthio, amino, nitro, cyano, oxo, carboxy, (C_1-C_6) alkyloxycarbonyl, mono-N- or di-N,N- (C_1-C_6) alkylamino, said (C_1-C_6) alkyl substituent also optionally has from one to nine fluorines;

wherein $R_{V.5}$ and $R_{V.6}$, or $R_{V.6}$ and $R_{V.7}$, and/or $R_{V.7}$ and $R_{V.8}$ may also be taken together and can form at least one ring that is a partially saturated or fully

unsaturated four to eight membered ring optionally having one to three heteroatoms independently selected from nitrogen, sulfur and oxygen;

wherein said rings formed by $R_{V.5}$ and $R_{V.6}$, or $R_{V.6}$ and $R_{V.7}$, and/or $R_{V.7}$ and $R_{V.8}$ are optionally mono-, di- or tri-substituted independently with halo, (C₁-C₆)alkyl, (C₁-C₄)alkylsulfonyl, (C₂-C₆)alkenyl, hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino wherein said (C₁-C₆)alkyl substituent is optionally mono-, di- or tri-substituted independently with hydroxy, (C₁-C₆)alkoxy, (C₁-C₄)alkylthio, amino, nitro, cyano, oxo, carboxy, (C₁-C₆)alkyloxycarbonyl, mono-N- or di-N,N-(C₁-C₆)alkylamino, said (C₁-C₆)alkyl substituent also optionally has from one to nine fluorines.

Compounds of Formula V and their methods of manufacture are disclosed in commonly assigned United States Patent No. 6,140,343, United States Patent Application Serial No. 09/671,221 filed September 27, 2000, and PCT Publication No. WO 00/17165, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula V:

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester;
[2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid tert-butyl ester;
[2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
[2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
[2S,4S] 4-[1-(3,5-bis-trifluoromethyl-benzyl)-ureido]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;
[2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;
[2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-methoxymethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid propyl ester;

[2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

5 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;

[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

10 [2S,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester;


[2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

[2S,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-cyclopropyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester;

15 [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-formyl-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester; and

[2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-methyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester.

20 Another class of CETP inhibitors that finds utility with the present invention consists of cycloalkano-pyridines having the Formula VI



Formula VI


and pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds;

25 in which

A_{VI} denotes an aryl containing 6 to 10 carbon atoms, which is optionally substituted with up to five identical or different substituents in the form of a halogen, nitro, hydroxyl, trifluoromethyl, trifluoromethoxy or a straight-chain or branched alkyl, acyl, hydroxyalkyl or alkoxy containing up to 7 carbon atoms
30 each, or in the form of a group according to the formula -BNR_{VI-3}R_{VI-4}, wherein

R_{VI-3} and R_{VI-4} are identical or different and denote a hydrogen, phenyl or a straight-chain or branched alkyl containing up to 6 carbon atoms,

D_{VI} denotes an aryl containing 6 to 10 carbon atoms, which is optionally substituted with a phenyl, nitro, halogen, trifluoromethyl or trifluoromethoxy, or a radical according to the formula R_{VI-5} - L_{VI-1} .



or R_{VI-9} - T_{VI} - V_{VI} - X_{VI} , wherein

R_{VI-5} , R_{VI-6} and R_{VI-9} denote, independently from one another, a cycloalkyl containing 3 to 6 carbon atoms, or an aryl containing 6 to 10 carbon atom or a 5- to 7-membered, optionally benzo-condensed, saturated or unsaturated, mono-, bi- or tricyclic heterocycle containing up to 4 heteroatoms from the series of S, N and/or O, wherein the rings are optionally substituted, in the case of the nitrogen-containing rings also via the N function, with up to five identical or different substituents in the form of a halogen, trifluoromethyl, nitro, hydroxyl, cyano, carboxyl, trifluoromethoxy, a straight-chain or branched acyl, alkyl, alkylthio, alkylalkoxy, alkoxy or alkoxycarbonyl containing up to 6 carbon atoms each, an aryl or trifluoromethyl-substituted aryl containing 6 to 10 carbon atoms each, or an optionally benzo-condensed, aromatic 5- to 7-membered heterocycle containing up to 3 heteroatoms from the series of S, N and/or O, and/or in the form of a group according to the formula BOR_{VI-10} , $-SR_{VI-11}$, $-SO_2R_{VI-12}$ or $BNR_{VI-13}R_{VI-14}$, wherein

R_{VI-10} , R_{VI-11} and R_{VI-12} denote, independently from one another, an aryl containing 6 to 10 carbon atoms, which is in turn substituted with up to two identical or different substituents in the form of a phenyl, halogen or a straight-chain or branched alkyl containing up to 6 carbon atoms,

R_{VI-13} and R_{VI-14} are identical or different and have the meaning of R_{VI-3} and R_{VI-4} given above, or


R_{VI-5} and/or R_{VI-6} denote a radical according to the formula

45



R_{VI-7} denotes a hydrogen or halogen, and

R_{VI-8} denotes a hydrogen, halogen, azido, trifluoromethyl, hydroxyl, trifluoromethoxy, a straight-chain or branched alkoxy or alkyl containing up to 6 carbon atoms each, or a radical according to the formula



wherein

R_{VI-15} and R_{VI-16} are identical or different and have the meaning of R_{VI-3} and R_{VI-4} given above, or

R_{VI-7} and R_{VI-8} together form a radical according to the formula $=O$ or $=NR_{VI-17}$, wherein

R_{VI-17} denotes a hydrogen or a straight-chain or branched alkyl, alkoxy or acyl containing up to 6 carbon atoms each,

L_{VI} denotes a straight-chain or branched alkylene or alkenylene chain containing up to 8 carbon atoms each, which are optionally substituted with up to two hydroxyl groups,

T_{VI} and X_{VI} are identical or different and denote a straight-chain or branched alkylene chain containing up to 8 carbon atoms, or


T_{VI} or X_{VI} denotes a bond,

V_{VI} denotes an oxygen or sulfur atom or an BNR_{VI-18} group, wherein

R_{VI-18} denotes a hydrogen or a straight-chain or branched alkyl containing up to 6 carbon atoms or a phenyl,

E_{VI} denotes a cycloalkyl containing 3 to 8 carbon atoms, or a straight-chain or branched alkyl containing up to 8 carbon atoms, which is optionally substituted with a cycloalkyl containing 3 to 8 carbon atoms or a hydroxyl, or a phenyl, which is optionally substituted with a halogen or trifluoromethyl,

R_{VI-1} and R_{VI-2} together form a straight-chain or branched alkylene chain containing up to 7 carbon atoms, which must be substituted with a carbonyl group and/or a radical according to the formula



wherein

a and b are identical or different and denote a number equaling 1, 2 or 3,

R_{VI-19} denotes a hydrogen atom, a cycloalkyl containing 3 to 7 carbon atoms, a straight-chain or branched silylalkyl containing up to 8 carbon atoms, or a straight-chain or branched alkyl containing up to 8 carbon atoms, which is optionally substituted with a hydroxyl, a straight-chain or a branched alkoxy containing up to 6 carbon atoms or a phenyl, which may in turn be substituted with a halogen, nitro, trifluoromethyl, trifluoromethoxy or phenyl or tetrazole-substituted phenyl, and an alkyl that is optionally substituted with a group according to the formula BOR_{VI-22} , wherein


R_{VI-22} denotes a straight-chain or branched acyl containing up to 4 carbon atoms or benzyl, or

R_{VI-19} denotes a straight-chain or branched acyl containing up to 20 carbon atoms or benzoyl, which is optionally substituted with a halogen, trifluoromethyl, nitro or trifluoromethoxy, or a straight-chain or branched fluoroacyl containing up to 8 carbon atoms,

R_{VI-20} and R_{VI-21} are identical or different and denote a hydrogen, phenyl or a straight-chain or branched alkyl containing up to 6 carbon atoms, or

R_{VI-20} and R_{VI-21} together form a 3- to 6-membered carbocyclic ring, and the carbocyclic rings formed are optionally substituted, optionally also geminally, with up to six identical or different substituents in the form of trifluoromethyl, hydroxyl, nitrile, halogen, carboxyl, nitro, azido, cyano, cycloalkyl or cycloalkyloxy containing 3 to 7 carbon atoms each, a straight-chain or branched alkoxy, carbonyl, alkoxy or alkylthio containing up to 6 carbon atoms each, or a straight-chain or branched alkyl containing up to 6 carbon atoms, which is in turn substituted with up to two identical or different substituents in the form of a hydroxyl, benzyloxy, trifluoromethyl, benzoyl, a straight-chain or branched alkoxy, oxyacyl or carboxyl containing up to 4 carbon atoms each and/or a phenyl, which may in turn be substituted with a halogen, trifluoromethyl or trifluoromethoxy, and/or the carbocyclic rings formed are optionally substituted, also geminally, with up to five identical or different substituents in the form of a phenyl, benzoyl, thiophenyl or sulfonylbenzyl, which in turn are optionally substituted with a halogen, trifluoromethyl, trifluoromethoxy or nitro, and/or optionally in the form of a radical according to the formula

47




wherein

c is a number equaling 1, 2, 3 or 4,

5 d is a number equaling 0 or 1,

$\text{R}_{\text{VI}-23}$ and $\text{R}_{\text{VI}-24}$ are identical or different and denote a hydrogen, cycloalkyl containing 3 to 6 carbon atoms, a straight-chain or branched alkyl containing up to 6 carbon atoms, benzyl or phenyl, which is optionally substituted with up to two identical or different substituents in the form of halogen, trifluoromethyl, cyano, phenyl or nitro, and/or the carbocyclic rings formed are optionally substituted with a spiro-linked radical according to the formula



wherein

15 W_{VI} denotes either an oxygen atom or a sulfur atom, Y_{VI} and $\text{Y}=\text{VI}$ together form a 2- to 6-membered straight-chain or branched alkylene chain,


e is a number equaling 1, 2, 3, 4, 5, 6 or 7,

f is a number equaling 1 or 2,

20 $\text{R}_{\text{VI}-25}$, $\text{R}_{\text{VI}-26}$, $\text{R}_{\text{VI}-27}$, $\text{R}_{\text{VI}-28}$, $\text{R}_{\text{VI}-29}$, $\text{R}_{\text{VI}-30}$ and $\text{R}_{\text{VI}-31}$ are identical or different and denote a hydrogen, trifluoromethyl, phenyl, halogen or a straight-chain or branched alkyl or alkoxy containing up to 6 carbon atoms each, or

$\text{R}_{\text{VI}-25}$ and $\text{R}_{\text{VI}-26}$ or $\text{R}_{\text{VI}-27}$ and $\text{R}_{\text{VI}-28}$ each together denote a straight-chain or branched alkyl chain containing up to 6 carbon atoms or

25 $\text{R}_{\text{VI}-25}$ and $\text{R}_{\text{VI}-26}$ or $\text{R}_{\text{VI}-27}$ and $\text{R}_{\text{VI}-28}$ each together form a radical according to the formula



wherein

 W_{VI} has the meaning given above,

g is a number equaling 1, 2, 3, 4, 5, 6 or 7,

R_{VI-32} and R_{VI-33} together form a 3- to 7-membered heterocycle, which contains an oxygen or sulfur atom or a group according to the formula SO, SO₂ or BNR_{VI-34}, wherein

5 R_{VI-34} denotes a hydrogen atom, a phenyl, benzyl, or a straight-chain or branched alkyl containing up to 4 carbon atoms, and salts and N oxides thereof, with the exception of 5(6H)-quinolones, 3-benzoyl-7,8-dihydro-2,7,7-trimethyl-4-phenyl.


Compounds of Formula VI and their methods of manufacture are disclosed
10 in European Patent Application No. EP 818448 A1, United States Patent No. 6,207,671 and United States Patent No. 6,069,148, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from one of the following compounds of Formula VI:

15 2-cyclopentyl-4-(4-fluorophenyl)-7,7-dimethyl-3-(4-trifluoromethylbenzoyl)-4,6,7,8-tetrahydro-1H-quinolin-5-one;
2-cyclopentyl-4-(4-fluorophenyl)-7,7-dimethyl-3-(4-trifluoromethylbenzoyl)-7,8-dihydro-6H-quinolin-5-one;
[2-cyclopentyl-4-(4-fluorophenyl)-5-hydroxy-7,7-dimethyl-5,6,7,8-
20 tetrahydroquinolin-3-yl]-(4-trifluoromethylphenyl)-methanone;
[5-(t-butyldimethylsilyloxy)-2-cyclopentyl-4-(4-fluorophenyl)-7,7-dimethyl-5,6,7,8-tetrahydroquinolin-3-yl]-(4-trifluoromethylphenyl)-methanone;
[5-(t-butyldimethylsilyloxy)-2-cyclopentyl-4-(4-fluorophenyl)-7,7-dimethyl-5,6,7,8-tetrahydroquinolin-3-yl]-(4-trifluoromethylphenyl)-methanol;
25 5-(t-butyldimethylsilyloxy)-2-cyclopentyl-4-(4-fluorophenyl)-3-[fluoro-(4-trifluoromethylphenyl)-methyl]-7,7-dimethyl-5,6,7,8-tetrahydroquinoline; and
2-cyclopentyl-4-(4-fluorophenyl)-3-[fluoro-(4-trifluoromethylphenyl)-methyl]-7,7-dimethyl-5,6,7,8-tetrahydroquinolin-5-ol.

Another class of CETP inhibitors that finds utility with the present
30 invention consists of substituted-pyridines having the Formula VII

49




Formula VII

or a pharmaceutically acceptable salt or tautomer thereof,
wherein

5 R_{VII-2} and R_{VII-6} are independently selected from the group consisting of hydrogen, hydroxy, alkyl, fluorinated alkyl, fluorinated aralkyl, chlorofluorinated alkyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, alkoxy, alkoxyalkyl, and alkoxycarbonyl; provided that at least one of R_{VII-2} and R_{VII-6} is fluorinated alkyl, chlorofluorinated alkyl or alkoxyalkyl;

10 R_{VII-3} is selected from the group consisting of hydroxy, amido, arylcarbonyl, heteroarylcarbonyl, hydroxymethyl -CHO, -CO₂R_{VII-7}, wherein R_{VII-7} is selected from the group consisting of hydrogen, alkyl and cyanoalkyl; and




15 wherein R_{VII-15a} is selected from the group consisting of hydroxy, hydrogen, halogen, alkylthio, alkenylthio, alkynylthio, arylthio, heteroarylthio, heterocyclylthio, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy and heterocycloxy, and

20 R_{VII-16a} is selected from the group consisting of alkyl, haloalkyl, alkenyl, haloalkenyl, alkynyl, haloalkynyl, aryl, heteroaryl, and heterocyclyl, arylalkoxy, trialkylsilyloxy;

25 R_{VII-4} is selected from the group consisting of hydrogen, hydroxy, halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, aryl, heteroaryl, heterocyclyl, cycloalkylalkyl, cycloalkenylalkyl, aralkyl, heteroarylalkyl, heterocyclylalkyl, cycloalkylalkenyl, cycloalkenylalkenyl, aralkenyl, heteroarylalkenyl, heterocyclylalkenyl, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, heterocyclioxy, alkanoyloxy, alkenoyloxy, alkynoyloxy,


- aryloxy, heteroaryloxy, heterocyclyloxy, alkoxycarbonyl,
 alkenoxycarbonyl, alkynoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl,
 heterocyclyloxycarbonyl, thio, alkylthio, alkenylthio, alkynylthio, arylthio,
 heteroarylthio, heterocyclylthio, cycloalkylthio, cycloalkenylthio, alkylthioalkyl,
 5 alkenylthioalkyl, alkynylthioalkyl, arylthioalkyl, heteroarylthioalkyl,
 heterocyclylthioalkyl, alkylthioalkenyl, alkenylthioalkenyl, alkynylthioalkenyl,
 arylthioalkenyl, heteroarylthioalkenyl, heterocyclylthioalkenyl, alkylamino,
 alkenylamino, alkynylamino, arylamino, heteroarylamino, heterocyclylamino,
 arylalkylamino, diarylamino, diheteroarylamino, alkylarylamino,
 10 alkylheteroarylamino, arylheteroarylamino, trialkylsilyl, trialkenylsilyl, triarylsilyl,
 -CO(O)N(R_{VII-8a}R_{VII-8b}), wherein R_{VII-8a} and R_{VII-8b} are independently selected from
 the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl, -
 SO₂R_{VII-9}, wherein R_{VII-9} is selected from the group consisting of hydroxy, alkyl,
 alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl, -OP(O)(OR_{VII-10a})(OR_{VII-10b}),
 15 wherein R_{VII-10a} and R_{VII-10b} are independently selected from the group consisting
 of hydrogen, hydroxy, alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl,
 and -OP(S)(OR_{VII-11a})(OR_{VII-11b}), wherein R_{VII-11a} and R_{VII-11b} are independently
 selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and
 heterocyclyl;
- 20 R_{VII-5} is selected from the group consisting of hydrogen, hydroxy,
 halogen, alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl,
 haloalkynyl, aryl, heteroaryl, heterocyclyl, alkoxy, alkenoxy, alkynoxy, aryloxy,
 heteroaryloxy, heterocyclyloxy, alkylcarbonyloxyalkyl, alkenylcarbonyloxyalkyl,
 alkynylcarbonyloxyalkyl, arylcarbonyloxyalkyl, heteroarylcarbonyloxyalkyl,
 25 heterocyclylcarbonyloxyalkyl, cycloalkylalkyl, cycloalkenylalkyl, aralkyl,
 heteroarylalkyl, heterocyclylalkyl, cycloalkylalkenyl, cycloalkenylalkenyl,
 aralkenyl, heteroarylalkenyl, heterocyclylalkenyl, alkylthioalkyl,
 cycloalkylthioalkyl, alkenylthioalkyl, alkynylthioalkyl, arylthioalkyl,
 heteroarylthioalkyl, heterocyclylthioalkyl, alkylthioalkenyl, alkenylthioalkenyl,
 30 alkynylthioalkenyl, arylthioalkenyl, heteroarylthioalkenyl, heterocyclylthioalkenyl,
 alkoxyalkyl, alkenoxyalkyl, alkynoxyalkyl, aryloxyalkyl, heteroaryloxyalkyl,
 heterocyclyloxyalkyl, alkoxyalkenyl, alkenoxyalkenyl, alkynoxyalkenyl,
 aryloxyalkenyl, heteroaryloxyalkenyl, heterocyclyloxyalkenyl, cyano,
 hydroxymethyl, -CO₂R_{VII-14}, wherein R_{VII-14} is selected from the group consisting
 35 of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl;

51




wherein R_{VII-15b} is selected from the group consisting of hydroxy, hydrogen, halogen, alkylthio, alkenylthio, alkynylthio, arylthio, heteroarylthio, heterocyclylthio, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, heterocycliloxy, aroyloxy, and alkylsulfonyloxy, and

$R_{VII-16b}$ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, arylalkoxy, and trialkylsilyloxy;



wherein R_{VII-17} and R_{VII-18} are independently selected from the group
10 consisting of alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl;




wherein R_{VII-19} is selected from the group consisting of alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, -SR_{VII-20}, -OR_{VII-21}, and BR_{VII-22}CO₂R_{VII-23}, wherein

15 R_{VII-20} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aminoalkyl, aminoalkenyl, aminoalkynyl, aminoaryl, aminoheteroaryl, aminoheterocyclyl, alkylheteroarylamino, arylheteroarylamino,


R_{VII-21} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl,

20 R_{VII-22} is selected from the group consisting of alkylene or arylene, and


R_{VII-23} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;






wherein R_{VII-24} is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, aralkyl, aralkenyl, and aralkynyl;



5 wherein R_{VII-25} is heterocyclidenyl;




wherein R_{VII-26} and R_{VII-27} are independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;



10


wherein R_{VII-28} and R_{VII-29} are independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;

15





53


wherein R_{VII-30} and R_{VII-31} are independently alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, and heterocyclyloxy; and



5 wherein R_{VII-32} and R_{VII-33} are independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;




wherein R_{VII-36} is selected from the group consisting of alkyl, alkenyl, aryl, heteroaryl and heterocyclyl;



10

wherein R_{VII-37} and R_{VII-38} are independently selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl;



15

wherein R_{VII-39} is selected from the group consisting of hydrogen, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, heterocyclyloxy, alkylthio, alkenylthio, alkynylthio, arylthio, heteroarylthio and heterocyclylthio, and


R_{VII-40} is selected from the group consisting of haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl, haloheterocyclyl, cycloalkyl, cycloalkenyl, heterocyclylalkoxy, heterocyclylalkenoxy, heterocyclylalkynoxy, alkylthio, alkenylthio, alkynylthio, arylthio, heteroarylthio and heterocyclylthio;

20




wherein R_{VII-41} is heterocyclidenyl;

54



wherein R_{VII-42} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, and heterocyclyl, and

R_{VII-43} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, cycloalkyl, cycloalkenyl, haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl, and haloheterocyclyl;



wherein R_{VII-44} is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl;

10 - N = S = O;

$$\cdot N = C = S:$$
$$-N = C = O:$$
$$-N=C=O;$$

- N₃;

- SR_{VII-45}


15 wherein R_{VII-45} is selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl, haloheterocyclyl, heterocyclyl, cycloalkylalkyl, cycloalkenylalkyl, aralkyl, heteroarylalkyl, heterocyclylalkyl, cycloalkylalkenyl, cycloalkenylalkenyl, aralkenyl, heteroarylalkenyl, heterocyclylalkenyl, alkylthioalkyl, alkenylthioalkyl, alkynylthioalkyl, arylthioalkyl, heteroarylthioalkyl, heterocyclylthioalkyl, alkylthioalkenyl, alkenylthioalkenyl, alkynylthioalkenyl, arylthioalkenyl, heteroarylthioalkenyl, heterocyclylthioalkenyl, aminocarbonylalkyl, aminocarbonylalkenyl, aminocarbonylalkynyl, aminocarbonylaryl, aminocarbonylheteroaryl, and aminocarbonylheterocyclyl,

$$-\text{SR}_{\text{VII-46}}, \text{ and } -\text{CH}_2\text{R}_{\text{VII-47}},$$

wherein R_{VII-46} is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl, and


R_{VII-47} is selected from the group consisting of hydrogen, alkyl, alkenyl, 30 alkynyl, aryl, heteroaryl and heterocyclyl; and

55




wherein $\text{R}_{\text{VII-48}}$ is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl, and


$\text{R}_{\text{VII-49}}$ is selected from the group consisting of alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy, heterocyclyloxy, haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl and haloheterocyclyl;



wherein $\text{R}_{\text{VII-50}}$ is selected from the group consisting of hydrogen, alkyl, cycloalkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, alkoxy, alkenoxy, alkynoxy, aryloxy, heteroaryloxy and heterocyclyloxy;



wherein $\text{R}_{\text{VII-51}}$ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl, heterocyclyl, haloalkyl, haloalkenyl, haloalkynyl, haloaryl, haloheteroaryl and haloheterocyclyl; and



wherein $\text{R}_{\text{VII-53}}$ is selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, heteroaryl and heterocyclyl;


provided that when $\text{R}_{\text{VII-5}}$ is selected from the group consisting of heterocyclylalkyl and heterocyclylalkenyl, the heterocyclyl radical of the corresponding heterocyclylalkyl or heterocyclylalkenyl is other than δ -lactone; and

provided that when $\text{R}_{\text{VII-4}}$ is aryl, heteroaryl or heterocyclyl, and one of $\text{R}_{\text{VII-2}}$ and $\text{R}_{\text{VII-6}}$ is trifluoromethyl, then the other of $\text{R}_{\text{VII-2}}$ and $\text{R}_{\text{VII-6}}$ is difluoromethyl.

Compounds of Formula VII and their methods of manufacture are disclosed in PCT Publication No. WO 9941237-A1, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor of Formula VII is dimethyl
 5 5,5-dithiobis[2-difluoromethyl-4-(2-methylpropyl)-6-(trifluoromethyl)-3-pyridine-carboxylate].

Another class of CETP inhibitors that finds utility with the present invention consists of substituted biphenyls having the Formula VIII



Formula VIII

10 or a pharmaceutically acceptable salt, enantiomers, or stereoisomers thereof, in which

A_{VIII} stands for aryl with 6 to 10 carbon atoms, which is optionally substituted up to 3 times in an identical manner or differently by halogen, hydroxy, trifluoromethyl, trifluoromethoxy, or by straight-chain or branched alkyl,
 15 acyl, or alkoxy with up to 7 carbon atoms each, or by a group of the formula

-NR_{VIII-1}R_{VIII-2}, wherein

R_{VIII-1} and R_{VIII-2} are identical or different and denote hydrogen, phenyl, or straight-chain or branched alkyl with up to 6 carbon atoms,

D_{VIII} stands for straight-chain or branched alkyl with up to 8 carbon
 20 atoms, which is substituted by hydroxy,

E_{VIII} and L_{VIII} are either identical or different and stand for straight-chain or branched alkyl with up to 8 carbon atoms, which is optionally substituted by cycloalkyl with 3 to 8 carbon atoms, or stands for cycloalkyl with 3 to 8 carbon atoms, or

25 E_{VIII} has the above-mentioned meaning and

L_{VIII} in this case stands for aryl with 6 to 10 carbon atoms, which is optionally substituted up to 3 times in an identical manner or differently by halogen, hydroxy, trifluoromethyl, trifluoromethoxy, or by straight-chain or branched alkyl, acyl, or alkoxy with up to 7 carbon atoms each, or by a group of
 30 the formula

-NR_{VIII-3}R_{VIII-4}, wherein

R_{VIII-3} and R_{VIII-4} are identical or different and have the meaning given above for R_{VIII-1} and R_{VIII-2}, or


5 E_{VIII} stands for straight-chain or branched alkyl with up to 8 carbon atoms, or stands for aryl with 6 to 10 carbon atoms, which is optionally substituted up to 3 times in an identical manner or differently by halogen, hydroxy, trifluoromethyl, trifluoromethoxy, or by straight-chain or branched alkyl, acyl, or alkoxy with up to 7 carbon atoms each, or by a group of the formula

-NR_{VIII-5}R_{VIII-6}, wherein

10 R_{VIII-5} and R_{VIII-6} are identical or different and have the meaning given above for R_{VIII-1} and R_{VIII-2}, and

L_{VIII} in this case stands for straight-chain or branched alkoxy with up to 8 carbon atoms or for cycloalkyloxy with 3 to 8 carbon atoms,

T_{VIII} stands for a radical of the formula



R_{VIII-7} and R_{VIII-8} are identical or different and denote cycloalkyl with 3 to 8 carbon atoms, or aryl with 6 to 10 carbon atoms, or denote a 5- to 7-member aromatic, optionally benzo-condensed, heterocyclic compound with up to 3 heteroatoms from the series S, N and/or O, which are optionally substituted up to 3 times in an identical manner or differently by trifluoromethyl, trifluoromethoxy, halogen, hydroxy, carboxyl, by straight-chain or branched alkyl, acyl, alkoxy, or alkoxycarbonyl with up to 6 carbon atoms each, or by phenyl, phenoxy, or thiophenyl, which can in turn be substituted by halogen, trifluoromethyl, or trifluoromethoxy, and/or the rings are substituted by a group of the formula

20

25

-NR_{VIII-11}R_{VIII-12}, wherein

R_{VIII-11} and R_{VIII-12} are identical or different and have the meaning given above for R_{VIII-1} and R_{VIII-2},

30 X_{VIII} denotes a straight or branched alkyl chain or alkenyl chain with 2 to 10 carbon atoms each, which are optionally substituted up to 2 times by hydroxy,

R_{VIII-9} denotes hydrogen, and

$R_{VIII-10}$ denotes hydrogen, halogen, azido, trifluoromethyl, hydroxy, mercapto, trifluoromethoxy, straight-chain or branched alkoxy with up to 5 carbon atoms, or a radical of the formula


$-NR_{VIII-13}R_{VIII-14}$, wherein

5 $R_{VIII-13}$ and $R_{VIII-14}$ are identical or different and have the meaning given above for R_{VIII-1} and R_{VIII-2} , or

R_{VIII-9} and $R_{VIII-10}$ form a carbonyl group together with the carbon atom.

Compounds of Formula VIII are disclosed in PCT Publication No. WO 9804528, which is incorporated herein by reference in its entirety for all
10 purposes.

Another class of CETP inhibitors that finds utility with the present invention consists of substituted 1,2,4-triazoles having the Formula IX



Formula IX

or a pharmaceutically acceptable salt or tautomer thereof;

15 wherein R_{IX-1} is selected from higher alkyl, higher alkenyl, higher alkynyl, aryl, aralkyl, aryloxyalkyl, alkoxyalkyl, alkylthioalkyl, arylthioalkyl, and cycloalkylalkyl;

wherein R_{IX-2} is selected from aryl, heteroaryl, cycloalkyl, and cycloalkenyl, wherein

20 R_{IX-2} is optionally substituted at a substitutable position with one or more radicals independently selected from alkyl, haloalkyl, alkylthio, alkylsulfinyl, alkylsulfonyl, alkoxy, halo, aryloxy, aralkyloxy, aryl, aralkyl, aminosulfonyl, amino, monoalkylamino and dialkylamino; and

wherein R_{IX-3} is selected from hydrido, -SH and halo;

25 provided R_{IX-2} cannot be phenyl or 4-methylphenyl when R_{IX-1} is higher alkyl and when R_{IX-3} is BSH.

Compounds of Formula IX and their methods of manufacture are disclosed in PCT Publication No. WO 9914204, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula IX:

- 2,4-dihydro-4-(3-methoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-4-(2-fluorophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 5 2,4-dihydro-4-(2-methylphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-4-(3-chlorophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2, 4-dihydro-4-(2-methoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-4-(3-methylphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 4-cyclohexyl-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 10 2,4-dihydro-4-(3-pyridyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-4-(2-ethoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-4-(2,6-dimethylphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-4-(4-phenoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 4-(1,3-benzodioxol-5-yl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-
- 15 thione;
- 4-(2-chlorophenyl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-4-(4-methoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-5-tridecyl-4-(3-trifluoromethylphenyl)-3H-1,2,4-triazole-3-
- thione;
- 20 2,4-dihydro-5-tridecyl-4-(3-fluorophenyl)-3H-1,2,4-triazole-3-thione;
- 4-(3-chloro-4-methylphenyl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-
- thione;
- 2,4-dihydro-4-(2-methylthiophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 4-(4-benzyloxyphenyl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 25 2,4-dihydro-4-(2-naphthyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-5-tridecyl-4-(4-trifluoromethylphenyl)-3H-1,2,4-triazole-3-
- thione;
- 2,4-dihydro-4-(1-naphthyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-4-(3-methylthiophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 30 2,4-dihydro-4-(4-methylthiophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;
- 2,4-dihydro-4-(3,4-dimethoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-
- thione;
- 2,4-dihydro-4-(2,5-dimethoxyphenyl)-5-tridecyl-3H-1,2,4-triazole-3-
- thione;

2,4-dihydro-4-(2-methoxy-5-chlorophenyl)-5-tridecyl-3H-1,2,4-triazole-3-thione;


4-(4-aminosulfonylphenyl)-2,4-dihydro-5-tridecyl-3H-1,2,4-triazole-3-thione;

5 2,4-dihydro-5-dodecyl-4-(3-methoxyphenyl)-3H-1,2,4-triazole-3-thione;
2,4-dihydro-4-(3-methoxyphenyl)-5-tetradecyl-3H-1,2,4-triazole-3-thione;
2,4-dihydro-4-(3-methoxyphenyl)-5-undecyl-3H-1,2,4-triazole-3-thione;

and

2,4-dihydro-(4-methoxyphenyl)-5-pentadecyl-3H-1,2,4-triazole-3-thione.

10 Another class of CETP inhibitors that finds utility with the present invention consists of hetero-tetrahydroquinolines having the Formula X



Formula X

and pharmaceutically acceptable salts, enantiomers, or stereoisomers or N-oxides of said compounds;

15 in which

A_x represents cycloalkyl with 3 to 8 carbon atoms or a 5 to 7-membered, saturated, partially saturated or unsaturated, optionally benzo-condensed heterocyclic ring containing up to 3 heteroatoms from the series comprising S, N and/or O, that in case of a saturated heterocyclic ring is bonded to a nitrogen function, optionally bridged over it, and in which the aromatic systems mentioned above are optionally substituted up to 5-times in an identical or different substituents in the form of halogen, nitro, hydroxy, trifluoromethyl, trifluoromethoxy or by a straight-chain or branched alkyl, acyl, hydroxyalkyl or alkoxy each having up to 7 carbon atoms or by a group of the formula BNR_x.

25 ₃R_{x-4},


in which

R_{x-3} and R_{x-4} are identical or different and denote hydrogen, phenyl or straight-chain or branched alkyl having up to 6 carbon atoms,


or

30

A_x represents a radical of the formula



- 5 D_x represents an aryl having 6 to 10 carbon atoms, that is optionally substituted by phenyl, nitro, halogen, trifluoromethyl or trifluoromethoxy, or it represents a radical of the formula



- 10 in which

- R_{x-5} , R_{x-6} and R_{x-9} independently of one another denote cycloalkyl having 3 to 6 carbon atoms, or an aryl having 6 to 10 carbon atoms or a 5- to 7-membered aromatic, optionally benzo-condensed saturated or unsaturated, mono-, bi-, or tricyclic heterocyclic ring from the series consisting of S, N and/or O, in which the rings are substituted, optionally, in case of the nitrogen containing aromatic rings via the N function, with up to 5 identical or different substituents in the form of halogen, trifluoromethyl, nitro, hydroxy, cyano, carbonyl, trifluoromethoxy, straight straight-chain or branched acyl, alkyl, alkylthio, alkylalkoxy, alkoxy, or alkoxycarbonyl each having up to 6 carbon atoms, by aryl or trifluoromethyl-substituted aryl each having 6 to 10 carbon atoms or by an, optionally benzo-condensed, aromatic 5- to 7-membered heterocyclic ring having up to 3 heteroatoms from the series consisting of S, N, and/or O, and/or substituted by a group of the formula BOR_{x-10} , $-SR_{x-11}$, SO_2R_{x-12} or $BNR_{x-13}R_{x-14}$,

- 25 in which



R_{x-10} , R_{x-11} and R_{x-12} independently from each other denote aryl having 6 to 10 carbon atoms, which is in turn substituted with up to 2 identical or different

substituents in the form of phenyl, halogen or a straight-chain or branched alkyl having up to 6 carbon atoms,

R_{X-13} and R_{X-14} are identical or different and have the meaning of R_{X-3} and R_{X-4} indicated above,

5 or

R_{X-5} and/or R_{X-6} denote a radical of the formula



or

R_{X-7} denotes hydrogen or halogen, and

10 R_{X-8} denotes hydrogen, halogen, azido, trifluoromethyl, hydroxy, trifluoromethoxy, straight-chain or branched alkoxy or alkyl having up to 6 carbon atoms or a radical of the formula

$\text{BNR}_{X-15}\text{R}_{X-16}$,

in which

15 R_{X-15} and R_{X-16} are identical or different and have the meaning of R_{X-3} and R_{X-4} indicated above,

or

R_{X-7} and R_{X-8} together form a radical of the formula $=\text{O}$ or $=\text{NR}_{X-17}$,

in which

20 R_{X-17} denotes hydrogen or straight chain or branched alkyl, alkoxy or acyl having up to 6 carbon atoms,

L_X denotes a straight chain or branched alkylene or alkenylene chain having up to 8 carbon atoms, that are optionally substituted with up to 2 hydroxy groups,

25 T_X and X_X are identical or different and denote a straight chain or branched alkylene chain with up to 8 carbon atoms

or


T_X or X_X denotes a bond,

V_X represents an oxygen or sulfur atom or an BNR_{X-18} -group, in which

30 R_{X-18} denotes hydrogen or straight chain or branched alkyl with up to 6 carbon atoms or phenyl,

E_x represents cycloalkyl with 3 to 8 carbon atoms, or straight chain or branched alkyl with up to 8 carbon atoms, that is optionally substituted by cycloalkyl with 3 to 8 carbon atoms or hydroxy, or represents a phenyl, that is optionally substituted by halogen or trifluoromethyl,

- 5 R_{x-1} and R_{x-2} together form a straight-chain or branched alkylene chain with up to 7 carbon atoms, that must be substituted by carbonyl group and/or by a radical with the formula



- 10 in which a and b are identical or different and denote a number equaling 1, 2, or 3,

- 15 R_{x-19} denotes hydrogen, cycloalkyl with 3 up to 7 carbon atoms, straight chain or branched silylalkyl with up to 8 carbon atoms or straight chain or branched alkyl with up to 8 carbon atoms, that are optionally substituted by hydroxyl, straight chain or branched alkoxy with up to 6 carbon atoms or by phenyl, which in turn might be substituted by halogen, nitro, trifluoromethyl, trifluoromethoxy or by phenyl or by tetrazole-substituted phenyl, and alkyl, optionally be substituted by a group with the formula BOR_{x-22} , in which



- 20 R_{x-22} denotes a straight chain or branched acyl with up to 4 carbon atoms or benzyl, or

- 25 R_{x-19} denotes straight chain or branched acyl with up to 20 carbon atoms or benzoyl, that is optionally substituted by halogen, trifluoromethyl, nitro or trifluoromethoxy, or it denotes straight chain or branched fluoroacyl with up to 8 carbon atoms and 9 fluorine atoms,

R_{x-20} and R_{x-21} are identical or different and denote hydrogen, phenyl or straight chain or branched alkyl with up to 6 carbon atoms, or

- 30 R_{x-20} and R_{x-21} together form a 3- to 6- membered carbocyclic ring, and the carbocyclic rings formed are optionally substituted, optionally also geminally, with up to six identical or different substituents in the form of trifluoromethyl, hydroxy, nitrile, halogen, carboxyl, nitro, azido, cyano, cycloalkyl or

cycloalkyloxy with 3 to 7 carbon atoms each, by straight chain or branched
alkoxycarbonyl, alkoxy or alkylthio with up to 6 carbon atoms each or by straight
chain or branched alkyl with up to 6 carbon atoms, which in turn is substituted
with up to 2 identically or differently by hydroxyl, benzyloxy, trifluoromethyl,
5 benzoyl, straight chain or branched alkoxy, oxyacyl or carbonyl with up to 4
carbon atoms each and/or phenyl, which may in turn be substituted with a
halogen, trifluoromethyl or trifluoromethoxy, and/or the formed carbocyclic rings
are optionally substituted, also geminally, with up to 5 identical or different
substituents in the form of phenyl, benzoyl, thiophenyl or sulfonylbenzyl, which
10 in turn are optionally substituted by halogen, trifluoromethyl, trifluoromethoxy or
nitro, and/or optionally are substituted by a radical with the formula




in which

15 c denotes a number equaling 1, 2, 3, or 4,

d denotes a number equaling 0 or 1,

R_{X-23} and R_{X-24} are identical or different and denote hydrogen, cycloalkyl
with 3 to 6 carbon atoms, straight chain or branched alkyl with up to 6 carbon
atoms, benzyl or phenyl, that is optionally substituted with up to 2 identically or
20 differently by halogen, trifluoromethyl, cyano, phenyl or nitro, and/or the formed
carbocyclic rings are substituted optionally by a spiro-linked radical with the
formula



25 in which

W_X denotes either an oxygen or a sulfur atom

Y_X and Y'_X together form a 2 to 6 membered straight chain or branched
alkylene chain,


e denotes a number equaling 1, 2, 3, 4, 5, 6, or 7,

30 f denotes a number equaling 1 or 2,

R_{X-25} , R_{X-26} , R_{X-27} , R_{X-28} , R_{X-29} , R_{X-30} and R_{X-31} are identical or different and denote hydrogen, trifluoromethyl, phenyl, halogen or straight chain or branched alkyl or alkoxy with up to 6 carbon atoms each,
or

5 R_{X-25} and R_{X-26} or R_{X-27} and R_{X-28} respectively form together a straight chain or branched alkyl chain with up to 6 carbon atoms,
or

R_{X-25} and R_{X-26} or R_{X-27} and R_{X-28} each together form a radical with the formula



10

in which

W_X has the meaning given above,

g denotes a number equaling 1, 2, 3, 4, 5, 6, or 7,

15 R_{X-32} and R_{X-33} form together a 3- to 7- membered heterocycle, which contains an oxygen or sulfur atom or a group with the formula SO , SO_2 or $-NR_{X-34}$,

in which

R_{X-34} denotes hydrogen, phenyl, benzyl or straight or branched alkyl with up to 4 carbon atoms.

20

Compounds of Formula X and their methods of manufacture are disclosed in PCT Publication No. WO 9914215, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula X:


25

2-cyclopentyl-5-hydroxy-7,7-dimethyl-4-(3-thienyl)-3-(4-trifluoromethylbenzoyl)-5,6,7,8-tetrahydroquinoline;

2-cyclopentyl-3-[fluoro-(4-trifluoromethylphenyl)methyl]-5-hydroxy-7,7-dimethyl-4-(3-thienyl)-5,6,7,8-tetrahydroquinoline; and

30 2-cyclopentyl-5-hydroxy-7,7-dimethyl-4-(3-thienyl)-3-(trifluoromethylbenzyl)-5,6,7,8-tetrahydroquinoline.


Another class of CETP inhibitors that finds utility with the present invention consists of substituted tetrahydro naphthalines and analogous compound having the Formula XI



Formula XI

and stereoisomers, stereoisomer mixtures, and salts thereof, in which


A_{XI} stands for cycloalkyl with 3 to 8 carbon atoms, or stands for aryl with 6 to 10 carbon atoms, or stands for a 5- to 7-membered, saturated, partially unsaturated or unsaturated, possibly benzocondensated, heterocycle with up to 4 heteroatoms from the series S, N and/or O, where aryl and the heterocyclic ring systems mentioned above are substituted up to 5-fold, identical or different, by cyano, halogen, nitro, carboxyl, hydroxy, trifluoromethyl, trifluoro-methoxy, or by straight-chain or branched alkyl, acyl, hydroxyalkyl, alkylthio, alkoxycarbonyl, oxyalkoxycarbonyl or alkoxy each with up to 7 carbon atoms, or by a group of the formula



in which

R_{XI-3} and R_{XI-4} are identical or different and denote hydrogen, phenyl, or straight-chain or branched alkyl with up to 6 carbon atoms

D_{XI} stands for a radical of the formula



in which

R_{XI-5}, R_{XI-6} and R_{XI-9}, independent of each other, denote cycloalkyl with 3 to 6 carbon atoms, or denote aryl with 6 to 10 carbon atoms, or denote a 5- to 7-membered, possibly benzocondensated, saturated or unsaturated, mono-, bi- or tricyclic heterocycle with up to 4 heteroatoms of the series S, N and/or O, where the cycles are possibly substituted in the case of the nitrogen-containing rings

also via the N-functionCup to 5-fold, identical or different, by halogen, trifluoromethyl, nitro, hydroxy, cyano, carboxyl, trifluoromethoxy, straight-chain or branched acyl, alkyl, alkylthio, alkylalkoxy, alkoxy or alkoxycarbonyl with up to 6 carbon atoms each, by aryl or trifluoromethyl substituted aryl with 6 to 10 carbon atoms each, or by a possibly benzocondensated aromatic 5- to 7-membered heterocycle with up to 3 heteroatoms of the series S, N and/or O, and/or are substituted by a group of the formula

$-OR_{XI-10}$, $-SR_{XI-11}$, $-SO_2R_{XI-12}$ or $-NR_{XI-13}R_{XI-14}$,

in which

R_{XI-10} , R_{XI-11} and R_{XI-12} , independent of each other, denote aryl with 6 to 10 carbon atoms, which itself is substituted up to 2-fold, identical or different, by phenyl, halogen, or by straight-chain or branched alkyl with up to 6 carbon atoms,

R_{XI-13} and R_{XI-14} are identical or different and have the meaning given above for R_{XI-3} and R_{XI-4} ,

or

R_{XI-5} and/or R_{XI-6} denote a radical of the formula



R_{XI-7} denotes hydrogen, halogen or methyl,

and

R_{XI-8} denotes hydrogen, halogen, azido, trifluoromethyl, hydroxy, trifluoromethoxy, straight-chain or branched alkoxy or alkyl with up to 6 carbon atoms each, or a radical of the formula $-NR_{XI-15}R_{XI-16}$,

in which

R_{XI-15} and R_{XI-16} are identical or different and have the meaning given above for R_{XI-3} and R_{XI-4} ,

or

R_{XI-7} and R_{XI-8} together form a radical of the formula $=O$ or $=NR_{XI-17}$, in which

R_{XI-17} denotes hydrogen or straight-chain or branched alkyl, alkoxy or acyl with up to 6 carbon atoms each,

L_{X1} denotes a straight-chain or branched alkylene- or alkenylene chain with up to 8 carbon atoms each, which is possibly substituted up to 2-fold by hydroxy,

T_{X1} and X_{X1} are identical or different and denote a straight-chain or branched alkylene chain with up to 8 carbon atoms,
5 or


T_{X1} and X_{X1} denotes a bond,

V_{X1} stands for an oxygen- or sulfur atom or for an $-NR_{X1-18}$ group,
in which

10 R_{X1-18} denotes hydrogen or straight-chain or branched alkyl with up to 6 carbon atoms, or phenyl,

E_{X1} stands for cycloalkyl with 3 to 8 carbon atoms, or stands for straight-chain or branched alkyl with up to 8 carbon atoms, which is possibly substituted by cycloalkyl with 3 to 8 carbon atoms or hydroxy, or stands for phenyl, which is
15 possibly substituted by halogen or trifluoromethyl,

R_{X1-1} and R_{X1-2} together form a straight-chain or branched alkylene chain with up to 7 carbon atoms, which must be substituted by a carbonyl group and/or by a radical of the formula



20 in which

a and b are identical or different and denote a number 1, 2 or 3

R_{X1-19} denotes hydrogen, cycloalkyl with 3 to 7 carbon atoms, straight-chain or branched silylalkyl with up to 8 carbon atoms, or straight-chain or branched alkyl with up to 8 carbon atoms, which is possibly substituted by
25 hydroxy, straight-chain or branched alkoxy with up to 6 carbon atoms, or by phenyl, which itself can be substituted by halogen, nitro, trifluoromethyl, trifluoromethoxy or by phenyl substituted by phenyl or tetrazol, and alkyl is possibly substituted by a group of the formula $-OR_{X1-22}$,



in which

30 R_{X1-22} denotes straight-chain or branched acyl with up to 4 carbon atoms, or benzyl,
or

R_{X1-19} denotes straight-chain or branched acyl with up to 20 carbon atoms or benzoyl, which is possibly substituted by halogen, trifluoromethyl, nitro or trifluoromethoxy, or denotes straight-chain or branched fluoroacyl with up to 8 carbon atoms and 9 fluorine atoms,

5 R_{X1-20} and R_{X1-21} are identical or different, denoting hydrogen, phenyl or straight-chain or branched alkyl with up to 6 carbon atoms, or

R_{X1-20} and R_{X1-21} together form a 3- to 6-membered carbocycle, and, possibly also geminally, the alkylene chain formed by R_{X1-1} and R_{X1-2} , is possibly
 10 substituted up to 6-fold, identical or different, by trifluoromethyl, hydroxy, nitrile, halogen, carboxyl, nitro, azido, cyano, cycloalkyl or cycloalkyloxy with 3 to 7 carbon atoms each, by straight-chain or branched alkoxy carbonyl, alkoxy or alkoxythio with up to 6 carbon atoms each, or by straight-chain or branched alkyl with up to 6 carbon atoms, which itself is substituted up to 2-fold,
 15 identical or different, by hydroxyl, benzyloxy, trifluoromethyl, benzoyl, straight-chain or branched alkoxy, oxyacyl or carboxyl with up to 4 carbon atoms each, and/or phenyl- which itself can be substituted by halogen, trifluoromethyl or trifluoromethoxy, and/or the alkylene chain formed by R_{X1-1} and R_{X1-2} is substituted, also geminally, possibly up to 5-fold, identical or different, by
 20 phenyl, benzoyl, thiophenyl or sulfobenzyl -which themselves are possibly substituted by halogen, trifluoromethyl, trifluoromethoxy or nitro, and/or the alkylene chain formed by R_{X1-1} and R_{X1-2} is possibly substituted by a radical of the formula



in which


c denotes a number 1, 2, 3 or 4,

d denotes a number 0 or 1,

30 R_{X1-23} and R_{X1-24} are identical or different and denote hydrogen, cycloalkyl with 3 to 6 carbon atoms, straight-chain or branched alkyl with up to 6 carbon atoms, benzyl or phenyl, which is possibly substituted up to 2-fold, identical or different, by halogen, trifluoromethyl, cyano, phenyl or nitro, and/or the alkylene

70

chain formed by R_{X1-1} and R_{X1-2} is possibly substituted by a spiro-jointed radical of the formula



5 in which

W_{X1} denotes either an oxygen or a sulfur atom,

Y_{X1} and Y'_{X1} together form a 2- to 6-membered straight-chain or branched alkylene chain,

e is a number 1, 2, 3, 4, 5, 6 or 7,

10 f denotes a number 1 or 2,


R_{X1-25} , R_{X1-26} , R_{X1-27} , R_{X1-28} , R_{X1-29} , R_{X1-30} and R_{X1-31} are identical or different and denote hydrogen, trifluoromethyl, phenyl, halogen, or straight-chain or branched alkyl or alkoxy with up to 6 carbon atoms each,

or

15 R_{X1-25} and R_{X1-26} or R_{X1-27} and R_{X1-28} together form a straight-chain or branched alkyl chain with up to 6 carbon atoms,

or

R_{X1-25} and R_{X1-26} or R_{X1-27} and R_{X1-28} together form a radical of the formula



20 in which

W_{X1} has the meaning given above,

g is a number 1, 2, 3, 4, 5, 6 or 7,


R_{X1-32} and R_{X1-33} together form a 3- to 7-membered heterocycle that contains an oxygen- or sulfur atom or a group of the formula SO, SO₂ or -NR_{X1-34},

25 in which

R_{X1-34} denotes hydrogen, phenyl, benzyl, or straight-chain or branched alkyl with up to 4 carbon atoms.

Compounds of Formula XI and their methods of manufacture are disclosed in PCT Publication No. WO 9914174, which is incorporated herein by reference in its entirety for all purposes.

Another class of CETP inhibitors that finds utility with the present invention consists of 2-aryl-substituted pyridines having the Formula (XII)



Formula XII

or pharmaceutically acceptable salts, enantiomers, or stereoisomers of said compounds,

in which

A_{XII} and E_{XII} are identical or different and stand for aryl with 6 to 10 carbon atoms which is possibly substituted, up to 5-fold identical or different, by halogen, hydroxy, trifluoromethyl, trifluoromethoxy, nitro or by straight-chain or branched alkyl, acyl, hydroxy alkyl or alkoxy with up to 7 carbon atoms each, or by a group of the formula $-NR_{XII-1}R_{XII-2}$,


where

R_{XII-1} and R_{XII-2} are identical or different and are meant to be hydrogen, phenyl or straight-chain or branched alkyl with up to 6 carbon atoms,

D_{XII} stands for straight-chain or branched alkyl with up to 8 carbon atoms, which is substituted by hydroxy,

L_{XII} stands for cycloalkyl with 3 to 8 carbon atoms or for straight-chain or branched alkyl with up to 8 carbon atoms, which is possibly substituted by cycloalkyl with 3 to 8 carbon atoms, or by hydroxy,

T_{XII} stands for a radical of the formula $R_{XII-3}-X_{XII}^{\cdot}$ or



where

R_{XII-3} and R_{XII-4} are identical or different and are meant to be cycloalkyl

with 3 to 8 carbon atoms, or aryl with 6 to 10 carbon atoms, or a 5- to 7-membered aromatic, possibly benzocondensated heterocycle with up to 3 heteroatoms from the series S, N and/or O, which are possibly substituted. up to 3-fold identical or different, by trifluoromethyl, trifluoromethoxy, halogen, hydroxy, carboxyl, nitro, by straight-chain or branched alkyl, acyl, alkoxy or alkoxy carbonyl with up to 6 carbon atoms each. or by phenyl, phenoxy or phenylthio which in turn can be substituted by halogen. trifluoromethyl or trifluoromethoxy, and/or where the cycles are possibly substituted by a group of the formula $-NR_{XII-7}R_{XII-8}$,

10 where

R_{XII-7} and R_{XII-8} are identical or different and have the meaning of R_{XII-1} and R_{XII-2} given above,

X_{XII} is a straight-chain or branched alkyl or alkenyl with 2 to 10 carbon atoms each, possibly substituted up to 2-fold by hydroxy or halogen,

15 R_{XII-5} stands for hydrogen.

and

R_{XII-6} means to be hydrogen, halogen, mercapto, azido, trifluoromethyl, hydroxy, trifluoromethoxy, straight-chain or branched alkoxy with up to 5 carbon atoms, or a radical of the formula $BNR_{XII-9}R_{XII-10}$.

20 where

R_{XII-9} and R_{XII-10} are identical or different and have the meaning of R_{XII-1} and R_{XII-2} given above,

or

R_{XII-5} and R_{XII-6} , together with the carbon atom, form a carbonyl group.

25 Compounds of Formula XII and their methods of manufacture are disclosed in EP 796846-A1, United States Patent No. 6,127,383 and United States Patent No. 5,925,645, all of which are incorporated herein by reference in their entireties for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula XII:

4,6-bis-(p-fluorophenyl)-2-isopropyl-3-[(p-trifluoromethylphenyl)-(fluoro)-methyl]-5-(1-hydroxyethyl)pyridine;


2,4-bis-(4-fluorophenyl)-6-isopropyl-5-[4-(trifluoromethylphenyl)-fluoromethyl]-3-hydroxymethylpyridine; and

35 2,4-bis-(4-fluorophenyl)-6-isopropyl-5-[2-(3-trifluoromethylphenyl)vinyl]-

3-hydroxymethyl)pyridine.

Another class of CETP inhibitors that finds utility with the present invention consists of compounds having the Formula (XIII)

5



Formula XIII

or pharmaceutically acceptable salts, enantiomers, stereoisomers, hydrates, or solvates of said compounds, in which

10

R_{XIII} is a straight chain or branched C_{1-10} alkyl; straight chain or branched C_{2-10} alkenyl; halogenated C_{1-4} lower alkyl; C_{3-10} cycloalkyl that may be substituted; C_{5-8} cycloalkenyl that may be substituted; C_{3-10} cycloalkyl C_{1-10} alkyl that may be substituted; aryl that may be substituted; aralkyl that may be substituted; or a 5- or 6-membered heterocyclic group having 1 to 3 nitrogen

15

atoms, oxygen atoms or sulfur atoms that may be substituted, X_{XIII-1} , X_{XIII-2} , X_{XIII-3} , X_{XIII-4} may be the same or different and are a hydrogen atom; halogen atom; C_{1-4} lower alkyl; halogenated C_{1-4} lower alkyl; C_{1-4} lower alkoxy; cyano group; nitro group; acyl; or aryl, respectively;

Y_{XIII} is $-CO-$; or BSO_2- ; and

20

Z_{XIII} is a hydrogen atom; or mercapto protective group.

Compounds of Formula XIII and their methods of manufacture are disclosed in PCT Publication No. WO 98/35937, which is incorporated herein by reference in its entirety for all purposes.

25

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula XIII:

N,N' -(dithiodi-2,1-phenylene)bis[2,2-dimethyl-propanamide];

N,N' -(dithiodi-2,1-phenylene)bis[1-methyl-cyclohexanecarboxamide];

N,N' -(dithiodi-2,1-phenylene)bis[1-(3-methylbutyl)-

cyclopentanecarboxamide];

N,N'-(dithiodi-2,1-phenylene)bis[1-(3-methylbutyl)-
cyclohexanecarboxamide];

N,N'-(dithiodi-2,1-phenylene)bis[1-(2-ethylbutyl)-
5 cyclohexanecarboxamide];


N,N'-(dithiodi-2,1-phenylene)bis-tricyclo[3.3.1.1^{3,7}]decane-1-
carboxamide;

propanethioic acid, 2-methyl-, S-[2-[[[1-(2-
ethylbutyl)cyclohexyl]carbonyl]amino]phenyl] ester;

10 propanethioic acid, 2,2-dimethyl-, S-[2-[[[1-(2-
ethylbutyl)cyclohexyl]carbonyl]amino]phenyl] ester; and

ethanethioic acid, S-[2-[[[1-(2-
ethylbutyl)cyclohexyl]carbonyl]amino]phenyl] ester.

Another class of CETP inhibitors that finds utility with the present
15 invention consists of polycyclic aryl and heteroaryl tertiary-heteroalkylamines
having the Formula XIV



Formula XIV

and pharmaceutically acceptable forms thereof, wherein:

n_{XIV} is an integer selected from 0 through 5;

R_{XIV-1} is selected from the group consisting of haloalkyl, haloalkenyl, haloalkoxyalkyl, and haloalkenyloxyalkyl;

X_{XIV} is selected from the group consisting of O, H, F, S, S(O), NH, N(OH), N(alkyl), and N(alkoxy);

R_{XIV-16} is selected from the group consisting of hydrido, alkyl, alkenyl, alkynyl, aryl, aralkyl, aryloxyalkyl, alkoxyalkyl, alkenyloxyalkyl, alkylthioalkyl, arylthioalkyl, aralkoxyalkyl, heteroaralkoxyalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl, cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, monocarboalkoxyalkyl, monocarboalkoxy, dicarboalkoxyalkyl, monocarboxamido, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, acyl, aroyl, heteroaroyl, heteroaryloxyalkyl, dialkoxyposphonoalkyl, trialkylsilyl, and a spacer selected from the group consisting of a covalent single bond and a linear spacer moiety having from 1 through 4 contiguous atoms linked to the point of bonding of an aromatic substituent selected from the group consisting of R_{XIV-4} , R_{XIV-8} , R_{XIV-9} , and R_{XIV-13} to form a heterocycli ring having from 5 through 10 contiguous members with the provisos that said spacer moiety is other than a covalent single bond when R_{XIV-2} is alkyl and there is no R_{XIV-16} wherein X is H or F;

D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} are independently selected from the group consisting of C, N, O, S and a covalent bond with the provisos that no more than one of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} is a covalent bond, no more than one of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} is O, no more than one of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} is S, one of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} must be a covalent bond when two of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} are O and S, and no more than four of D_{XIV-1} , D_{XIV-2} , J_{XIV-1} , J_{XIV-2} and K_{XIV-1} are N;

D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} are independently selected from the group consisting of C, N, O, S and a covalent bond with the provisos that no more than one of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} is a covalent bond, no more than one of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} is O, no more than one of D_{XIV-3} ,

D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} is S, one of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} must be a covalent bond when two of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} are O and S, and no more than four of D_{XIV-3} , D_{XIV-4} , J_{XIV-3} , J_{XIV-4} and K_{XIV-2} and K_{XIV-2} are N;

R_{XIV-2} is independently selected from the group consisting of hydrido,
 5 hydroxy, hydroxyalkyl, amino, aminoalkyl, alkylamino, dialkylamino, alkyl,
 alkenyl, alkynyl, aryl, aralkyl, aralkoxyalkyl, aryloxyalkyl, alkoxyalkyl,
 heteroaryloxyalkyl, alkenyloxyalkyl, alkylthioalkyl, aralkylthioalkyl, arylthioalkyl,
 cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl, cycloalkenylalkyl,
 haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy,
 10 aloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy,
 halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl,
 perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl,
 heteroaralkylthioalkyl, monocarboalkoxyalkyl, dicarboalkoxyalkyl,
 monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl,
 15 alkylsulfonyl, alkylsulfinylalkyl, alkylsulfonylalkyl, haloalkylsulfinyl,
 haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl,
 aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl,
 cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylsulfonylalkyl,
 heteroarylsulfinyl, heteroarylsulfonyl, heteroarylsulfinylalkyl, aralkylsulfinylalkyl,
 20 aralkylsulfonylalkyl, carboxy, carboxyalkyl, carboalkoxy, carboxamide,
 carboxamidoalkyl, carboaralkoxy, dialkoxyphosphono, diaralkoxyphosphono,
 dialkoxyphosphonoalkyl, and diaralkoxyphosphonoalkyl;

R_{XIV-2} and R_{XIV-3} are taken together to form a linear spacer moiety
 25 selected from the group consisting of a covalent single bond and a moiety
 having from 1 through 6 contiguous atoms to form a ring selected from the
 group consisting of a cycloalkyl having from 3 through 8 contiguous members, a
 cycloalkenyl having from 5 through 8 contiguous members, and a heterocyclyl
 having from 4 through 8 contiguous members;

R_{XIV-3} is selected from the group consisting of hydrido, hydroxy, halo,
 30 cyano, aryloxy, hydroxyalkyl, amino, alkylamino, dialkylamino, acyl, sulfhydryl,
 acylamido, alkoxy, alkylthio, arylthio, alkyl, alkenyl, alkynyl, aryl,
 aralkyl, aryloxyalkyl, alkoxyalkyl, heteroarylthio, aralkylthio, aralkoxyalkyl,
 alkylsulfinylalkyl, alkylsulfonylalkyl, aroyl, heteroaroyl, aralkylthioalkyl,
 heteroaralkylthioalkyl, heteroaryloxyalkyl, alkenyloxyalkyl, alkylthioalkyl,
 35 arylthioalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl,

cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl, 5 monocarboalkoxyalkyl, dicarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl, aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl, cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylsulfonylalkyl, 10 heteroarylsulfinyl, heteroarylsulfonyl, heteroarylsulfinylalkyl, aralkylsulfinylalkyl, aralkylsulfonylalkyl, carboxy, carboxyalkyl, carboalkoxy, carboxamide, carboxamidoalkyl, carboaralkoxy, dialkoxyposphono, diaralkoxyposphono, dialkoxyposphonoalkyl, and diaralkoxyposphonoalkyl;

Y_{XIV} is selected from a group consisting of a covalent single 15 bond, $(C(R_{XIV-14})_2)_{q_{XIV}}$ wherein q_{XIV} is an integer selected from 1 and 2 and $(CH(R_{XIV-14}))_{g_{XIV}}-W_{XIV}-(CH(R_{XIV-14}))_{p_{XIV}}$ wherein g_{XIV} and p_{XIV} are integers independently selected from 0 and 1;

R_{XIV-14} is independently selected from the group consisting of hydrido, hydroxy, halo, cyano, aryloxy, amino, alkylamino, dialkylamino, hydroxyalkyl, 20 acyl, aroyl, heteroaroyl, heteroaryloxyalkyl, sulfhydryl, acylamido, alkoxy, alkylthio, arylthio, alkyl, alkenyl, alkynyl, aryl, aralkyl, aryloxyalkyl, aralkoxyalkylalkoxy, alkylsulfinylalkyl, alkylsulfonylalkyl, aralkylthioalkyl, heteroaralkoxythioalkyl, alkoxyalkyl, heteroaryloxyalkyl, alkenyloxyalkyl, alkylthioalkyl, arylthioalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, 25 cycloalkenyl, cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl, heteroaralkylthioalkyl, monocarboalkoxyalkyl, 30 dicarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl, aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl, cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylsulfonylalkyl, heteroarylsulfinyl, 35 heteroarylsulfonyl, heteroarylsulfinylalkyl, aralkylsulfinylalkyl,

aralkylsulfonylalkyl, carboxy, carboxyalkyl, carboalkoxy, carboxamide, carboxamidoalkyl, carboaralkoxy, dialkoxyposphono, diaralkoxyposphono, dialkoxyposphonoalkyl, diaralkoxyposphonoalkyl, a spacer selected from a moiety having a chain length of 3 to 6 atoms connected to the point of bonding selected from the group consisting of R_{XIV-8} and R_{XIV-13} to form a ring selected from the group consisting of a cycloalkenyl ring having from 5 through 8 contiguous members and a heterocyclyl ring having from 5 through 8 contiguous members and a spacer selected from a moiety having a chain length of 2 to 5 atoms connected to the point of bonding selected from the group consisting of R_{XIV-4} and R_{XIV-8} to form a heterocyclyl having from 5 through 8 contiguous members with the proviso that, when Y_{XIV} is a covalent bond, an R_{XIV-14} substituent is not attached to Y_{XIV} ;

R_{XIV-14} and R_{XIV-14} , when bonded to the different atoms, are taken together to form a group selected from the group consisting of a covalent bond, alkylene, haloalkylene, and a spacer selected from a group consisting of a moiety having a chain length of 2 to 5 atoms connected to form a ring selected from the group of a saturated cycloalkyl having from 5 through 8 contiguous members, a cycloalkenyl having from 5 through 8 contiguous members, and a heterocyclyl having from 5 through 8 contiguous members;

R_{XIV-14} and R_{XIV-14} , when bonded to the same atom are taken together to form a group selected from the group consisting of oxo, thiono, alkylene, haloalkylene, and a spacer selected from the group consisting of a moiety having a chain length of 3 to 7 atoms connected to form a ring selected from the group consisting of a cycloalkyl having from 4 through 8 contiguous members, a cycloalkenyl having from 4 through 8 contiguous members, and a heterocyclyl having from 4 through 8 contiguous members;

W_{XIV} is selected from the group consisting of O, C(O), C(S), C(O)N(R_{XIV-14}), C(S)N(R_{XIV-14}), (R_{XIV-14})NC(O), (R_{XIV-14})NC(S), S, S(O), S(O)₂, S(O)₂N(R_{XIV-14}), (R_{XIV-14})NS(O)₂, and N(R_{XIV-14}) with the proviso that R_{XIV-14} is selected from other than halo and cyano;

Z_{XIV} is independently selected from a group consisting of a covalent single bond, $(C(R_{XIV-15})_2)_{q_{XIV-2}}$ wherein q_{XIV-2} is an integer selected from 1 and 2, $(CH(R_{XIV-15}))_{j_{XIV}}-W-(CH(R_{XIV-15}))_{k_{XIV}}$ wherein j_{XIV} and k_{XIV} are integers independently selected from 0 and 1 with the proviso that, when Z_{XIV} is a covalent single bond, an R_{XIV-15} substituent is not attached to Z_{XIV} ;

R_{XIV-15} is independently selected, when Z_{XIV} is $(C(R_{XIV-15})_2)_{q_{XIV}}$ wherein q_{XIV} is an integer selected from 1 and 2, from the group consisting of hydrido, hydroxy, halo, cyano, aryloxy, amino, alkylamino, dialkylamino, hydroxyalkyl, acyl, aroyl, heteroaroyl, heteroaryloxyalkyl, sulfhydryl, acylamido, alkoxy,

5 alkylthio, arylthio, alkyl, alkenyl, alkynyl, aryl, aralkyl, aryloxyalkyl, aralkoxyalkyl, alkylsulfinylalkyl, alkylsulfonylalkyl, aralkylthioalkyl, heteroaralkylthioalkyl, alkoxyalkyl, heteroaryloxyalkyl, alkenyloxyalkyl, alkylthioalkyl, arylthioalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl, cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy,

10 haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl, heteroaralkylthioalkyl, monocarboalkoxyalkyl, dicarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfinyl,

15 haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl, aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl, cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylulfonylalkyl, heteroarylulfonyl,

20 heteroarylulfonyl, heteroarylulfonylalkyl, aralkylsulfinylalkyl, aralkylsulfonylalkyl, carboxy, carboxyalkyl, carboalkoxy, carboxamide, carboxamidoalkyl, carboaralkoxy, dialkoxyphosphono, diaralkoxyphosphono, dialkoxyphosphonoalkyl, diaralkoxyphosphonoalkyl, a spacer selected from a moiety having a chain length of 3 to 6 atoms connected to the point of bonding

25 selected from the group consisting of R_{XIV-4} and R_{XIV-8} to form a ring selected from the group consisting of a cycloalkenyl ring having from 5 through 8 contiguous members and a heterocyclyl ring having from 5 through 8 contiguous members, and a spacer selected from a moiety having a chain length of 2 to 5 atoms connected to the point of bonding selected from the group consisting of R_{XIV-9}

30 and R_{XIV-13} to form a heterocyclyl having from 5 through 8 contiguous members; R_{XIV-15} and R_{XIV-15} , when bonded to the different atoms, are taken together to form a group selected from the group consisting of a covalent bond, alkylene, haloalkylene, and a spacer selected from a group consisting of a moiety having a chain length of 2 to 5 atoms connected to form a ring selected

35 from the group of a saturated cycloalkyl having from 5 through 8 contiguous

members, a cycloalkenyl having from 5 through 8 contiguous members, and a heterocyclyl having from 5 through 8 contiguous members;

R_{XIV-15} and R_{XIV-15} , when bonded to the same atom are taken together to form a group selected from the group consisting of oxo, thiono, alkylene, haloalkylene, and a spacer selected from the group consisting of a moiety having a chain length of 3 to 7 atoms connected to form a ring selected from the group consisting of a cycloalkyl having from 4 through 8 contiguous members, a cycloalkenyl having from 4 through 8 contiguous members, and a heterocyclyl having from 4 through 8 contiguous members;

R_{XIV-15} is independently selected, when Z_{XIV} is $(CH(R_{XIV-15}))_{j_{XIV}}-W-(CH(R_{XIV-15}))_{k_{XIV}}$ wherein j_{XIV} and k_{XIV} are integers independently selected from 0 and 1, from the group consisting of hydrido, halo, cyano, aryloxy, carboxyl, acyl, aroyl, heteroaroyl, hydroxyalkyl, heteroaryloxyalkyl, acylamido, alkoxy, alkylthio, arylthio, alkyl, alkenyl, alkynyl, aryl, aralkyl, aryloxyalkyl, alkoxyalkyl, heteroaryloxyalkyl, aralkoxyalkyl, heteroaralkoxyalkyl, alkylsulfonylalkyl, alkylsulfinylalkyl, alkenyloxyalkyl, alkylthioalkyl, arylthioalkyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkenyl, cycloalkenyl, cycloalkenylalkyl, haloalkyl, haloalkenyl, halocycloalkyl, halocycloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, halocycloalkenyloxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, heteroarylalkyl, heteroarylthioalkyl, heteroaralkylthioalkyl, monocarboalkoxyalkyl, dicarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, alkylsulfinyl, alkylsulfonyl, haloalkylsulfinyl, haloalkylsulfonyl, arylsulfinyl, arylsulfinylalkyl, arylsulfonyl, arylsulfonylalkyl, aralkylsulfinyl, aralkylsulfonyl, cycloalkylsulfinyl, cycloalkylsulfonyl, cycloalkylsulfinylalkyl, cycloalkylsulfonylalkyl, heteroarylulfonylalkyl, heteroarylulfonyl, heteroarylulfonyl, heteroarylulfonylalkyl, aralkylsulfinylalkyl, aralkylsulfonylalkyl, carboxyalkyl, carboalkoxy, carboxamide, carboxamidoalkyl, carboaralkoxy, dialkoxyposphonoalkyl, diaralkoxyposphonoalkyl, a spacer selected from a linear moiety having a chain length of 3 to 6 atoms connected to the point of bonding selected from the group consisting of R_{XIV-4} and R_{XIV-8} to form a ring selected from the group consisting of a cycloalkenyl ring having from 5 through 8 contiguous members and a heterocyclyl ring having from 5 through 8 contiguous members, and a spacer

selected from a linear moiety having a chain length of 2 to 5 atoms connected to the point of bonding selected from the group consisting of R_{XIV-9} and R_{XIV-13} to form a heterocyclyl ring having from 5 through 8 contiguous members;

- R_{XIV-4} , R_{XIV-5} , R_{XIV-6} , R_{XIV-7} , R_{XIV-8} , R_{XIV-9} , R_{XIV-10} , R_{XIV-11} , R_{XIV-12} , and R_{XIV-13}
- 5 are independently selected from the group consisting of perhaloaryloxy, alkanoylalkyl, alkanoylalkoxy, alkanoyloxy, N-aryl-N-alkylamino, heterocyclylalkoxy, heterocyclylthio, hydroxyalkoxy, carboxamidoalkoxy, alkoxycarbonylalkoxy, alkoxycarbonylalkenyloxy, aralkanoylalkoxy, aralkenoyl, N-alkylcarboxamido, N-haloalkylcarboxamido, N-cycloalkylcarboxamido,
- 10 N-arylcarboxamidoalkoxy, cycloalkylcarbonyl, cyanoalkoxy, heterocyclylcarbonyl, hydrido, carboxy, heteroaralkylthio, heteroaralkoxy, cycloalkylamino, acylalkyl, acylalkoxy, aroylalkoxy, heterocycliloxy, aralkylaryl, aralkyl, aralkenyl, aralkynyl, heterocyclyl, perhaloaralkyl, aralkylsulfonyl, aralkylsulfonylalkyl, aralkylsulfinyl, aralkylsulfinylalkyl, halocycloalkyl,
- 15 halocycloalkenyl, cycloalkylsulfinyl, cycloalkylsulfinylalkyl, cycloalkylsulfonyl, cycloalkylsulfonylalkyl, heteroaryl-amino, N-heteroaryl-amino-N-alkylamino, heteroaryl-aminoalkyl, haloalkylthio, alkanoyloxy, alkoxy, alkoxyalkyl, haloalkoxyalkyl, heteroaralkoxy, cycloalkoxy, cycloalkenyloxy, cycloalkoxyalkyl, cycloalkylalkoxy, cycloalkenyloxyalkyl, cycloalkylenedioxy, halocycloalkoxy,
- 20 halocycloalkoxyalkyl, halocycloalkenyloxy, halocycloalkenyloxyalkyl, hydroxy, amino, thio, nitro, lower alkylamino, alkylthio, alkylthioalkyl, arylamino, aralkylamino, arylthio, arylthioalkyl, heteroaralkoxyalkyl, alkylsulfinyl, alkylsulfinylalkyl, arylsulfinylalkyl, arylsulfonylalkyl, heteroaryl-sulfinylalkyl, heteroaryl-sulfonylalkyl, alkylsulfonyl, alkylsulfonylalkyl, haloalkylsulfinylalkyl,
- 25 haloalkylsulfonylalkyl, alkylsulfonamido, alkylaminosulfonyl, amidosulfonyl, monoalkylamid-sulfonyl, dialkyl amid-sulfonyl, monoarylamid-sulfonyl, arylsulfonamido, diarylamidosulfonyl, monoalkyl monoaryl amid-sulfonyl, arylsulfinyl, arylsulfonyl, heteroarylthio, heteroaryl-sulfinyl, heteroaryl-sulfonyl, heterocyclyl-sulfonyl, heterocyclylthio, alkanoyl, alkenoyl, aroyl, heteroaroyl,
- 30 aralkanoyl, heteroaralkanoyl, haloalkanoyl, alkyl, alkenyl, alkynyl, alkenyloxy, alkenyloxyalkyl, alkylenedioxy, haloalkylenedioxy, cycloalkyl, cycloalkylalkanoyl, cycloalkenyl, lower cycloalkylalkyl, lower cycloalkenylalkyl, halo, haloalkyl, haloalkenyl, haloalkoxy, hydroxyhaloalkyl, hydroxyaralkyl, hydroxyalkyl,
- 35 hydroxyheteroaralkyl, haloalkoxyalkyl, aryl, heteroaralkynyl, aryloxy, aralkoxy, aryloxyalkyl, saturated heterocyclyl, partially saturated heterocyclyl, heteroaryl,

heteroaryloxy, heteroaryloxyalkyl, arylalkenyl, heteroarylalkenyl, carboxyalkyl, carboalkoxy, alkoxycarboxamido, alkylamidocarbonylamido, arylamidocarbonylamido, carboalkoxyalkyl, carboalkoxyalkenyl, carboaralkoxy, carboxamido, carboxamidoalkyl, cyano, carbohaloalkoxy, phosphono, 5 phosphonoalkyl, diaralkoxyphosphono, and diaralkoxyphosphonoalkyl with the proviso that there are one to five non-hydrido ring substituents R_{XIV-4} , R_{XIV-5} , R_{XIV-6} , R_{XIV-7} , and R_{XIV-8} present, that there are one to five non-hydrido ring substituents R_{XIV-9} , R_{XIV-10} , R_{XIV-11} , R_{XIV-12} , and R_{XIV-13} present, and R_{XIV-4} , R_{XIV-5} , R_{XIV-6} , R_{XIV-7} , R_{XIV-8} , R_{XIV-9} , R_{XIV-10} , R_{XIV-11} , R_{XIV-12} , and R_{XIV-13} are each 10 independently selected to maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen;

R_{XIV-4} and R_{XIV-5} , R_{XIV-5} and R_{XIV-6} , R_{XIV-6} and R_{XIV-7} , R_{XIV-7} and R_{XIV-8} , R_{XIV-8} and R_{XIV-9} , R_{XIV-9} and R_{XIV-10} , R_{XIV-10} and R_{XIV-11} , R_{XIV-11} and R_{XIV-12} , and R_{XIV-12} and 15 R_{XIV-13} are independently selected to form spacer pairs wherein a spacer pair is taken together to form a linear moiety having from 3 through 6 contiguous atoms connecting the points of bonding of said spacer pair members to form a ring selected from the group consisting of a cycloalkenyl ring having 5 through 8 contiguous members, a partially saturated heterocyclyl ring having 5 through 8 20 contiguous members, a heteroaryl ring having 5 through 6 contiguous members, and an aryl with the provisos that no more than one of the group consisting of spacer pairs R_{XIV-4} and R_{XIV-5} , R_{XIV-5} and R_{XIV-6} , R_{XIV-6} and R_{XIV-7} , and R_{XIV-7} and R_{XIV-8} are used at the same time and that no more than one of the group consisting of spacer pairs R_{XIV-9} and R_{XIV-10} , R_{XIV-10} and R_{XIV-11} , R_{XIV-11} and R_{XIV-12} , 25 and R_{XIV-12} and R_{XIV-13} are used at the same time;

R_{XIV-4} and R_{XIV-9} , R_{XIV-4} and R_{XIV-13} , R_{XIV-8} and R_{XIV-9} , and R_{XIV-8} and R_{XIV-13} are independently selected to form a spacer pair wherein said spacer pair is taken together to form a linear moiety wherein said linear moiety forms a ring 30 selected from the group consisting of a partially saturated heterocyclyl ring having from 5 through 8 contiguous members and a heteroaryl ring having from 5 through 6 contiguous members with the proviso that no more than one of the group consisting of spacer pairs R_{XIV-4} and R_{XIV-9} , R_{XIV-4} and R_{XIV-13} , R_{XIV-8} and R_{XIV-9} , and R_{XIV-8} and R_{XIV-13} is used at the same time.

Compounds of Formula XIV and their methods of manufacture are disclosed in PCT Publication No. WO 00/18721, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula XIV:

- 5 3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-isopropylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[3-(3-cyclopropylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-(2-furyl)phenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-(2,3-dichlorophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-methylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[3-(3,5-dimethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-ethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-*t*-butylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 35

3-[[3-(3-methylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

5 3-[[3-(phenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

10 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethoxy)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethyl)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3,5-dimethylphenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

15 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethylthio)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3,5-difluorophenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

20 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[cyclohexylmethoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

25 3-[[3-(3-difluoromethoxyphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

30 3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(pentafluoroethyl)methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[3-(3-isopropylphenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[3-(3-cyclopropylphenoxy)phenyl][[3-(pentafluoroethyl) phenyl]methyl]-
amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-(2-furyl)phenoxy)phenyl][[3-(pentafluoroethyl) phenyl]methyl]-
amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[3-(2,3-dichlorophenoxy)phenyl][[3-(pentafluoroethyl) phenyl]methyl]-
amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-methylphenoxy)phenyl][[3-(pentafluoroethyl)
10 phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[3-
(pentafluoroethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[3-(pentafluoroethyl)phenyl]-
methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3,5-dimethylphenoxy)phenyl][[3-(pentafluoroethyl) phenyl]methyl]-
20 amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-ethylphenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-t-butylphenoxy)phenyl][[3-(pentafluoroethyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[3-(3-methylphenoxy)phenyl][[3-pentafluoroethyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[3-
(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(phenoxy)phenyl][[3-(pentafluoroethyl)phenyl]methyl]
30 amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[3-
(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(pentafluoroethyl)phenyl]methyl][3-[[3-(trifluoromethoxy)phenyl]-
methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[[3-(trifluoromethyl)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[[3-(trifluoromethylthio)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[[3,5-difluorophenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(pentafluoroethyl)phenyl]methyl][3-[cyclohexylmethoxy]phenyl]-amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[[3-(3-difluoromethoxyphenoxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[3-(pentafluoroethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-isopropylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[[3-(3-cyclopropylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-(2-furyl)phenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(2,3-dichlorophenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[[3-(4-fluorophenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(4-methylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(heptafluoropropyl) phenyl]-
methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-(heptafluoropropyl)
phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[3-
(heptafluoropropyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[3-
(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3,5-dimethylphenoxy)phenyl][[3-(heptafluoropropyl)
phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[3-(3-ethylphenoxy)phenyl][[3-(heptafluoropropyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-*i*-butylphenoxy)phenyl][[3-(heptafluoropropyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-(3-methylphenoxy)phenyl][[3-(heptafluoropropyl)
phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[3-
(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(phenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]
amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[3-[3-(*N,N*-dimethylamino)phenoxy]phenyl][[3-
(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3-
(trifluoromethoxy)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3-(trifluoromethyl)phenyl]-
methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3,5-
dimethylphenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3-
(trifluoromethylthio)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3,5-
difluorophenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(heptafluoropropyl)phenyl]methyl][3-(cyclohexylmethoxy)phenyl]-
amino]-1,1,1-trifluoro-2-propanol;
- 35 3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[3-

- (heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl]][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-difluoromethoxyphenoxy)phenyl]][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 5 3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl]][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl]][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 10 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-cyclopropylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 15 3-[[3-(3-(2-furyl)phenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(2,3-dichlorophenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 20 3-[[3-(4-fluorophenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(4-methylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 25 3-[[3-(2-fluoro-5-bromophenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(4-chloro-3-ethylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 30 3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3,5-dimethylphenoxy)phenyl]][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[3-(3-ethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-t-butylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[3-(3-methylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(phenoxy)phenyl][[2-fluoro-5-(trifluoromethyl) phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethoxy)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethyl)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[cyclohexylmethoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-difluoromethoxyphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-isopropylphenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[3-(3-cyclopropylphenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-(2-furyl)phenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2,3-dichlorophenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[3-(4-fluorophenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl) phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-methylphenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl) phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-(2-fluoro-5-bromophenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-chloro-3-ethylphenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl]][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl]][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3,5-dimethylphenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[3-(3-ethylphenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl) phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-t-butylphenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[3-(3-methylphenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl]][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(phenoxy)phenyl]][[2-fluoro-4-(trifluoromethyl) phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 35

91

3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethoxy)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

5 3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethyl)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

10 3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[cyclohexylmethoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;

15 3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;


3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

20 3-[[3-(3-difluoromethoxyphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol; and

3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol.

25 Another class of CETP inhibitors that finds utility with the present invention consists of substituted N-Aliphatic-N-Aromatic *tertiary*-Heteroalkylamines having the Formula XV




Formula XV


and pharmaceutically acceptable forms thereof, wherein:

5 n_{XV} is an integer selected from 1 through 2;

AQ-1



AQ-2



A_{XV} and Q_{XV} are independently selected from the group consisting of
 $-\text{CH}_2(\text{CR}_{XV-37}\text{R}_{XV-38})_{v_{XV}}-(\text{CR}_{XV-33}\text{R}_{XV-34})_{u_{XV}}-\text{T}_{XV}-(\text{CR}_{XV-35}\text{R}_{XV-36})_{w_{XV}}-\text{H}$,

with the provisos that one of A_{XV} and Q_{XV} must be AQ-1 and that one of A_{XV} and Q_{XV} must be selected from the group consisting of AQ-2 and $-\text{CH}_2(\text{CR}_{XV-37}\text{R}_{XV-38})_{v_{XV}}-(\text{CR}_{XV-33}\text{R}_{XV-34})_{u_{XV}}-\text{T}_{XV}-(\text{CR}_{XV-35}\text{R}_{XV-36})_{w_{XV}}-\text{H}$;

T_{XV} is selected from the group consisting of a single covalent bond, O, S, S(O), S(O)₂, C(R_{XV-33})=C(R_{XV-35}), and

C \equiv C;

v_{XV} is an integer selected from 0 through 1 with the proviso that v_{XV} is 1 when any one of R_{XV-33}, R_{XV-34}, R_{XV-35}, and R_{XV-36} is aryl or heteroaryl;

u_{XV} and w_{XV} are integers independently selected from 0 through 6;

A_{XV-1} is C(R_{XV-30});

10 D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} are independently selected from the group consisting of C, N, O, S and a covalent bond with the provisos that no more than one of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} is a covalent bond, no more than one of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} is O, no more than one of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} is S, one of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} must be

15 a covalent bond when two of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} are O and S, and no more than four of D_{XV-1} , D_{XV-2} , J_{XV-1} , J_{XV-2} , and K_{XV-1} are N;

B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are independently selected from the group consisting of C, C(R_{XV-30}), N, O, S and a covalent bond with the provisos that no more than 5 of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are

20 a covalent bond, no more than two of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are O, no more than two of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are S, no more than two of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are simultaneously O and S, and no more than two of B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are N;

25 B_{XV-1} and D_{XV-3} , D_{XV-3} and J_{XV-3} , J_{XV-3} and K_{XV-2} , K_{XV-2} and J_{XV-4} , J_{XV-4} and D_{XV-4} , and D_{XV-4} and B_{XV-2} are independently selected to form an in-ring spacer pair wherein said

spacer pair is selected from the group consisting of C(R_{XV-33})=C(R_{XV-35}) and N=N with the provisos that AQ-2 must be a ring of at least five contiguous members,

30 that no more than two of the group of said spacer pairs are simultaneously C(R_{XV-33})=C(R_{XV-35}) and that no more than one of the group of said spacer pairs can be N=N unless the other spacer pairs are other than C(R_{XV-33})=C(R_{XV-35}), O, N, and S;

R_{XV-1} is selected from the group consisting of haloalkyl and haloalkoxymethyl;

R_{XV-2} is selected from the group consisting of hydrido, aryl, alkyl, alkenyl, haloalkyl, haloalkoxy, haloalkoxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl and heteroaryl;

R_{XV-3} is selected from the group consisting of hydrido, aryl, alkyl, alkenyl, haloalkyl, and haloalkoxyalkyl;

Y_{XV} is selected from the group consisting of a covalent single bond, $(CH_2)_q$ wherein q is an integer selected from 1 through 2 and $(CH_2)_j-O-(CH_2)_k$ wherein j and k are integers independently selected from 0 through 1;

Z_{XV} is selected from the group consisting of covalent single bond, $(CH_2)_q$ wherein q is an integer selected from 1 through 2, and $(CH_2)_j-O-(CH_2)_k$ wherein j and k are integers independently selected from 0 through 1;

R_{XV-4} , R_{XV-8} , R_{XV-9} and R_{XV-13} are independently selected from the group consisting of hydrido, halo, haloalkyl, and alkyl;

R_{XV-30} is selected from the group consisting of hydrido, alkoxy, alkoxyalkyl, halo, haloalkyl, alkylamino, alkylthio, alkylthioalkyl, alkyl, alkenyl, haloalkoxy, and haloalkoxyalkyl with the proviso that R_{XV-30} is selected to maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen;

R_{XV-30} , when bonded to A_{XV-1} , is taken together to form an intra-ring linear spacer connecting the A_{XV-1} -carbon at the point of attachment of R_{XV-30} to the point of bonding of a group selected from the group consisting of R_{XV-10} , R_{XV-11} , R_{XV-12} , R_{XV-31} , and R_{XV-32} wherein said intra-ring linear spacer is selected from the group consisting of a covalent single bond and a spacer moiety having from 1 through 6 contiguous atoms to form a ring selected from the group consisting of a cycloalkyl having from 3 through 10 contiguous members, a cycloalkenyl having from 5 through 10 contiguous members, and a heterocyclyl having from 5 through 10 contiguous members;

R_{XV-30} , when bonded to A_{XV-1} , is taken together to form an intra-ring branched spacer connecting the A_{XV-1} -carbon at the point of attachment of R_{XV-30} to the points of bonding of each member of any one of substituent pairs selected from the group consisting of substituent pairs R_{XV-10} and R_{XV-11} , R_{XV-10} and R_{XV-31} , R_{XV-10} and R_{XV-32} , R_{XV-10} and R_{XV-12} , R_{XV-11} and R_{XV-31} , R_{XV-11} and R_{XV-32} , R_{XV-11} and R_{XV-12} , R_{XV-31} and R_{XV-32} , R_{XV-31} and R_{XV-12} , and R_{XV-32} and R_{XV-12} and

wherein said intra-ring branched spacer is selected to form two rings selected from the group consisting of cycloalkyl having from 3 through 10 contiguous members, cycloalkenyl having from 5 through 10 contiguous members, and heterocyclyl having from 5 through 10 contiguous members;

- 5 R_{XV-4} , R_{XV-5} , R_{XV-6} , R_{XV-7} , R_{XV-8} , R_{XV-9} , R_{XV-10} , R_{XV-11} , R_{XV-12} , R_{XV-13} , R_{XV-31} ,
 R_{XV-32} , R_{XV-33} , R_{XV-34} , R_{XV-35} , and R_{XV-36} are independently selected from the
group consisting of hydrido, carboxy, heteroaralkylthio, heteroaralkoxy,
cycloalkylamino, acylalkyl, acylalkoxy, aroylalkoxy, heterocyclyloxy, aralkylaryl,
aralkyl, aralkenyl, aralkynyl, heterocyclyl, perhaloaralkyl, aralkylsulfonyl,
10 aralkylsulfonylalkyl, aralkylsulfinyl, aralkylsulfinylalkyl, halocycloalkyl,
halocycloalkenyl, cycloalkylsulfinyl, cycloalkylsulfinylalkyl, cycloalkylsulfonyl,
cycloalkylsulfonylalkyl, heteroaryl amino, N-heteroaryl amino-N-alkyl amino,
heteroaryl aminoalkyl, haloalkylthio, alkanoyloxy, alkoxy, alkoxyalkyl,
haloalkoxyalkyl, heteroaralkoxy, cycloalkoxy, cycloalkenyloxy, cycloalkoxyalkyl,
15 cycloalkylalkoxy,
cycloalkenyloxyalkyl, cycloalkylenedioxy, halocycloalkoxy,
halocycloalkoxyalkyl, halocycloalkenyloxy, halocycloalkenyloxyalkyl, hydroxy,
amino, thio, nitro, lower alkyl amino, alkylthio, alkylthioalkyl, aryl amino,
aralkyl amino, arylthio, arylthioalkyl, heteroaralkoxyalkyl, alkylsulfinyl,
20 alkylsulfinylalkyl, arylsulfinylalkyl, arylsulfonylalkyl, heteroaryl sulfinylalkyl,
heteroaryl sulfonylalkyl, alkylsulfonyl, alkylsulfonylalkyl, haloalkylsulfinylalkyl,
haloalkylsulfonylalkyl, alkylsulfonamido, alkylaminosulfonyl, amidosulfonyl,
monoalkylamidossulfonyl, dialkyl amidossulfonyl, monoarylamidossulfonyl,
aryl sulfonamido, diarylamidossulfonyl, monoalkyl monoaryl amidossulfonyl,
25 arylsulfinyl, arylsulfonyl, heteroarylthio, heteroaryl sulfinyl, heteroaryl sulfonyl,
heterocyclisulfonyl, heterocyclylthio, alkanoyl, alkenoyl, aroyl, heteroaroyl,
aralkanoyl, heteroaralkanoyl, haloalkanoyl, alkyl, alkenyl, alkynyl, alkenyloxy,
alkenyloxyalkyl, alkylenedioxy, haloalkylenedioxy, cycloalkyl, cycloalkylalkanoyl,
cycloalkenyl, lower cycloalkylalkyl, lower cycloalkenylalkyl, halo, haloalkyl,
30 haloalkenyl, haloalkoxy, hydroxyhaloalkyl,
hydroxyaralkyl, hydroxyalkyl, hydroxyheteroaralkyl, haloalkoxyalkyl, aryl,
heteroaralkynyl, aryloxy, aralkoxy, aryloxyalkyl, saturated heterocyclyl, partially
saturated heterocyclyl, heteroaryl, heteroaryl oxy, heteroaryl oxyalkyl, arylalkenyl,
heteroarylalkenyl, carboxyalkyl, carboalkoxy, alkoxycarboxamido,
35 alkylamidocarbonylamido, alkylamidocarbonylamido, carboalkoxyalkyl,

carboalkoxyalkenyl, carboaralkoxy, carboxamido, carboxamidoalkyl, cyano, carbohaloalkoxy, phosphono, phosphonoalkyl, diaralkoxyphosphono, and diaralkoxyphosphonoalkyl with the provisos that R_{XV-4} , R_{XV-5} , R_{XV-6} , R_{XV-7} , R_{XV-8} , R_{XV-9} , R_{XV-10} , R_{XV-11} , R_{XV-12} , R_{XV-13} , R_{XV-31} , R_{XV-32} , R_{XV-33} , R_{XV-34} , R_{XV-35} , and R_{XV-36} are each independently selected to maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen, that no more than three of the R_{XV-33} and R_{XV-34} substituents are simultaneously selected from other than the group consisting of hydrido and halo, and that no more than three of the R_{XV-35} and R_{XV-36} substituents are simultaneously selected from other than the group consisting of hydrido and halo;

R_{XV-9} , R_{XV-10} , R_{XV-11} , R_{XV-12} , R_{XV-13} , R_{XV-31} , and R_{XV-32} are independently selected to be oxo with the provisos that B_{XV-1} , B_{XV-2} , D_{XV-3} , D_{XV-4} , J_{XV-3} , J_{XV-4} , and K_{XV-2} are independently selected from the group consisting of C and S, no more than two of R_{XV-9} , R_{XV-10} , R_{XV-11} , R_{XV-12} , R_{XV-13} , R_{XV-31} , and R_{XV-32} are simultaneously oxo, and that R_{XV-9} , R_{XV-10} , R_{XV-11} , R_{XV-12} , R_{XV-13} , R_{XV-31} , and R_{XV-32} are each independently selected to maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen;

R_{XV-4} and R_{XV-5} , R_{XV-5} and R_{XV-6} , R_{XV-6} and R_{XV-7} , R_{XV-7} and R_{XV-8} , R_{XV-9} and R_{XV-10} , R_{XV-10} and R_{XV-11} , R_{XV-11} and R_{XV-31} , R_{XV-31} and R_{XV-32} , R_{XV-32} and R_{XV-12} , and R_{XV-12} and R_{XV-13} are independently selected to form spacer pairs wherein a spacer pair is taken together to form a linear moiety having from 3 through 6 contiguous atoms connecting the points of bonding of said spacer pair members to form a ring selected from the group consisting of a cycloalkenyl ring having 5 through 8 contiguous members, a partially saturated heterocyclyl ring having 5 through 8 contiguous members, a heteroaryl ring having 5 through 6 contiguous members, and an aryl with the provisos that no more than one of the group consisting of spacer pairs R_{XV-4} and R_{XV-5} , R_{XV-5} and R_{XV-6} , R_{XV-6} and R_{XV-7} , R_{XV-7} and R_{XV-8} is used at the same time and that no more than one of the group consisting of spacer pairs R_{XV-9} and R_{XV-10} , R_{XV-10} and R_{XV-11} , R_{XV-11} and R_{XV-31} , R_{XV-31} and R_{XV-32} , R_{XV-32} and R_{XV-12} , and R_{XV-12} and R_{XV-13} are used at the same time;

R_{XV-9} and R_{XV-11} , R_{XV-9} and R_{XV-12} , R_{XV-9} and R_{XV-13} , R_{XV-9} and R_{XV-31} , R_{XV-9} and R_{XV-32} , R_{XV-10} and R_{XV-12} , R_{XV-10} and R_{XV-13} , R_{XV-10} and R_{XV-31} , R_{XV-10} and R_{XV-32} , R_{XV-11} and R_{XV-12} , R_{XV-11} and R_{XV-13} , R_{XV-11} and R_{XV-31} , R_{XV-11} and R_{XV-32} , R_{XV-12} and R_{XV-13} , R_{XV-12} and R_{XV-31} , R_{XV-12} and R_{XV-32} , R_{XV-13} and R_{XV-31} , R_{XV-13} and R_{XV-32} , R_{XV-31} and R_{XV-32} are used at the same time;

32, R_{XV-11} and R_{XV-12}, R_{XV-11} and R_{XV-13}, R_{XV-11} and R_{XV-32}, R_{XV-12} and R_{XV-31}, R_{XV-13} and R_{XV-31}, and R_{XV-13} and R_{XV-32} are independently selected to form a spacer pair wherein said spacer pair is taken together to form a linear spacer moiety selected from the group consisting of a covalent single bond and a moiety
 5 having from 1 through 3 contiguous atoms to form a ring selected from the group consisting of a cycloalkyl having from 3 through 8 contiguous members, a cycloalkenyl having from 5 through 8 contiguous members, a saturated heterocyclyl having from 5 through 8 contiguous members and a partially saturated heterocyclyl having from 5 through 8 contiguous members with the
 10 provisos that no more than one of said group of spacer pairs is used at the same time;

R_{XV-37} and R_{XV-38} are independently selected from the group consisting of hydrido, alkoxy, alkoxyalkyl, hydroxy, amino, thio, halo, haloalkyl, alkylamino, alkylthio, alkylthioalkyl, cyano, alkyl, alkenyl, haloalkoxy, and
 15 haloalkoxyalkyl.

Compounds of Formula XV and their methods of manufacture are disclosed in PCT Publication No. WO 00/18723, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the
 20 following compounds of Formula XV:

3-[[3-(4-chloro-3-ethylphenoxy)phenyl]
 (cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(4-chloro-3-ethylphenoxy)phenyl]
 (cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 25 3-[[3-(4-chloro-3-ethylphenoxy)phenyl]
 (cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][(3-trifluoromethyl)cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][(3-pentafluoroethyl)
 30 cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][(3-trifluoromethoxy)
 cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(4-chloro-3-ethylphenoxy)phenyl][(3-(1,1,2,2-
 tetrafluoroethoxy)cyclo-hexylmethyl]amino]-1,1,1-trifluoro-2-propanol;
 35 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]

- (cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]
 (cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]
 5 (cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl][(3-trifluoromethyl)cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]](3-pentafluoroethyl)cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 10 3-[[3-(3-trifluoromethoxyphenoxy)phenyl][(3-
 trifluoromethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-trifluoromethoxyphenoxy)phenyl]](3-(1,1,2,2-
 tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl](cyclohexylmethyl)amino]-1,1,1-
 15 trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl](cyclopentylmethyl)amino]-1,1,1-
 trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl](cyclopropylmethyl)amino]-1,1,1-
 trifluoro-2-propanol;
 20 3-[[3-(3-isopropylphenoxy)phenyl][(3-trifluoromethyl) cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl][(3-pentafluoroethyl) cyclohexyl-
 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl][(3-trifluoromethoxy) cyclohexyl-
 25 methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(3-isopropylphenoxy)phenyl][3-(1,1,2,2-
 tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(2,3-dichlorophenoxy)phenyl](cyclohexylmethyl)amino]-1,1,1-
 trifluoro-2-propanol;
 30 3-[[3-(2,3-dichlorophenoxy)phenyl](cyclopentylmethyl)
 amino]-1,1,1-trifluoro-2-propanol;
 3-[[3-(2,3-dichlorophenoxy)phenyl](cyclopropylmethy)amino]-1,1,1-
 trifluoro-2-propanol;
 3-[[3-(2,3-dichlorophenoxy)phenyl][(3-trifluoromethyl)
 35 cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;

- 3-[[3-(2,3-dichlorophenoxy)phenyl][(3-pentafluoroethyl) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(2,3-dichlorophenoxy)phenyl][(3-trifluoromethoxy) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[3-(2,3-dichlorophenoxy)phenyl][(3-(1,1,2,2-tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl](cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl](cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[3-(4-fluorophenoxy)phenyl](cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][(3-trifluoromethyl) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[3-(4-fluorophenoxy)phenyl][(3-pentafluoroethyl) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][(3-trifluoromethoxy) cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(4-fluorophenoxy)phenyl][(3-(1,1,2,2-tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl](cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl](cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl](cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl][(3-trifluoromethyl)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl][(3-pentafluoroethyl)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl][(3-trifluoromethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[3-(3-trifluoromethoxybenzyloxy)phenyl][(3-(1,1,2,2-tetrafluoroethoxy)-cyclohexylmethyl]amino]-1,1,1-trifluoro-2-propanol;
- 35 3-[[3-(3-trifluoromethylbenzyloxy)phenyl]

- (cyclohexylmethyl)amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(3-trifluoromethylbenzyloxy)phenyl]
(cyclopentylmethyl)amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(3-trifluoromethylbenzyloxy)phenyl]
5 (cyclopropylmethyl)amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(3-trifluoromethylbenzyloxy)phenyl][(3-trifluoromethyl)cyclohexyl-
methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(3-trifluoromethylbenzyloxy)phenyl][(3-pentafluoroethyl)cyclohexyl-
methyl]amino]-1,1,1-trifluoro-2-propanol;
10 3-[[3-(3-trifluoromethylbenzyloxy)phenyl][(3-
trifluoromethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[3-(3-trifluoromethylbenzyloxy)phenyl][3-(1,1,2,2-
tetrafluoroethoxy)cyclohexyl-methyl]amino]-1,1,1-trifluoro-2-propanol;
3-[[[(3-trifluoromethyl)phenyl]methyl](cyclohexyl)amino]-1,1,1-trifluoro-2-
15 propanol;
3-[[[(3-pentafluoroethyl)phenyl]methyl](cyclohexyl)amino]-1,1,1-trifluoro-
2-propanol;
3-[[[(3-trifluoromethoxy)phenyl]methyl](cyclohexyl)amino]-1,1,1-trifluoro-
2-propanol;
20 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]
methyl](cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
3-[[[(3-trifluoromethyl)phenyl]methyl]
(4-methylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
3-[[[(3-pentafluoroethyl)phenyl]methyl]
25 (4-methylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
3-[[[(3-trifluoromethoxy)phenyl]methyl]
(4-methylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](4-
methylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
30 3-[[[(3-trifluoromethyl)phenyl]methyl](3-
trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
3-[[[(3-pentafluoroethyl)phenyl]methyl](3-
trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
3-[[[(3-trifluoromethoxy)phenyl]methyl](3-
35 trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;

- 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](3-trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)cyclohexyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 3-[[[3-(3-pentafluoroethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)cyclohexyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethoxy)phenyl]methyl][3-(4-chloro-3-methylphenoxy)cyclohexyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)-cyclohexyl]amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethyl)phenyl]methyl](3-phenoxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-pentafluoroethyl)phenyl]methyl](3-phenoxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 15 3-[[[3-(3-trifluoromethoxy)phenyl]methyl](3-phenoxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](3-phenoxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethyl)phenyl]methyl](3-isopropoxycyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 20 3-[[[3-(3-pentafluoroethyl)phenyl]methyl](3-isopropoxycyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethoxy)phenyl]methyl](3-isopropoxycyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 25 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](3-isopropoxycyclohexyl)-amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethyl)phenyl]methyl](3-cyclopentyloxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 30 3-[[[3-(3-pentafluoroethyl)phenyl]methyl](3-cyclopentyloxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(3-trifluoromethoxy)phenyl]methyl](3-cyclopentyloxy-cyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
- 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl](3-cyclopentyloxy-cyclohexyl)-amino]-1,1,1-trifluoro-2-propanol;
- 35 3-[[[3-(2-trifluoromethyl)pyrid-6-yl]methyl](3-

- isopropoxycyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-cyclopentyloxycyclohexyl)-
 amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-phenoxy-cyclohexyl)amino]-
 5 1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-
 trifluoromethylcyclohexyl)amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl][3-(4-chloro-3-
 ethylphenoxy)cyclo-hexyl]amino]-1,1,1-trifluoro-2-propanol;
 10 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl][3-(1,1,2,2-
 tetrafluoroethoxy)cyclo-hexyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-pentafluoroethylcyclohexyl)-
 amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(2-trifluoromethyl)pyrid-6-yl]methyl](3-trifluoromethoxycyclohexyl)-
 15 amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)propyl]-
 amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-pentafluoroethyl)phenyl]methyl][3-(4-chloro-3-
 ethylphenoxy)propyl]-amino]-1,1,1-trifluoro-2-propanol;
 20 3-[[[(3-trifluoromethoxy)phenyl]methyl][3-(4-chloro-3-
 ethylphenoxy)propyl]-amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-(4-chloro-3-
 ethylphenoxy)-propyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)-2,2,-di-
 25 fluoropropyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-pentafluoroethyl)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)-2,2,-
 di-fluoropropyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethoxy)phenyl]methyl][3-(4-chloro-3-ethylphenoxy)-2,2,-
 di-fluoropropyl]amino]-1,1,1-trifluoro-2-propanol;
 30 3-[[[(3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-(4-chloro-3-
 ethylphenoxy)-2,2,-difluoropropyl]amino]-1,1,1-trifluoro-2-propanol;
 3-[[[(3-trifluoromethyl)phenyl]methyl][3-(isopropoxy)propyl]amino]-1,1,1-
 trifluoro-2-propanol;
 3-[[[(3-pentafluoroethyl)phenyl]methyl][3-(isopropoxy)propyl]amino]-
 35 1,1,1-trifluoro-2-propanol;

103


3-[[[(3-trifluoromethoxy)phenyl]methyl][3-(isopropoxy)propyl]amino]-
1,1,1-trifluoro-2-propanol;

3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-
(isopropoxy)propyl]amino]-1,1,1-trifluoro-2-propanol; and

5 3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-
(phenoxy)propyl]amino]-1,1,1-trifluoro-2-propanol.

Another class of CETP inhibitors that finds utility with the present
invention consists of (R)-chiral halogenated 1-substituted amino-(n+1)-alkanols
having the Formula XVI

10




Formula XVI

and pharmaceutically acceptable forms thereof, wherein:


15 n_{xvi} is an integer selected from 1 through 4;

X_{xvi} is oxy;

R_{xvi-1} is selected from the group consisting of haloalkyl, haloalkenyl,
haloalkoxymethyl, and haloalkenyloxymethyl with the proviso that R_{xvi-1} has a
higher Cahn-Ingold-Prelog stereochemical system ranking than both R_{xvi-2} and
20 $(CH_{xvi-3})_n-N(A_{xvi})Q_{xvi}$ wherein A_{xvi} is Formula XVI-(II) and Q is Formula XVI-
(III);



XVI-II



XVI-III

R_{XVI-16} is selected from the group consisting of hydrido, alkyl, acyl, aroyl, heteroaroyl, trialkylsilyl, and a spacer selected from the group consisting of a
 5 covalent single bond and a linear spacer moiety having a chain length of 1 to 4 atoms linked to the point of bonding of any aromatic substituent selected from the group consisting of R_{XVI-4} , R_{XVI-8} , R_{XVI-9} , and R_{XVI-13} to form a heterocyclyl ring having from 5 through 10 contiguous members;

D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} are independently selected from the
 10 group consisting of C, N, O, S and covalent bond with the provisos that no more than one of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} is a covalent bond, no more than one D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} is be O, no more than one of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} is S, one of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} must be a covalent bond when two of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} are O and S, and
 15 no more than four of D_{XVI-1} , D_{XVI-2} , J_{XVI-1} , J_{XVI-2} and K_{XVI-1} is N;

D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} are independently selected from the group consisting of C, N, O, S and covalent bond with the provisos that no more than one is a covalent bond, no more than one of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} is O, no more than one of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} is S, no more
 20 than two of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} is O and S, one of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} must be a covalent bond when two of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} are O and S, and no more than four of D_{XVI-3} , D_{XVI-4} , J_{XVI-3} , J_{XVI-4} and K_{XVI-2} are N;

R_{XVI-2} is selected from the group consisting of hydrido, aryl, aralkyl, alkyl, alkenyl, alkenyloxyalkyl, haloalkyl, haloalkenyl, halocycloalkyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, halocycloalkoxy, halocycloalkoxyalkyl, perhaloaryl, perhaloaralkyl, perhaloaryloxyalkyl, heteroaryl, dicyanoalkyl, and carboalkoxycyanoalkyl, with the proviso that R_{XVI-2} has a lower Cahn-Ingold-Prelog system ranking than both R_{XVI-1} and $(CHR_{XVI-3})_n-N(A_{XVI})Q_{XVI}$;

R_{XVI-3} is selected from the group consisting of hydrido, hydroxy, cyano, aryl, aralkyl, acyl, alkoxy, alkyl, alkenyl, alkoxyalkyl, heteroaryl, alkenyloxyalkyl, haloalkyl, haloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, monocyanoalkyl, dicyanoalkyl, carboxamide, and carboxamidoalkyl, with the provisos that $(CHR_{XVI-3})_n-N(A_{XVI})Q_{XVI}$ has a lower Cahn-Ingold-Prelog stereochemical system ranking than R_{XVI-1} and a higher Cahn-Ingold-Prelog stereochemical system ranking than R_{XVI-2} ;

Y_{XVI} is selected from a group consisting of a covalent single bond, $(C(R_{XVI-14})_2)_q$ wherein q is an integer selected from 1 and 2 and $(CH(R_{XVI-14}))_g-W_{XVI}-(CH(R_{XVI-14}))_p$ wherein g and p are integers independently selected from 0 and 1;

R_{XVI-14} is selected from the group consisting of hydrido, hydroxy, cyano, hydroxyalkyl, acyl, alkoxy, alkyl, alkenyl, alkynyl, alkoxyalkyl, haloalkyl, haloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, monocarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, carboalkoxy, carboxamide, and carboxamidoalkyl;

Z_{XVI} is selected from a group consisting of a covalent single bond, $(C(R_{XVI-15})_2)_q$, wherein q is an integer selected from 1 and 2, and $(CH(R_{XVI-15}))_j-W_{XVI}-(CH(R_{XVI-15}))_k$ wherein j and k are integers independently selected from 0 and 1;

X_{XVI} is selected from the group consisting of O, C(O), C(S), C(O)N(R_{XVI-14}), C(S)N(R_{XVI-14}), (R_{XVI-14})NC(O), (R_{XVI-14})NC(S), S, S(O), S(O)₂, S(O)₂N(R_{XVI-14}), (R_{XVI-14})NS(O)₂, and N(R_{XVI-14}) with the proviso that R_{XVI-14} is other than cyano;

R_{XVI-15} is selected, from the group consisting of hydrido, cyano, hydroxyalkyl, acyl, alkoxy, alkyl, alkenyl, alkynyl, alkoxyalkyl, haloalkyl, haloalkenyl, haloalkoxy, haloalkoxyalkyl, haloalkenyloxyalkyl, monocarboalkoxyalkyl, monocyanoalkyl, dicyanoalkyl, carboalkoxycyanoalkyl, carboalkoxy, carboxamide, and carboxamidoalkyl;

R_{XVI-4} , R_{XVI-5} , R_{XVI-6} , R_{XVI-7} , R_{XVI-8} , R_{XVI-9} , R_{XVI-10} , R_{XVI-11} , R_{XVI-12} , and R_{XVI-13}
 are independently selected from the group consisting of hydrido, carboxy,
 heteroaralkylthio, heteroaralkoxy, cycloalkylamino, acylalkyl, acylalkoxy,
 aroylalkoxy, heterocycloxy, aralkylaryl, aralkyl, aralkenyl, aralkynyl,
 5 heterocyclyl, perhaloaralkyl, aralkylsulfonyl, aralkylsulfonylalkyl, aralkylsulfinyl,
 aralkylsulfinylalkyl, halocycloalkyl, halocycloalkenyl, cycloalkylsulfinyl,
 cycloalkylsulfinylalkyl, cycloalkylsulfonyl, cycloalkylsulfonylalkyl,
 heteroaryl amino, N-heteroaryl amino-N-alkyl amino, heteroaralkyl,
 heteroaryl aminoalkyl, haloalkylthio, alkanoyloxy, alkoxy, alkoxyalkyl,
 10 haloalkoxylalkyl, heteroaralkoxy, cycloalkoxy, cycloalkenyl oxy, cycloalkoxyalkyl,
 cycloalkylalkoxy, cycloalkenyl oxyalkyl, cycloalkylenedioxy, halocycloalkoxy,
 halocycloalkoxyalkyl, halocycloalkenyl oxy, halocycloalkenyl oxyalkyl, hydroxy,
 amino, thio, nitro, lower alkyl amino, alkylthio, alkylthioalkyl, aryl amino,
 aralkyl amino, arylthio, arylthioalkyl, heteroaralkoxyalkyl, alkylsulfinyl,
 15 alkylsulfinylalkyl, arylsulfinylalkyl, arylsulfonylalkyl, heteroaryl sulfinylalkyl,
 heteroaryl sulfonylalkyl, alkylsulfonyl, alkylsulfonylalkyl, haloalkylsulfinylalkyl,
 haloalkylsulfonylalkyl, alkylsulfonamido, alkylaminosulfonyl, amidosulfonyl,
 monoalkyl amidosulfonyl, dialkyl, amidosulfonyl, monoaryl amidosulfonyl,
 arylsulfonamido, diarylamidosulfonyl, monoalkyl monoaryl amidosulfonyl,
 20 arylsulfinyl, arylsulfonyl, heteroarylthio, heteroaryl sulfinyl, heteroaryl sulfonyl,
 heterocyclylsulfonyl, heterocyclylthio, alkanoyl, alkenoyl, aroyl, heteroaroyl,
 aralkanoyl, heteroaralkanoyl, haloalkanoyl, alkyl, alkenyl, alkynyl, alkenyloxy,
 alkenyloxyalkyl, alkylenedioxy, haloalkylenedioxy, cycloalkyl, cycloalkylalkanoyl,
 cycloalkenyl, lower cycloalkylalkyl, lower cycloalkenylalkyl, halo, haloalkyl,
 25 haloalkenyl, haloalkoxy, hydroxyhaloalkyl,
 hydroxyaralkyl, hydroxyalkyl, hydroxyheteroaralkyl, haloalkoxyalkyl, aryl,
 heteroaralkynyl, aryloxy, aralkoxy, aryloxyalkyl, saturated heterocyclyl, partially
 saturated heterocyclyl, heteroaryl, heteroaryl oxy, heteroaryl oxyalkyl, arylalkenyl,
 heteroarylalkenyl, carboxyalkyl, carboalkoxy, alkoxycarboxamido,
 30 alkylamidocarbonylamido, arylamidocarbonylamido, carboalkoxyalkyl,
 carboalkoxyalkenyl, carboaralkoxy, carboxamido, carboxamidoalkyl, cyano,
 carbohaloalkoxy, phosphono, phosphonoalkyl, diaralkoxyphosphono, and
 diaralkoxyphosphonoalkyl with the proviso that R_{XVI-4} , R_{XVI-5} , R_{XVI-6} , R_{XVI-7} , R_{XVI-8} ,
 R_{XVI-9} , R_{XVI-10} , R_{XVI-11} , R_{XVI-12} , and R_{XVI-13} are each independently selected to

maintain the tetravalent nature of carbon, trivalent nature of nitrogen, the divalent nature of sulfur, and the divalent nature of oxygen;

R_{XVI-4} and R_{XVI-5} , R_{XVI-5} and R_{XVI-6} , R_{XVI-6} and R_{XVI-7} , R_{XVI-7} and R_{XVI-8} , R_{XVI-9} and R_{XVI-10} , R_{XVI-10} and R_{XVI-11} , R_{XVI-11} and R_{XVI-12} , and R_{XVI-12} and R_{XIV-13} are
 5 independently selected to form spacer pairs wherein a spacer pair is taken together to form a linear moiety having from 3 through 6 contiguous atoms connecting the points of bonding of said spacer pair members to form a ring selected from the group consisting of a cycloalkenyl ring having 5 through 8 contiguous members, a partially saturated heterocyclyl ring having 5 through 8
 10 contiguous members, a heteroaryl ring having 5 through 6 contiguous members, and an aryl with the provisos that no more than one of the group consisting of spacer pairs R_{XVI-4} and R_{XVI-5} , R_{XVI-5} and R_{XVI-6} , R_{XVI-6} and R_{XVI-7} , and R_{XVI-7} and R_{XVI-8} is used at the same time and that no more than one of the group consisting of spacer pairs R_{XIV-9} and R_{XVI-10} , R_{XVI-10} and R_{XVI-11} , R_{XVI-11} and R_{XVI-12} ,
 15 and R_{XVI-12} and R_{XVI-13} can be used at the same time;

R_{XVI-4} and R_{XVI-9} , R_{XVI-4} and R_{XVI-13} , R_{XVI-8} and R_{XVI-9} , and R_{XVI-8} and R_{XVI-13} is independently selected to form a spacer pair wherein said spacer pair is taken together to form a linear moiety wherein said linear moiety forms a ring selected from the group consisting of a partially saturated heterocyclyl ring
 20 having from 5 through 8 contiguous members and a heteroaryl ring having from 5 through 6 contiguous members with the proviso that no more than one of the group consisting of spacer pairs R_{XVI-4} and R_{XVI-9} , R_{XVI-4} and R_{XVI-13} , R_{XVI-8} and R_{XVI-9} , and R_{XVI-8} and R_{XVI-13} is used at the same time.

Compounds of Formula XVI and their methods of manufacture are
 25 disclosed in PCT Publication No. WO 00/18724, which is incorporated herein by reference in its entirety for all purposes.

In a preferred embodiment, the CETP inhibitor is selected from the following compounds of Formula XVI:

(2R)-3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(1,1,2,2-
 30 tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(3-isopropylphenoxy)phenyl][[3-(1,1,2,2-
 tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(3-cyclopropylphenoxy)phenyl][[3-(1,1,2,2-
 tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;

- (2R)-3-[[3-(3-(2-furyl)phenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2,3-dichlorophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[3-(4-fluorophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(4-methylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 (2R)-3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3,5-dimethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-ethylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 (2R)-3-[[3-(3-t-butylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-methylphenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 (2R)-3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(phenoxy)phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(N,N-dimethylamino)phenoxy]phenyl][[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethoxy)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethyl)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;

- (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3-(trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(1,1,2,2-tetrafluoroethoxy)phenyl]methyl][3-(cyclohexylmethoxy)-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 (2R)-3-[[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(3-difluoromethoxyphenoxy)phenyl][3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][3-(1,1,2,2-tetrafluoroethoxy)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(3-trifluoromethoxyphenoxy)phenyl][3-(pentafluoroethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 (2R)-3-[[[3-(3-isopropylphenoxy)phenyl][3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(3-cyclopropylphenoxy)phenyl][3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 25 (2R)-3-[[[3-(3-(2-furyl)phenoxy)phenyl][3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(2,3-dichlorophenoxy)phenyl][3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(4-fluorophenoxy)phenyl][3-(pentafluoroethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 (2R)-3-[[[3-(4-methylphenoxy)phenyl][3-(pentafluoroethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(2-fluoro-5-bromophenoxy)phenyl][3-(pentafluoroethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 35 (2R)-3-[[[3-(4-chloro-3-ethylphenoxy)phenyl][3-

- (pentafluoroethyl)phenyl)methyl]-amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][3-(pentafluoroethyl)-phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 5 (2R)-3-[[3-(3,5-dimethylphenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]-amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(3-ethylphenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 10 (2R)-3-[[3-(3-*t*-butylphenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(3-methylphenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 15 (2R)-3-[[3-(phenoxy)phenyl][3-(pentafluoroethyl)phenyl)methyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(*N,N*-dimethylamino)phenoxy]phenyl][3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
 20 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-(3-(trifluoromethoxy)phenyl)-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[[3-(trifluoromethyl)-phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 25 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[[3,5-dimethylphenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[[3-(trifluoromethylthio)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-[[3,5-difluorophenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
 30 (2R)-3-[[[3-(pentafluoroethyl)phenyl)methyl][3-(cyclohexylmethoxy)phenyl]-amino]-1,1,1-trifluoro-2-propanol;
 (2R)-3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;

- (2R)-3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-difluoromethoxyphenoxy)phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[3-(pentafluoroethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[3-(pentafluoroethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 (2R)-3-[[3-(3-isopropylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-cyclopropylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[3-(3-(2-furyl)phenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2,3-dichlorophenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(4-fluorophenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 (2R)-3-[[3-(4-methylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 (2R)-3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[3-(heptafluoropropyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 (2R)-3-[[3-(3,5-dimethylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-ethylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

- (2R)-3-[[3-(3-*t*-butylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-methylphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(phenoxy)phenyl][[3-(heptafluoropropyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(*N,N*-dimethylamino)phenoxy]phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 10 (2R)-3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3-(trifluoromethoxy)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3-(trifluoromethyl)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3,5-dimethylphenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3-(trifluoromethylthio)phenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[[3,5-difluorophenyl]methoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 (2R)-3-[[[3-(heptafluoropropyl)phenyl]methyl][3-[3-(cyclohexylmethoxy)phenyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 (2R)-3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-difluoromethoxyphenoxy)phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[3-(heptafluoropropyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 (2R)-3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[3-(heptafluoropropyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-trifluoromethoxyphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

- (2R)-3-[[3-(3-isopropylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-cyclopropylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[3-(3-(2-furyl)phenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2,3-dichlorophenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(4-fluorophenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-3-propanol;
- 10 (2R)-3-[[3-(4-methylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[2-fluoro-5-(trifluoro-methyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 20 (2R)-3-[[3-(3,5-dimethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-ethylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 25 (2R)-3-[[3-(3-t-butylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-methylphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 30 (2R)-3-[[3-(phenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(N,N-dimethylamino,phenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 35 (2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-

- (trifluoromethoxy)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-3-propanol;
(2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethyl)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
5 (2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
10 (2R)-3-[[[2-fluoro-5-(trifluoromethyl)phenyl]methyl][3-[cyclohexylmethoxyl-phenyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
15 (2R)-3-[[[3-(3-difluoromethoxyphenoxy)phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[2-fluoro-5-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
20 (2R)-3-[[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[2-fluoro-5-(trifluoro-methyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(3-trifluoromethoxyphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(3-isopropylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
25 (2R)-3-[[[3-(3-cyclopropylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(3-(2-furyl)phenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
30 (2R)-3-[[[3-(2,3-dichlorophenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(4-fluorophenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
(2R)-3-[[[3-(4-methylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
35 (trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;

- (2R)-3-[[3-(2-fluoro-5-bromophenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(4-chloro-3-ethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- 5 (2R)-3-[[3-[3-(1,1,2,2-tetrafluoroethoxy)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(pentafluoroethyl)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3,5-dimethylphenoxy)phenyl][[2-fluoro-4-
- 10 (trifluoromethyl)phenyl]-methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-ethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(3-*t*-butylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- 15 (2R)-3-[[3-(3-methylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]-amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(5,6,7,8-tetrahydro-2-naphthoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-(phenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)
- 20 phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[3-[3-(*N,N*-dimethylamino)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethoxy)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- 25 (3R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-(trifluoromethyl)phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3,5-dimethylphenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3-
- 30 (trifluoromethylthio)-phenyl]methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[[3,5-difluorophenyl]-methoxy]phenyl]amino]-1,1,1-trifluoro-2-propanol;
- (2R)-3-[[[2-fluoro-4-(trifluoromethyl)phenyl]methyl][3-[cyclohexylmethoxy]-phenyl]amino]-1,1,1-trifluoro-2-propanol;

(2R)-3-[[3-(2-difluoromethoxy-4-pyridyloxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;


(2R)-3-[[3-(2-trifluoromethyl-4-pyridyloxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

5 (2R)-3-[[3-(3-difluoromethoxyphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol;

(2R)-3-[[[3-(3-trifluoromethylthio)phenoxy]phenyl][[2-fluoro-4-(trifluoromethyl)-phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol; and

10 (2R)-3-[[3-(4-chloro-3-trifluoromethylphenoxy)phenyl][[2-fluoro-4-(trifluoromethyl)phenyl]methyl]amino]-1,1,1-trifluoro-2-propanol.

Another class of CETP inhibitors that finds utility with the present invention consists of quinolines of Formula XVII



Formula XVII


15

and pharmaceutically acceptable forms thereof, wherein:

A_{XVII} denotes an aryl containing 6 to 10 carbon atoms, which is optionally substituted with up to five identical or different substituents in the form of a halogen, nitro, hydroxyl, trifluoromethyl, trifluoromethoxy or a straight-chain or
20 branched alkyl, acyl, hydroxyalkyl or alkoxy containing up to 7 carbon atoms each, or in the form of a group according to the formula -NR_{XVII-4}R_{XVII-5}, wherein

R_{XVII-4} and R_{XVII-5} are identical or different and denote a hydrogen, phenyl or a straight-chain or branched alkyl containing up to 6 carbon atoms,

D_{XVII} denotes an aryl containing 6 to 10 carbon atoms, which is optionally
25 substituted with a phenyl, nitro, halogen, trifluoromethyl or trifluoromethoxy, or a radical according to the formula



wherein


R_{XVII-6} , R_{XVII-7} , $R_{XVII-10}$ denote, independently from one another, a
 cycloalkyl containing 3 to 6 carbon atoms, or an aryl containing 6 to 10 carbon
 atom or a 5- to 7-membered, optionally benzo-condensed, saturated or
 5 unsaturated, mono-, bi- or tricyclic heterocycle containing up to 4 heteroatoms
 from the series of S, N and/or O, wherein the rings are optionally substituted, in
 the case of the nitrogen-containing rings also via the N function, with up to five
 identical or different substituents in the form of a halogen, trifluoromethyl, nitro,
 10 hydroxyl, cyano, carboxyl, trifluoromethoxy, a straight-chain or branched acyl,
 alkyl, alkylthio, alkylalkoxy, alkoxy or alkoxycarbonyl containing up to 6 carbon
 atoms each, an aryl or trifluoromethyl-substituted aryl containing 6 to 10 carbon
 atoms each, or an optionally benzo-condensed, aromatic 5- to 7-membered
 heterocycle containing up to 3 heteroatoms from the series of S, N and/or O,
 15 and/or in the form of a group according to the formula $-OR_{XVII-11}$, $-SR_{XVII-12}$,
 $-SO_2R_{XVII-13}$, or $-NR_{XVII-14}R_{XVII-15}$;

$R_{XVII-11}$, $R_{XVII-12}$, and $R_{XVII-13}$ denote, independently from one another, an
 aryl containing 6 to 10 carbon atoms, which is in turn substituted with up to two
 identical or different substituents in the form of a phenyl, halogen or a straight-
 20 chain or branched alkyl containing up to 6 carbon atoms,

$R_{XVII-14}$ and $R_{XVII-15}$ are identical or different and have the meaning of
 R_{XVII-4} and R_{XVII-5} given above, or

R_{XVII-6} and/or R_{XVII-7} denote a radical according to the formula

118



R_{XVII-8} denotes a hydrogen or halogen, and

5 R_{XVII-9} denotes a hydrogen, halogen, azido, trifluoromethyl, hydroxyl, trifluoromethoxy, a straight-chain or branched alkoxy or alkyl containing up to 6 carbon atoms each, or a radical according to the formula $NR_{XVII-16}R_{XVII-17}$;

$R_{XVII-16}$ and $R_{XVII-17}$ are identical or different and have the meaning of R_{XVII-4} and R_{XVII-5} above; or

10 R_{XVII-8} and R_{XVII-9} together form a radical according to the formula $=O$ or $=NR_{XVII-18}$;

$R_{XVII-18}$ denotes a hydrogen or a straight-chain or branched alkyl, alkoxy or acyl containing up to 6 carbon atoms each;

15 L_{XVII} denotes a straight-chain or branched alkylene or alkenylene chain containing up to 8 carbon atoms each, which are optionally substituted with up to two hydroxyl groups;

T_{XVII} and X_{XVII} are identical or different and denote a straight-chain or branched alkylene chain containing up to 8 carbon atoms; or

T_{XVII} and X_{XVII} denotes a bond;

V_{XVII} denotes an oxygen or sulfur atom or $-NR_{XVII-19}$;

20 $R_{XVII-19}$ denotes a hydrogen or a straight-chain or branched alkyl containing up to 6 carbon atoms or a phenyl;

25 E_{XVII} denotes a cycloalkyl containing 3 to 8 carbon atoms, or a straight-chain or branched alkyl containing up to 8 carbon atoms, which is optionally substituted with a cycloalkyl containing 3 to 8 carbon atoms or a hydroxyl, or a phenyl, which is optionally substituted with a halogen or trifluoromethyl;

R_{XVII-1} and R_{XVII-2} are identical or different and denote a cycloalkyl containing 3 to 8 carbon atoms, hydrogen, nitro, halogen, trifluoromethyl, trifluoromethoxy, carboxy, hydroxy, cyano, a straight-chain or branched acyl, alkoxycarbonyl or alkoxy with up to 6 carbon atoms, or $NR_{XVII-20}R_{XVII-21}$;

30 $R_{XVII-20}$ and $R_{XVII-21}$ are identical or different and denote hydrogen, phenyl, or a straight-chain or branched alkyl with up to 6 carbon atoms; and or

R_{XVII-1} and/or R_{XVII-2} are straight-chain or branched alkyl with up to 6 carbon atoms, optionally substituted with halogen, trifluoromethoxy, hydroxy, or a straight-chain or branched alkoxy with up to 4 carbon atoms, aryl containing 6-10 carbon atoms optionally substituted with up to five of the same or different substituents selected from halogen, cyano, hydroxy, trifluoromethyl, trifluoromethoxy, nitro, straight-chain or branched alkyl, acyl, hydroxyalkyl, alkoxy with up to 7 carbon atoms and $NR_{XVII-22}R_{XVII-23}$;

$R_{XVII-22}$ and $R_{XVII-23}$ are identical or different and denote hydrogen, phenyl or a straight-chain or branched alkyl up to 6 carbon atoms; and/or

R_{XVII-1} and R_{XVII-2} taken together form a straight-chain or branched alkene or alkane with up to 6 carbon atoms optionally substituted with halogen, trifluoromethyl, hydroxy or straight-chain or branched alkoxy with up to 5 carbon atoms;


R_{XVII-3} denotes hydrogen, a straight-chain or branched acyl with up to 20 carbon atoms, a benzoyl optionally substituted with halogen, trifluoromethyl, nitro or trifluoromethoxy, a straight-chained or branched fluoroacyl with up to 8 carbon atoms and 7 fluoro atoms, a cycloalkyl with 3 to 7 carbon atoms, a straight chained or branched alkyl with up to 8 carbon atoms optionally substituted with hydroxyl, a straight-chained or branched alkoxy with up to 6 carbon atoms optionally substituted with phenyl which may in turn be substituted with halogen, nitro, trifluoromethyl, trifluoromethoxy, or phenyl or a tetrazol substituted phenyl, and/or an alkyl that is optionally substituted with a group according to the formula $-OR_{XVII-24}$;

$R_{XVII-24}$ is a straight-chained or branched acyl with up to 4 carbon atoms or benzyl.

Compounds of Formula XVII and their methods of manufacture are disclosed in PCT Publication No. WO 98/39299, which is incorporated herein by reference in its entirety for all purposes.

Another class of CETP inhibitors that finds utility with the present invention consists of 4-Phenyltetrahydroquinolines of Formula XVIII

120




Formula XVIII

5 , N oxides thereof, and pharmaceutically acceptable forms thereof, wherein:

A_{XVIII} denotes a phenyl optionally substituted with up to two identical or different substituents in the form of halogen, trifluoromethyl or a straight-chain or branched alkyl or alkoxy containing up to three carbon atoms;

D_{XVIII} denotes the formula



10

$R_{XVIII-5}$ and $R_{XVIII-6}$ are taken together to form $=O$; or

$R_{XVIII-5}$ denotes hydrogen and $R_{XVIII-6}$ denotes halogen or hydrogen; or

$R_{XVIII-5}$ and $R_{XVIII-6}$ denote hydrogen;

15 $R_{XVIII-7}$ and $R_{XVIII-8}$ are identical or different and denote phenyl, naphthyl, benzothiazolyl, quinoliny, pyrimidyl or pyridyl with up to four identical or different substituents in the form of halogen, trifluoromethyl, nitro, cyano, trifluoromethoxy, $-SO_2-CH_3$ or $NR_{XVIII-9}R_{XVIII-10}$;

$R_{XVIII-9}$ and $R_{XVIII-10}$ are identical or different and denote hydrogen or a straight-chained or branched alkyl of up to three carbon atoms;

20 E_{XVIII} denotes a cycloalkyl of from three to six carbon atoms or a straight-chained or branched alkyl of up to eight carbon atoms;

$R_{XVIII-1}$ denotes hydroxy;

$R_{XVIII-2}$ denotes hydrogen or methyl;

25 $R_{XVIII-3}$ and $R_{XVIII-4}$ are identical or different and denote straight-chained or branched alkyl of up to three carbon atoms; or

R_{XVIII-3} and R_{XVIII-4} taken together form an alkenylene made up of between two and four carbon atoms.

Compounds of Formula XVIII and their methods of manufacture are disclosed in PCT Publication No. WO 99/15504 and United States Patent No. 6,291,477, both of which are incorporated herein by reference in their entireties for all purposes.


The invention is particularly advantageous for the class of drugs which are both acid-sensitive and low-solubility. Exemplary acid-sensitive, low-solubility drugs include (+)-N-{3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl}-N-hydroxyurea; omeprazole; etoposide; famotidine; erythromycin; quinapril; lansoprazole; progabide; as well as CCR1 inhibitors such as quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3-fluorobenzyl-2(S),7-dihydroxy-7-methyloctyl]amide and quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide.

The invention is useful for improving the intrinsic dissolution rate of compounds selected from the following. The intrinsic dissolution rate is defined as the rate of dissolution of a pure pharmaceutical active ingredient when conditions such as surface area, agitation-stirring speed, pH and ionic-strength of the dissolution medium are kept constant. Intrinsic dissolution rate is further defined as being measured in water at 37°C using a USP II dissolution apparatus equipped with a Wood's apparatus (Wood, JH; Syarto, JE and Letterman, H: J.Pharm. Sci. 54 (1965), 1068) with a stirring speed of 50 rpm. The intrinsic dissolution rate is defined in terms of mg of drug dissolved per minute from a unit surface area, therefore, the intrinsic dissolution rate is referred to in units of mg/min.cm².

The compositions and methods of the invention are particularly useful for compounds with an intrinsic dissolution rate of preferably less than 0.1 mg/min.cm² and more preferably with less than 0.05 mg/min.cm².

Turning now to the chemical structures of specific CCR1 inhibitors, one class of CCR1 inhibitors that finds utility with the present invention consists of dihydroxyhexanoic acid derivatives having the Formula CCR1-I

122



CCR1-I

- wherein R_1 is (C_2-C_9) heteroaryl optionally substituted with one, two or three substituents independently selected from the group consisting of hydrogen, halo, cyano, (C_1-C_6)alkyl optionally substituted with one, two or three fluorine atoms, hydroxy, hydroxy- (C_1-C_6) alkyl, (C_1-C_6) alkoxy, (C_1-C_6) alkoxy (C_1-C_6) alkyl, $HO-(C=O)-$, (C_1-C_6) alkyl- $O-(C=O)-$, $HO-(C=O)-$ (C_1-C_6) alkyl, (C_1-C_6) alkyl- $O-(C=O)-$ (C_1-C_6) alkyl, (C_1-C_6) alkyl- $(C=O)-O-$, (C_1-C_6) alkyl- $(C=O)-O-(C_1-C_6)$ alkyl, $H(O=C)-$, $H(O=C)-(C_1-C_6)$ alkyl, (C_1-C_6) alkyl- $(O=C)-$, (C_1-C_6) alkyl- $(O=C)-(C_1-C_6)$ alkyl, NO_2 , amino, (C_1-C_6) alkylamino, $[(C_1-C_6)alkyl]_2$ amino, amino (C_1-C_6) alkyl, (C_1-C_6) alkylamino (C_1-C_6) alkyl, $[(C_1-C_6)alkyl]_2$ amino (C_1-C_6) alkyl, $H_2N-(C=O)-$, (C_1-C_6) alkyl- $NH-(C=O)-$, $[(C_1-C_6)alkyl]_2N-(C=O)-$, $H_2N(C=O)-(C_1-C_6)$ alkyl, (C_1-C_6) alkyl- $HN(C=O)-(C_1-C_6)$ alkyl, $[(C_1-C_6)alkyl]_2N-(C=O)-(C_1-C_6)$ alkyl, $H(O=C)-NH-$, (C_1-C_6) alkyl- $(C=O)-NH-$, (C_1-C_6) alkyl- $(C=O)-[NH](C_1-C_6)$ alkyl, (C_1-C_6) alkyl- $(C=O)-[N(C_1-C_6)alkyl](C_1-C_6)$ alkyl, (C_1-C_6) alkyl- $S-$, (C_1-C_6) alkyl- $(S=O)-$, (C_1-C_6) alkyl- SO_2- , (C_1-C_6) alkyl- SO_2-NH- , H_2N-SO_2- , $H_2N-SO_2-(C_1-C_6)$ alkyl, (C_1-C_6) alkyl- $HN-SO_2-(C_1-C_6)$ alkyl, $[(C_1-C_6)alkyl]_2N-SO_2-(C_1-C_6)$ alkyl, CF_3SO_3- , (C_1-C_6) alkyl- SO_3- , phenyl, (C_3-C_{10}) cycloalkyl, (C_2-C_9) heterocycloalkyl, and (C_2-C_9) heteroaryl;

- wherein R_2 is phenyl- $(CH_2)_m-$, naphthyl- $(CH_2)_m-$, (C_3-C_{10}) cycloalkyl- $(CH_2)_m-$, (C_1-C_6) alkyl or (C_2-C_9) heteroaryl- $(CH_2)_m-$, wherein each of said phenyl, naphthyl, (C_3-C_{10}) cycloalkyl or (C_2-C_9) heteroaryl moieties of said phenyl- $(CH_2)_m-$, naphthyl- $(CH_2)_m-$, (C_3-C_{10}) cycloalkyl- $(CH_2)_m-$ or (C_2-C_9) heteroaryl- $(CH_2)_m-$ groups may optionally be substituted with one, two, or three substituents independently selected from the group consisting of hydrogen, halo, cyano, (C_1-C_6)alkyl, hydroxy, hydroxy- (C_1-C_6) alkyl, (C_1-C_6) alkoxy, (C_1-C_6) alkoxy (C_1-C_6) alkyl, $HO-(C=O)-$, (C_1-C_6) alkyl- $O-(C=O)-$, $HO-(C=O)-$ (C_1-C_6) alkyl, (C_1-C_6) alkyl- $O-(C=O)-$ (C_1-C_6) alkyl, (C_1-C_6) alkyl- $(C=O)-O-$, (C_1-C_6) alkyl- $(C=O)-O-(C_1-C_6)$ alkyl, $H(O=C)-$, $H(O=C)-(C_1-C_6)$ alkyl, (C_1-C_6) alkyl- $(O=C)-$, (C_1-C_6) alkyl- $(O=C)-(C_1-C_6)$ alkyl, NO_2 , amino,

(C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 5 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-
 (S=O)-, (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-
 (C₁-C₆)alkyl, (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl,
 CF₃SO₃-, (C₁-C₆)alkyl-SO₃-, phenyl, phenoxy, benzyloxy, (C₃-C₁₀)cycloalkyl,
 10 (C₂-C₉)heterocycloalkyl, and (C₂-C₉)heteroaryl;
 wherein R³ is hydrogen, (C₁-C₁₀)alkyl, (C₃-C₁₀)cycloalkyl-(CH₂)_n-, (C₂-
 C₉)heterocycloalkyl-(CH₂)_n-, (C₂-C₉)heteroaryl-(CH₂)_n- or aryl-(CH₂)_n-; wherein n
 is an interger from zero to six;

wherein said R₃ (C₁-C₁₀)alkyl group may optionally be substituted with
 15 one or more substituents, (preferably from one to three substituents)
 independently selected from hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-
 (C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-
 (C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-
 (C=O)-O-, (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl,
 20 (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 25 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 30 (C₂-C₉)heteroaryl; and wherein any of the carbon-carbon single bonds of said
 (C₁-C₁₀)alkyl may optionally be replaced by a carbon-carbon double bond;
 wherein the (C₃-C₁₀)cycloalkyl moiety of said R₃ (C₃-C₁₀)cycloalkyl-
 (CH₂)_n- group may optionally be substituted by one to three substituents
 independently selected from the group consisting of hydrogen, halo, CN,
 35 (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy,

- (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 5 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 10 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl HN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl;
 15 wherein the (C₂-C₉)heterocycloalkyl moiety of said R₃ (C₂-
 C₉)heterocycloalkyl-(CH₂)_n- group may contain from one to three heteroatoms
 independently selected from nitrogen, sulfur, oxygen, >S(=O), >SO₂ or >NR⁶,
 wherein said (C₂-C₉)heterocycloalkyl moiety of said (C₂-C₉)heterocycloalkyl-
 (CH₂)_n- group may optionally be substituted on any of the ring carbon atoms
 20 capable of forming an additional bond (preferably one to three substituents per
 ring) with a substituent independently selected from the group consisting of
 hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy,
 (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 25 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 30 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,

(C₁-C₆)alkyl-SO₃⁻, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and (C₂-C₉)heteroaryl;

wherein the (C₂-C₉)heteroaryl moiety of said R³ (C₂-C₉)heteroaryl-(CH₂)_n- group may contain from one to three heteroatoms independently selected from
 5 nitrogen, sulfur or oxygen, wherein said (C₂-C₉)heteroaryl moiety of said (C₂-C₉)heteroaryl-(CH₂)_n- group may optionally be substituted on any of the ring carbon atoms capable of forming an additional bond (preferably one to three substituents per ring) with a substituent selected from the group consisting of
 10 hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 15 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-, (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 20 (C₁-C₆)alkyl-SO₂⁻, (C₁-C₆)alkyl-SO₂⁻NH-, H₂N-SO₂⁻, H₂N-SO₂⁻(C₁-C₆)alkyl, (C₁-C₆)alkylHN-SO₂⁻(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂⁻(C₁-C₆)alkyl, CF₃SO₃⁻,
 (C₁-C₆)alkyl-SO₃⁻, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and (C₂-C₉)heteroaryl; and

wherein said aryl moiety of said R₃ aryl-(CH₂)_n- group is optionally
 25 substituted phenyl or naphthyl, wherein said phenyl and naphthyl may optionally be substituted with from one to three substituents independently selected from the group consisting of hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 30 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 35 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-

NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl HN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 5 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl;

or R₃ and the carbon to which it is attached form a five to seven
 membered carbocyclic ring, wherein any of the carbon atoms of said five
 membered carbocyclic ring may optionally be substituted with a substituent
 10 selected from the group consisting of hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy,
 hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-,
 (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-, (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-,
 H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂,
 15 amino, (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 NH-, (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 20 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl; wherein one of the carbon-carbon bonds of said five to seven
 25 membered carbocyclic ring may optionally be fused to an optionally substituted
 phenyl ring, wherein said substituents may be independently selected from
 hydrogen, halo, CN, (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy,
 (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 30 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl,
 (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino,
 (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 35 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-

NH-, (C₁-C₆)alkyl(C=O)-NH-, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-,
 (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 5 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl;

wherein R₄ is hydrogen, (C₁-C₆)alkyl, hydroxy, (C₁-C₆)alkoxy, hydroxy(C₁-
 C₆)alkyl, (C₁-C₆)alkoxy(C=O)-, (C₃-C₁₀)cycloalkyl-(CH₂)_q-,
 (C₂-C₉)heterocycloalkyl-(CH₂)_q-, (C₂-C₉)heteroaryl-(CH₂)_q-, phenyl-(CH₂)_q-, or
 10 naphthyl-(CH₂)_q-; wherein said (C₂-C₉)heterocycloalkyl, (C₂-C₉)heteroaryl,
 phenyl and naphthyl groups may be optionally substituted with one or two
 substituents from the group consisting of hydrogen, halo, cyano, (C₁-C₆)alkyl,
 hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-
 (C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-
 15 (C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-, (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-,
 H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂,
 amino, (C₁-C₆)alkylamino, [(C₁-C₆)alkyl]₂ amino, amino(C₁-C₆)alkyl,
 (C₁-C₆)alkylamino (C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-,
 (C₁-C₆)alkyl-NH-(C=O)-, [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl,
 20 (C₁-C₆)alkyl-HN(C=O)-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-
 NH-, (C₁-C₆)alkyl(C=O)-NH-, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl,
 (C₁-C₆)alkyl(C=O)-[N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-
 , (C₁-C₆)alkyl-SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl,
 (C₁-C₆)alkylHN-SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-,
 25 (C₁-C₆)alkyl-SO₃-, phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and
 (C₂-C₉)heteroaryl;

wherein R₅ is hydrogen, (C₁-C₆)alkyl or amino; or

R₄ and R₅ together with the nitrogen atom to which they are attached
 form a (C₂-C₉)heterocycloalkyl group optionally substituted with one or two
 30 substituents selected from the group consisting of hydrogen, halo, cyano,
 (C₁-C₆)alkyl, hydroxy, hydroxy-(C₁-C₆)alkyl, (C₁-C₆)alkoxy,
 (C₁-C₆)alkoxy(C₁-C₆)alkyl, HO-(C=O)-, (C₁-C₆)alkyl-O-(C=O)-, HO-(C=O)-
 (C₁-C₆)alkyl, (C₁-C₆)alkyl-O-(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-(C=O)-O-,
 (C₁-C₆)alkyl-(C=O)-O-(C₁-C₆)alkyl, H(O=C)-, H(O=C)-(C₁-C₆)alkyl, (C₁-C₆)
 35 alkyl(O=C)-, (C₁-C₆)alkyl(O=C)-(C₁-C₆)alkyl, NO₂, amino, (C₁-C₆)alkylamino,

[(C₁-C₆)alkyl]₂ amino, amino(C₁-C₆)alkyl, (C₁-C₆)alkylamino (C₁-C₆)alkyl,
 [(C₁-C₆)alkyl]₂amino(C₁-C₆)alkyl, H₂N-(C=O)-, (C₁-C₆)alkyl-NH-(C=O)-,
 [(C₁-C₆)alkyl]₂N-(C=O)-, H₂N(C=O)-(C₁-C₆)alkyl, (C₁-C₆)alkyl-HN(C=O)-
 (C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-(C=O)-(C₁-C₆)alkyl, H(O=C)-NH-,
 5 (C₁-C₆)alkyl(C=O)-NH, (C₁-C₆)alkyl(C=O)-[NH](C₁-C₆)alkyl, (C₁-C₆)alkyl(C=O)-
 [N(C₁-C₆)alkyl](C₁-C₆)alkyl, (C₁-C₆)alkyl-S-, (C₁-C₆)alkyl-(S=O)-, (C₁-C₆)alkyl-
 SO₂-, (C₁-C₆)alkyl-SO₂-NH-, H₂N-SO₂-, H₂N-SO₂-(C₁-C₆)alkyl, (C₁-C₆)alkylHN-
 SO₂-(C₁-C₆)alkyl, [(C₁-C₆)alkyl]₂N-SO₂-(C₁-C₆)alkyl, CF₃SO₃-, (C₁-C₆)alkyl-SO₃-,
 phenyl, (C₃-C₁₀)cycloalkyl, (C₂-C₉)heterocycloalkyl, and (C₂-C₉)heteroaryl;
 10 wherein R⁶ is hydrogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy-(CH₂)_g-,
 (C₁-C₆)alkoxy(C=O)-(CH₂)_g-, (C₁-C₆)alkyl-(SO₂)-(CH₂)_g-, (C₆-C₁₀)aryloxy-(CH₂)_g-,
 (C₆-C₁₀)aryloxy(C=O)-(CH₂)_g-, or (C₆-C₁₀)aryl-(SO₂)-(CH₂)_g-;
 wherein g is an integer from zero to four;
 wherein m is an integer from zero to four;
 15 wherein n is an integer from zero to six;
 with the proviso that when one of R⁴ or R⁵ is hydrogen, and the other of R⁴ or
 R⁵ is (C₁-C₆)alkyl; R² is (C₃-C₁₀)cycloalkyl or isopropyl and R³ is (C₃-C₅)alkyl,
 phenyl, methylvinyl, dimethylvinyl, halovinyl, hydroxy(C₁-C₃)alkyl or amino(C₁-
 C₄)alkyl then R¹ must be other than indol-5-yl, 6-azaindol-2-yl, 2,3-dichloro-
 20 pyrrol-5-yl, 4-hydroxyquinolin-3-yl, 2-hydroxyquinoxalin-3-yl, 6-azaindolin-3-yl,
 or optionally substituted indol-2 or 3-yl;
 and the pharmaceutically acceptable salts of such compounds.

Unless otherwise indicated, the alkyl and alkenyl groups referred to
 herein, as well as the alkyl moieties of other groups referred to herein (e.g.,
 25 alkoxy), may be linear or branched, and they may also be cyclic (e.g.,
 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl) or be linear or
 branched and contain cyclic moieties. Such alkyl and alkoxy groups may be
 substituted with one, two or three halogen and/or hydroxy atoms, preferably
 fluorine atoms.

30 Unless otherwise indicated, "halogen" includes fluorine, chlorine,
 bromine, and iodine.

"(C₃-C₁₀)cycloalkyl" when used herein refers to cycloalkyl groups
 containing zero to two levels of unsaturation such as cyclopropyl, cyclobutyl,
 cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, 1,3-cyclohexadiene,
 35 cycloheptyl, cycloheptenyl, bicyclo[3.2.1]octane, norbornanyl, and the like.

"(C₂-C₉)heterocycloalkyl" when used herein refers to pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydropyranyl, pyranyl, thiopyranyl, aziridinyl, oxiranyl, methylenedioxy, chromenyl, isoxazolidinyl, 1,3-oxazolidin-3-yl, isothiazolidinyl, 1,3-thiazolidin-3-yl, 1,2-pyrazolidin-2-yl, 1,3-pyrazolidin-1-yl, 5 piperidinyl, thiomorpholinyl, 1,2-tetrahydrothiazin-2-yl, 1,3-tetrahydrothiazin-3-yl, tetrahydrothiadiazinyl, morpholinyl, 1,2-tetrahydrodiazin-2-yl, 1,3-tetrahydrodiazin-1-yl, tetrahydroazepinyl, piperazinyl, chromanyl, and the like. One of ordinary skill in the art will understand that the connection of said (C₂-C₉)heterocycloalkyl rings is through a carbon or a sp³ hybridized nitrogen heteroatom. 10

"(C₂-C₉)heteroaryl" when used herein refers to furyl, thienyl, thiazolyl, pyrazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyrrolyl, triazolyl, tetrazolyl, imidazolyl, 1,3,5-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,3-oxadiazolyl, 1,3,5-thiadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, pyridyl, pyrimidyl, pyrazinyl, 15 pyridazinyl, 1,2,4-triazinyl, 1,2,3-triazinyl, 1,3,5-triazinyl, pyrazolo[3,4-b]pyridinyl, cinnolinyl, pteridinyl, purinyl, 6,7-dihydro-5H-[1]pyrindinyl, benzo[b]thiophenyl, 5, 6, 7, 8-tetrahydro-quinolin-3-yl, benzoxazolyl, benzothiazolyl, benzisothiazolyl, benzisoxazolyl, benzimidazolyl, thianaphthenyl, isothianaphthenyl, benzofuranyl, isobenzofuranyl, isoindolyl, indolyl, indoliziny, 20 indazolyl, isoquinolyl, quinolyl, phthalazinyl, quinoxaliny, quinazolinyl, benzoxazinyl, and the like. One of ordinary skill in the art will understand that the connection of said (C₂-C₉)heterocycloalkyl rings is through a carbon atom or a sp³ hybridized nitrogen heteroatom.

"Aryl" when used herein refers to phenyl or naphthyl.

25 "Protected amine" and "protected amino" refers to an amine group with one of the hydrogen atoms replaced with a protecting group (P). Any suitable protecting group may be used for amine protection. Suitable protecting groups include carbobenzyloxy, t-butoxy carbonyl (BOC) or 9-fluorenyl-methylenoxy carbonyl.

30 Compounds of Formula CCR1-I and their methods of manufacture are disclosed in commonly assigned United States Patent Application Serial No. 09/380,269, filed February 5, 1998, United States Patent Application Serial No. 09/403,218, filed January 18, 1999, PCT Publication No. WO98/38167, and PCT Publication No. WO99/40061, all of which are incorporated herein by 35 reference in their entireties for all purposes.

In a preferred embodiment, the CCR1 inhibitor is selected from one of the following compounds of Formula CCR1-I:

quinoxaline-2-carboxylic acid 4(R)-carbamoyl-1(S)-(3-chloro-benzyl)-2(S),7-dihydroxy-7-methyl-octyl]-amide;

5 7,8-difluoro-quinoline-3-carboxylic acid (1S)-benzyl-4(R)-carbamoyl-2(S),7-dihydroxy-7-methyl-octyl]-amide;

6,7,8-trifluoro-quinoline-3-carboxylic acid (1(S)-benzyl-4(R)-carbamoyl-2(S),7-dihydroxy-7-methyl-octyl)-amide;

10 quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-(3-fluoro-benzyl)-2(S),7-dihydroxy-7-methyl-octyl]-amide;

quinoxaline-2-carboxylic acid (1(S)-benzyl-2(S),7-dihydroxy-4(R)-hydroxycarbamoyl-7-methyl-octyl)-amide;

quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-(2-chloro-benzyl)-2(S),7-dihydroxy-7-methyl-octyl]-amide;

15 quinoxaline-2-carboxylic acid [1(S)-(2-fluoro-benzyl)-2(S),7-dihydroxy-4(R)-hydroxycarbamoyl-7-methyl-octyl]-amide;

quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-(2-fluoro-benzyl)-2(S),7-dihydroxy-7-methyl-octyl]-amide;

20 quinoxaline-2-carboxylic acid [1(S)-(3,4-difluoro-benzyl)-2(S),7-dihydroxy-4(R)-hydroxycarbamoyl-7-methyl-octyl]-amide;

quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-(3,4-difluoro-benzyl)-2(S),7-dihydroxy-7-methyl-octyl]-amide;

quinoxaline-2-carboxylic acid (4(R)-carbamoyl-2(S),7-dihydroxy-7-methyl-1(S)-naphthalen-1-ylmethyl-octyl)-amide;

25 7,8-difluoro-quinoline-3-carboxylic acid 1(S)-benzyl-2(S)-hydroxy-7-methyl-4(R)-methylcarbamoyl-octyl]-amide;

8-fluoro-quinoline-3-carboxylic acid 1(S)-benzyl-2(S)-hydroxy-7-methyl-4(R)-methylcarbamoyl-octyl)-amide;

30 quinoxaline-2-carboxylic acid [4(R)-carbamoyl-7-fluoro-1-(3(S)-fluoro-benzyl)-2(S)-hydroxy-7-methyl-octyl]-amide;

quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1-(2(S)-fluoro-benzyl)-2(S)-hydroxy-7-methyl-octyl]-amide;

quinoxaline-2-carboxylic acid [1(S)-benzyl-4(S)-carbamoyl-4(S)-(2,6-dimethyl-tetrahydro-pyran-4-yl)-2(S)-hydroxy-butyl]-amide;

quinoxaline-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-7-fluoro-2(S)-hydroxy-7-methyl-octyl)-amide;

quinoxaline-2-carboxylic acid 1(S)-benzyl-5-cyclohexyl-2(S)-hydroxy-4(R)-methylcarbamoyl-pentyl)-amide;

5 quinoxaline-2-carboxylic acid 1(S)-cyclohexylmethyl-2(S)-hydroxy-7-methyl-4(R)-methylcarbamoyl-octyl)-amide;

quinoxaline-2-carboxylic acid [1(S)-benzyl-2(S)-hydroxy-4(S)-hydroxycarbamoyl-4-(1-hydroxy-4-methyl-cyclohexyl)-butyl]-amide;

10 quinoxaline-2-carboxylic acid [1(S)-benzyl-4(S)-(4,4-difluoro-1-hydroxy-cyclohexyl)-2(S)-hydroxy-4-hydroxycarbamoyl-butyl]-amide;

quinoxaline-2-carboxylic acid [1(S)-benzyl-4(S)-carbamoyl-4(S)-(4,4-difluoro-cyclohexyl)-2(S)-hydroxy-butyl]-amide;

quinoline-3-carboxylic acid (1(S)-benzyl-4(S)-carbamoyl-4-cyclohexyl-2(S)-hydroxy-butyl)-amide;

15 quinoxaline-2-carboxylic acid (4(R)-carbamoyl-2(S)-hydroxy-7-methyl-1(S)-thiophen-2-ylmethyl-octyl)-amide;

quinoxaline-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-7-chloro-2(S)-hydroxy-oct-6-enyl)-amide;

20 quinoxaline-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-2(S)-hydroxy-5-phenyl-pentyl)-amide;

N-1(S)-benzyl-4(R)-carbamoyl-7-fluoro-2(S)-hydroxy-7-methyl-octyl)-5,6-dichloro-nicotinamide;


quinoxaline-2-carboxylic acid (4(R)-carbamoyl-2(S)-hydroxy-7-methyl-1(S)-thiazol-4(R)-ylmethyl-octyl)-amide;

25 benzothiazole-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-7-fluoro-2(S)-hydroxy-7-methyl-octyl)-amide; and

benzofuran-2-carboxylic acid 1(S)-benzyl-4(R)-carbamoyl-7-fluoro-2(S)-hydroxy-7-methyl-octyl)-amide.

In another preferred embodiment, the CCR1 compound has a formula

30 Ia-1:



wherein the substituents are as defined above.

In a preferred method of making the compound Ia-1, the reaction is started with Scheme 1. In the herein described processes, the substituents are as defined for CCR1-I, and the following:

R_7 is hydroxy, (C₁-C₆)alkyl, or phenyl wherein the phenyl group unsubstituted or substituted with one, two, or three (C₁-C₆)alkyl, hydroxy, or halogen groups;

R_8 is hydroxy or halogen;

R_9 is phenyl, naphthyl, (C₃-C₁₀)cycloalkyl, (C₁-C₆)alkyl or (C₂-C₉)heteroaryl, wherein each of said phenyl, naphthyl, (C₃-C₁₀)cycloalkyl or (C₂-C₉)heteroaryl groups may be unsubstituted or substituted with one, two, or three substituents independently selected from the group consisting of halogen, cyano, and (C₁-C₆)alkyl;


P is a protecting group;

X is hydroxy or halogen; and

q is 0, 1, 2, 3, or 4.

133

Scheme 1



In scheme 1 step 1, a compound of the formula (VI-1) is reduced with a reducing agent under heat to form a compound of the formula (VId-1). In one embodiment, the reducing agent is aluminum triisopropoxide and isopropanol. Preferably, the temperature is maintained above room temperature, more preferably between about 60°C and about 82°C. The product alcohol can be isolated by either cooling the reaction mixture to room temperature, diluting with more isopropanol and collecting the crystalline material or by cooling the reaction to room temperature and adding 1N HCL and water and collecting the crystalline material.

Step 2 of scheme 1 includes reacting a compound of the formula $R_7\text{-SO}_2\text{-X}$ and a compound of the formula (VId-1) in the presence of a base to form the compound of the formula (Vle-1). Any amine base is suitable, including pyridine, triethylamine, N-methylmorpholine, and diisopropylethylamine. In one embodiment, $R_7\text{-SO}_2\text{-R}_8$ is p-toluenesulfonic acid, methanesulfonic acid, sulfuric acid, or methanesulfonyl chloride. In another embodiment, the conversion of hydroxy dioxane (VId-1) to dioxane oxazolidinone (Vle-1) can be achieved by treatment of the hydroxy dioxane (VId-1) with methanesulfonyl chloride and triethylamine in tetrahydrofuran solution and heating the mixture to cause the cyclization of the mesylate formed in situ to the oxazolidinone.

In step 3 of scheme 1, a compound of the formula (VIf-1) may be formed by heating the compound of the formula (Vle-1). The reaction may proceed by dissolving compound Vle-1 in a solvent such as pyridine or N-methyl imidazole and heating the mixture for several hours at temperature from about 50°C to about 100°C; preferably at about 80°C. The mesylate (VIf-1) may be recovered by extraction into an organic solvent such as ethyl acetate and removal of the amine solvents by extraction of the solution with aqueous acid.

Step 4 of scheme 1 depicts reacting hydroxylamine hydrochloride, a compound of the formula $R_7\text{-SO}_2\text{-X}$, and a compound of the formula (VIf-1) to form a compound of the formula (VIg-1). In one embodiment, $R_7\text{-SO}_2\text{-X}$ is p-toluenesulfonic acid, methanesulfonic acid, sulfuric acid, or methanesulfonyl chloride. The reaction may occur in a solvent, such as methanol. In one embodiment, the reaction occurs in methanol with tosic acid at reflux for 8 to 24 hours. The resulting nitrile oxazolidinone contains a small amount of the corresponding ethyl ester which is not removed since it also is converted to the


desired lactone in subsequent steps.

Step 5 of scheme 1 includes a) hydrolyzing a compound of the formula (VIg-1) with an aqueous solution in the presence of a base, b) protecting the amine group of the compound so formed, and c) cyclizing the compound so
5 formed with heat and an acid catalyst. In one embodiment, the compound VIg-1 is hydrolyzed with sodium hydroxide. The pH is adjusted to approximately 10 and tetrahydrofuran and BOC dicarbonate are added. This provides the protected hydroxy acid, which may be heated in 10% acetic acid and toluene to provide the protected amine lactone (V-1).

10 The compound of formula (V-1) may also be produced according to scheme 2.

136

Scheme 2



In step 1 of scheme 2, a compound of the formula (VI-1) may be reacted with ozone to form a compound of the formula (VIa-1). The compound VI-1 may be present in a solvent, such as ethyl acetate, and the ozone introduced through sparging at a temperature below room temperature, preferably at about -15°C, until the starting dioxane ketone is substantially reacted. Any excess ozone may be removed by bubbling nitrogen through the solution. The resulting crude ketone ester mixture may be isolated after treatment with aqueous sodium bisulfite to remove any hydroperoxides.

Alternatively, in step 1 of scheme 2, the compound of the formula (VIa-1) may be formed by reacting hypochlorous acid and a compound of the formula (VI-1). Such an oxidation reaction typically produces chlorinated forms of the compound VIa-1 as side products in addition to the compound VIa-1. This
5 oxidation reaction proceeds by mixing the compound VI-1 in solvent, such as acetic acid and/or acetone, and adding sodium hypochlorite, while keeping the mixture at a low temperature, preferably at or below about 0°C.


As a means to convert the side product chlorinated forms of the compound VIa-1 to compounds of the formula V-1, the compounds formed from
10 the hypochlorous acid oxidation reaction may optionally be hydrogenated by reaction with hydrogen in the presence of a catalyst. The hydrogenation may include introducing the products from the hypochlorous acid oxidation reaction into a solvent system of tetrahydrofuran and water, followed by addition of a Pd/C catalyst. The resulting mixture is subjected to hydrogen above
15 atmospheric pressure and temperature. In one embodiment, the pressure is about 80 pounds per square inch and the temperature is maintained from about 60°C to about 70°C until the reaction is substantially complete.

In step 2 of scheme 2, the compound of the formula (VIb-1) may be formed by reacting a silylating agent and a compound of the formula (VIa-1) and
20 reacting the compound so formed with a reducing agent. In one embodiment, the reducing agent is N-selectride. In another embodiment, the silylating agent is 1,1,1,3,3,3-hexamethyl-disilazane. The reduction reaction may occur at temperatures below about 0°C, preferably below about -20°C, more preferably below about -50°C. In addition, the reducing agent may be present in slight
25 excess.

In step 3 of scheme 2, the compound of the formula (V-1) is formed by heating a compound of the formula (VIb-1) in the presence of an acid catalyst, such as acetic acid. In one embodiment, the cyclization reaction occurs by
30 introducing the compound VIb-1 into a solvent mixture, such as toluene and 10% acetic acid, at the solvent reflux temperature for 8 to 16 hours. This provides the desired lactone as a crystalline solid after work up.

One method of making the compound of the formula (VI-1) is by reacting a compound of the formula (VII-1)


138



(VII-1)

with a Grinard reagent formed in situ by addition of 2-(2-bromb-ethyl)-
[1,3]dioxane to a mixture comprising magnesium and the compound of the
formula (VII-1). In one embodiment, the mixture further comprises methyl
magnesium chloride and/or methyl magnesium bromide in a solvent. Any
exotherm formed from the reaction may be controlled by the rate of addition of
the bromide.

The compound of the formula (VII-1) may be formed by coupling N,O-
dimethylhydroxylamine hydrochloride and a compound of the formula (VIII-1)




(VIII-1)

This coupling reaction may be performed by mixed anhydride procedure. In one
mixed anhydride procedure, compound VIII-1 is combined with methylene
chloride and N-methylmorpholine is added followed by isobutyl chloroformate.
In a separate mixture, a slurry of N,O-dimethylhydroxylamine hydrochloride is
treated with N-methylmorpholine. The two reaction mixtures are combined and
then quenched with a solution of citric acid in water. This procedure preferably
operates at a temperature below about 20°C, more preferably below about 0°C.

Compounds of formula (V-1) may be used to produce compounds of the
formula (IVa1-1) according to scheme 3:

139

Scheme 3




In step 1 of scheme 3, the compound of the formula (IVa1-1) may be formed by reacting 4-halo-2-methyl-2-butene and a compound of the formula (V-1) in the presence of a base. Exemplary bases include lithium dialkyl amides such as lithium N-isopropyl-N-cyclohexylamide, lithium bis(trimethylsilyl)amide, lithium di-isopropylamide, and potassium hydride. Suitable solvents include aprotic polar solvents such as ethers (such as tetrahydrofuran, glyme or dioxane), benzene, or toluene, preferably tetrahydrofuran. The aforesaid reaction is conducted at a temperature from about -78°C to about 0°C, preferably at about -78°C. In one embodiment, alkylation of the lactone (V-1) is accomplished by reacting the lactone (V-1) with lithium bis(trimethylsilyl)amide and dimethylallyl bromide in tetrahydrofuran at a temperature from about -78°C to about -50°C. Reaction times range from several hours or if an additive such as dimethyl imidazolidinone is present, the reaction may be complete in minutes.

Compounds of formula (IVa1-1) may be used to produce compounds of the formula (Ia-1) according to scheme 4:

140

Scheme 4



In step 1 of scheme 4, a compound of the formula (IIIa1-1) is formed by
 5 reacting a compound of the formula (IVa1-1) with phosphoric acid. Preferably,
 this reaction occurs in any suitable solvent, such as non-alcoholic solvents. Two
 preferred solvents include tetrahydrofuran and dichloroethane. The reaction
 may take place at any suitable temperature, preferably from about -25°C to
 about 120°C, more preferably from about 15°C to about 40°C. Reaction time is

dependent on temperature and batch size, amount other factors, but typically reaction time is from about 2 hours to about 14 hours.


Step 2 of scheme 4 depicts coupling a compound IIIa1-1 with a compound having the formula $R_1\text{-CO-X}$ to form a compound having the formula (IIa1-1). This coupling reaction is generally conducted at a temperature from about -30°C to about 80°C , preferably from about 0°C to about 25°C . The coupling reaction may occur with a coupling reagent that activates the acid functionality. Exemplary coupling reagents include dicyclohexylcarbodiimide/hydroxybenzotriazole (DCC/HBT), N-3-dimethylaminopropyl-N'-ethylcarbodiimide (EDC/HBT), 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), carbonyl diimidazole (CDI), and diethylphosphoryl cyanide. The coupling is conducted in an inert solvent, preferably an aprotic solvent, such as tetrahydrofuran, acetonitrile, dichloromethane, chloroform, or N,N-dimethylformamide. One preferred solvent is tetrahydrofuran. In one embodiment, quinoxaline acid is combined with CDI in anhydrous tetrahydrofuran and heated to provide the acyl imidazole. Compound IIIa1-1 is added to the acyl imidazole at room temperature to form the compound IIa1-1.

Step 3 of scheme 4 includes reacting the compound of formula IIa1-1 with an amine having a formula NHR_4R_5 to form a compound of the formula (Ia-1). In one embodiment, the amine is ammonia either anhydrous in an organic solvent or as an aqueous solution of ammonium hydroxide added to a polar solvent at a temperature from about -10°C to about 35°C , preferably at about 30°C . Suitable solvents include, alcohols, such as methanol, ethanol, or butanols; ethers such as tetrahydrofuran, glyme or dioxane; or a mixture thereof, including aqueous mixtures. Preferably the solvent is methanol. In one embodiment, the compound IIa1-1 is dissolved in methanol which has been saturated with ammonia gas. In another embodiment, the compound IIa1-1 in methanol is treated with ammonium hydroxide in tetrahydrofuran at room temperature.

Scheme 5 represents an alternative method to form compounds of formula Ia-1 from compounds of formula IVa1-1.

142

Scheme 5



In step 1 of scheme 5, a compound of the formula (IVa1-1) is reacted with a compound of the formula $R_9\text{-SO}_2\text{-X}$ to form a compound of the formula (IVa2-1). Any suitable acidic deprotection reaction may be performed. In one example, an excess of p-toluenesulfonic acid hydrate in ethyl acetate is introduced to the compound IVa1-1 at room temperature. Suitable solvents include ethyl acetate, alcohols, tetrahydrofuran, and mixtures thereof. The reaction may proceed at ambient or elevated temperatures. Typically, the reaction is substantially complete within two and twelve hours. The resulting compound IVa2-1 may be crystallized and separated from the reaction mixture, and may be further purified to remove impurities by recrystallization from hot ethyl acetate.

In step 2 of scheme 5, the compound IVa2-1 may be coupled with a compound having the formula $R_1\text{-CO-X}$ to form a compound of the formula (IIIa2-1). This coupling reaction is generally conducted at a temperature from about -30°C to about 80°C, preferably from about 0°C to about 25°C. The coupling reaction may occur with a coupling reagent that activates the acid functionality. Exemplary coupling reagents include dicyclohexylcarbodiimide/hydroxybenzotriazole (DCC/HBT), N-3-dimethylaminopropyl-N'-ethylcarbodiimide (EDC/HBT), 2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), carbonyl diimidazole (CDI)/dimethylaminopyridine (DMAP), and diethylphosphorylcyanide. The coupling is conducted in an inert solvent, preferably an aprotic solvent, such as acetonitrile, dichloromethane, chloroform, or N,N-dimethylformamide. One preferred solvent is methylene chloride. In one embodiment, quinoxaline acid is combined with methylene chloride, oxalyl chloride and a catalytic amount of N,N-dimethylformamide to form an acid chloride complex. The compound IVa2-1 is added to the acid chloride complex followed by triethylamine at a temperature from about 0°C to about 25°C to form the compound IIIa2-1.

Step 3 of scheme 5 includes reacting a compound IIIa2-1 with trifluoroacetic acid to produce a compound of the formula (IIa2-1). In one embodiment, the hydration with trifluoroacetic acid occurs in methylene chloride solution at room temperature. The hydration may take several hours to complete at room temperature. A catalytic amount of sulfuric acid can be added to the reaction solution to increase the rate of reaction.

Step 4 of scheme 5 includes reacting the compound of formula IIa2-1 with an amine having a formula NHR_4R_5 to form a compound of the formula (Ia-1). In one embodiment, the amine is ammonia either anhydrous in an organic solvent or as an aqueous solution of ammonium hydroxide added to a polar solvent at a temperature from about -10°C to about 35°C , preferably at about 30°C . Suitable solvents include, alcohols, such as methanol, ethanol, or butanols; ethers such as tetrahydrofuran, glyme or dioxane; or a mixture thereof, including aqueous mixtures. Preferably the solvent is methanol. In one embodiment, the compound IIa2-1 is dissolved in methanol which has been saturated with ammonia gas. In another embodiment, the compound IIa2-1 in methanol is treated with ammonium hydroxide in tetrahydrofuran at room temperature.

NEUTRAL DISPERSION POLYMERS

The neutral polymers suitable for use in the dispersions of the present invention are pharmaceutically acceptable and should be inert, in the sense that they do not chemically react with the drug in an adverse manner when present in the composition. The polymer should also have an aqueous-solubility of at least 0.1 mg/mL over at least a portion of the pH range of 1-8.

The polymer is "neutral," meaning that the dispersion polymer possesses substantially no acidic functional groups. By "substantially no acidic functional groups" is meant that the number of acidic groups covalently attached to the polymer is less than about 0.05 milliequivalents per gram of polymer. Preferably, the number is less than about 0.02 milliequivalents per gram of polymer. By "acidic groups" is meant functional groups that, when attached to the polymer, have pK_a values in a humid or aqueous environment of about 5 or less. Preferably, the pK_a value of the functional groups on the neutral polymer is greater than about 6. Thus, the neutral polymers may contain ionic groups as long as the groups are not acidic.

However, for many drugs, particularly those that are also sensitive to basic conditions, e.g., "base-sensitive drugs," it is preferred that the neutral polymers be substantially nonionizable. By "substantially nonionizable" is meant that the number of "ionizable groups" covalently attached to the polymer is less than about 0.05 milliequivalents per gram of polymer and preferably less than about 0.02 milliequivalents per gram of polymer. "Ionizable groups" are

those that are at least about 10% ionized over at least a portion of the physiologically relevant pH 1 to 8 range and thus such groups have pK_a values of about 0 to 9.

5 The neutral dispersion polymer may be cellulosic or non-cellulosic. A preferred class of neutral cellulosic dispersion polymers are those with at least one ester- and/or ether- linked substituent in which the polymer has a degree of substitution of at least 0.05 for each substituent. It should be noted that in the polymer nomenclature used herein, ether-linked substituents are recited prior to "cellulose" as the moiety attached to the ether group; for example, "methyl
10 cellulose" has a methyl moiety ether-linked to the polymer. In contrast, ester-linked substituents are recited after "cellulose" as the carboxylate; for example, "cellulose acetate" has an acetate moiety ester-linked to the polymer.

It should also be noted that a polymer name such as "cellulose acetate butyrate" refers to any of the family of cellulosic polymers that have acetate and
15 butyrate groups attached via ester linkages to a significant fraction of the cellulosic polymer's hydroxyl groups. Generally, the degree of substitution of each substituent group can range from about 0.05 to 2.9 as long as the other criteria of the polymer are met. "Degree of substitution" refers to the average number of the three hydroxyls per saccharide repeat unit on the cellulose chain that have been substituted. For example, if all of the hydroxyls on the cellulose
20 chain have been butyrate substituted, the butyrate degree of substitution is 3. Also included within each polymer family type are cellulosic polymers that have additional substituents added in relatively small amounts that do not substantially alter the performance of the polymer.

25 Polymer substituents may be either non-ionizable or ionizable; however, the ionizable groups may not be acidic groups. Exemplary ether-linked non-ionizable substituents include: alkyl groups, such as methyl, ethyl, propyl, butyl, etc.; hydroxy alkyl groups such as hydroxymethyl, hydroxyethyl, hydroxypropyl, etc.; and aryl groups such as phenyl. Exemplary ester-linked non-ionizable
30 groups include: alkylate groups, such as acetate, propionate, butyrate, etc.; and arylate groups such as phenylate. However, when ester-linked non-ionizable groups are included, the polymer may need to include a sufficient amount of a hydrophilic substituent so that the polymer has at least some water solubility at any physiologically relevant pH of from 1 to 8. In the case of neutral dispersion
35 polymers, such hydrophilic substituents may consist of non-acidic ionizable

groups such as amino-functionalized groups or phenolate groups. In the case of non-ionizable neutral dispersion polymers, such hydrophilic groups are non-ionizable substituents such as alcohol, ether or ester groups.

Exemplary neutral non-ionizable cellulosic polymers that may be used to form the dispersion include: hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and hydroxyethyl ethyl cellulose.

Exemplary neutral, but ionizable cellulosic polymers, include aminoethyl cellulose, aminoethyl cellulose acetate, hydroxypropyl amino ethyl cellulose and hydroxybenzyl cellulose.

Another class of neutral dispersion polymers is non-cellulosic, neutral polymers. Such polymers may be either non-ionizable or ionizable. Exemplary non-ionizable, neutral polymers include vinyl polymers and copolymers having substituents of hydroxyl, alkylacyloxy, and cyclicamido. Exemplary non-cellulosic, neutral polymers include hydroxyethyl methacrylate, polyvinylhydroxyethyl ether, polyethylene glycol, and polyoxyethylene-polyoxypropylene block copolymers also known as poloxamers.

Exemplary ionizable neutral polymers include amine-functionalized polyacrylates and polymethacrylates, some of which are also sold as EUDRAGITS manufactured by Rohm Tech Inc., and neutral proteins.

A preferred subset of neutral polymers are those that are also concentration-enhancing. Dispersion polymers that are both neutral and also concentration-enhancing are generally amphiphilic in that they possess substituents that are relatively hydrophobic and substituents that are relatively hydrophilic.

Amphiphilic cellulose may be prepared by substituting the cellulose at any or all of the 3 hydroxyl substituents present on each saccharide repeat unit with at least one relatively hydrophobic substituent. Hydrophobic substituents may be essentially any substituent that, if substituted to a high enough level or degree of substitution, can render the cellulosic polymer essentially aqueous insoluble. Hydrophilic regions of the polymer can be either those portions that are relatively unsubstituted, since the unsubstituted hydroxyls are themselves relatively hydrophilic, or those regions that are substituted with hydrophilic substituents. Examples of hydrophobic substituents include ether-linked alkyl

groups such as methyl, ethyl, propyl, butyl, etc.; or ester-linked alkyl groups such as acetate, propionate, butyrate, etc.; and ether- and/or ester-linked aryl groups such as phenyl, benzoate, or phenylate. As discussed above, hydrophilic substituents may consist of non-acidic ionizable groups such as amino-functionalized groups or phenolate groups. In the case of non-ionizable neutral dispersion polymers, such hydrophilic groups are non-ionizable substituents such as alcohol, ether or ester groups.


Exemplary amphiphilic polymers include non-ionizable cellulose such as hydroxypropyl methyl cellulose, hydroxyethyl methyl cellulose and hydroxyethyl acetate; non-acidic ionizable cellulose such as amino ethyl cellulose acetate and hydroxybenzyl cellulose; and non-ionizable non-cellulose such as polyvinylpyrrolidone ethylene/vinyl alcohol copolymers and polyoxyethylene-polyoxypropylene block copolymers (also referred to as poloxamers); and ionizable non-cellulose such as amine-functionalized polyacrylates and polymethacrylates. When the amphiphilic polymer is hydroxypropyl methyl cellulose, it is preferred that it is provided in a grade that possesses a dynamic viscosity that is 60 mPa-s or less, more preferably 10 mPa-s or less and most preferably between 1 mPa-s and 6 mPa-s, when measured in a 2% weight/volume solution at 20°C.

A preferred class of neutral non-cellulosic polymers are comprised of vinyl copolymers of a hydrophilic, hydroxyl-containing repeat unit and a hydrophobic, alkyl- or aryl-containing repeat unit. Such neutral vinyl copolymers are termed "amphiphilic hydroxyl-functional vinyl copolymers." Amphiphilic hydroxyl-functional vinyl copolymers are exceptional in that they are both non-ionic and yet, surprisingly, when used as dispersion polymers for low-solubility drugs, yield solid amorphous dispersions that provide high levels of drug concentration enhancement when dosed to an aqueous environment of use. Such polymers may be used with any low-solubility drug, and not simply acid-sensitive drugs.

Dispersions of such copolymers are believed to provide high concentration enhancements due to the amphiphilicity of these copolymers which provide both sufficient hydrophobic groups to interact with the hydrophobic, low-solubility drugs and also sufficient hydrophilic groups to have sufficient aqueous solubility for good dissolution. The copolymeric structure of the amphiphilic hydroxyl-functional vinyl copolymers also allows their

hydrophilicity and hydrophobicity to be adjusted to maximize performance with a specific low-solubility drug.

The preferred copolymers have the general structure:



wherein A and B represent "hydrophilic, hydroxyl-containing" and "hydrophobic" substituents, respectively, and n and m represent the average number of hydrophilic vinyl repeat units and average number of hydrophobic vinyl repeat units respectively per polymer molecule. Copolymers may be block copolymers, random copolymers or they may have structures anywhere between these two extremes. The sum of n and m is generally from about 50 to about 20,000 and therefore the polymers have molecular weights from about 2,500 to about 1,000,000 daltons.

The hydrophilic, hydroxyl-containing repeat units, "A," may simply be hydroxyl (-OH) or it may be any short-chain, 1 to 6 carbon, alkyl with one or more hydroxyls attached thereto. The hydroxyl-substituted alkyl may be attached to the vinyl backbone via carbon-carbon or ether linkages. Thus, exemplary "A" structures include, in addition to hydroxyl itself, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxymethoxy, hydroxyethoxy and hydroxypropoxy.

The hydrophobic substituent, "B," may simply be: hydrogen (-H), in which case the hydrophobic repeat unit is ethylene; an alkyl or aryl substituent with up to 12 carbons attached via a carbon-carbon bond such as methyl, ethyl or phenyl; an alkyl or aryl substituent with up to 12 carbons attached via an ether linkage such as methoxy, ethoxy or phenoxy; an alkyl or aryl substituent with up to 12 carbons attached via an ester linkage such as acetate, propionate, butyrate or benzoate. The amphiphilic hydroxyl-functional vinyl copolymers of the present invention may be synthesized by any conventional method used to prepare substituted vinyl copolymers. Some substituted vinyl copolymers such as polyvinyl alcohol/polyvinyl acetate are well known and commercially available.

A particularly convenient subclass of amphiphilic hydroxyl-functional vinyl copolymers to synthesize are those where the hydrophobic substituent "B" comprises the hydrophilic substituent "A" to which an alkylate or arylate group is attached via an ester linkage to one or more of the hydroxyls of A. Such

copoly- mers may be synthesized by first forming the homopolymer of the hydrophobic vinyl repeat unit having the substi- tuent B, followed by hydrolysis of a portion of the ester groups to convert a portion of the hydrophobic repeat units to hydrophilic, hydroxyl-containing repeat units having the substituent A.

5 For example, partial hydrolysis of the homopolymer, polyvinylbutyrate, yields the copolymer, vinylalcohol/vinylbutyrate copolymer for which A is hydroxyl (-OH) and B is butyrate (-OOC-CH₂-CH₂-CH₃).

For all types of copolymers, the value of n must be sufficiently large relative to the value of m that the resulting copolymer is at least partially water soluble. Although the value of the ratio, n/m varies depending on the identity of A and B, it is generally at least about 1 and more commonly about 2 or more. The ratio n/m can be as high as 200. When the copolymer is formed by hydrolysis of the hydrophobic homopolymer, the relative values of n and m are typically reported in "percent hydrolysis," which is the fraction (expressed as a percent) of the total repeat units of the copolymer that are in the hydrolyzed or hydroxyl form. The percent hydrolysis, H, is given as

10 15

$$H = 100 \cdot \left(\frac{n}{n + m} \right)$$

Thus, vinylbutyrate/vinylalcohol copolymer (formed by hydrolysis of a portion of the butyrate groups) having a percent hydrolysis of 75% has an n/m ratio of 3.

20

A particularly preferred family of amphiphilic hydroxyl-functional vinyl copolymers are those where A is hydroxyl and B is acetate. Such copolymers are termed vinylacetate/vinylalcohol copolymers. Some commercial grades are also sometimes referred to simply as polyvinylalcohol. However, the true homopolymer, polyvinylalcohol is not amphiphilic, is almost entirely water insoluble, and is not a part of this invention. Preferred vinylacetate/vinylalcohol copolymers are those where H is between about 67% and 99.5%, or n/m has a value between about 2 and 200. The preferred average molecular weight is between about 2500 and 1,000,000 daltons and more preferably between about 3000 and about 100,000 daltons.

25 30

In order to form dispersions that have a high level of concentration and bioavailability enhancement, it is important that the identity of A and B, the n/m value, and average molecular weight be chosen such that the polymers are sufficiently hydrophobic that they strongly interact with the hydrophobic, low-solubility drug and yet are still significantly water soluble. Typically good results

are obtained when the aqueous solubility of the copolymer at 37°C is between about 0.04 and 10.0 wt% and preferably between about 0.1 and 2.0 wt%.

While specific polymers are discussed as being suitable for use in the compositions of the present invention, blends of such polymers may also be suitable. Thus the term "polymer" is intended to include blends of polymers in addition to a single species of polymer.

CONCENTRATION ENHANCEMENT

As described above, it is often preferred that the neutral dispersion polymer is also concentration-enhancing. The neutral concentration-enhancing dispersion polymers improve the concentration of a low-solubility drug in a use environment, and thereby preferably improve bioavailability of the drug.

The term "concentration-enhancing" means that the polymer is present in a sufficient amount in the composition so as to improve the concentration of the drug in a use environment relative to a control composition free from the concentration-enhancing polymer. As used herein, a "use environment" can be either the *in vivo* environment of the GI tract, subdermal, intranasal, buccal, intrathecal, ocular, intraaural, subcutaneous spaces, vaginal tract, arterial and venous blood vessels, pulmonary tract or intramuscular tissue of an animal, such as a mammal and particularly a human, or the *in vitro* environment of a test solution, such as phosphate buffered saline (PSB) or a Model Fasted Duodenal (MFD) solution. Concentration enhancement may be determined through either *in vitro* dissolution tests or through *in vivo* tests. It has been determined that enhanced drug concentration in *in vitro* dissolution tests in Model Fasted Duodenal (MFD) solution or Phosphate Buffered Saline (PBS) is a good indicator of *in vivo* performance and bioavailability. An appropriate PBS solution is an aqueous solution comprising 20 mM sodium phosphate (Na_2HPO_4), 47 mM potassium phosphate (KH_2PO_4), 87 mM NaCl, and 0.2 mM KCl, adjusted to pH 6.5 with NaOH. An appropriate MFD solution is the same PBS solution wherein additionally is present 7.3 mM sodium taurocholic acid and 1.4 mM of 1-palmitoyl-2-oleyl-sn-glycero-3-phosphocholine. In particular, a composition containing a concentration-enhancing polymer may be dissolution-tested by adding it to MFD or PBS solution and agitating to promote dissolution.

In one aspect, a composition containing a concentration-enhancing polymer of the present invention provides a Maximum Drug Concentration

(MDC) that is at least 1.25-fold the MDC provided by a control composition. In other words, if the MDC provided by the control composition is 100 μ g/mL, then a composition of the present invention containing a concentration-enhancing polymer provides an MDC of at least 125 μ g/mL. More preferably, the MDC of drug achieved with the compositions of the present invention are at least 2-fold, even more preferably at least 3-fold, and most preferably at least 10-fold that of the control composition.

The appropriate control composition depends on whether a dispersion polymer is being tested, or whether a composition comprised of a mixture of a dispersion and concentration-enhancing polymer is being tested. Where the composition is simply the dispersion itself, and the dispersion polymer is being tested to determine whether it is a concentration-enhancing polymer, the control composition is conventionally the undispersed drug alone (e.g., typically, the crystalline drug alone in its most thermodynamically stable crystalline form, or in cases where a crystalline form of the drug is unknown, the control may be the amorphous drug alone) or the drug plus a weight of inert diluent equivalent to the weight of polymer in the test composition. By inert is meant that the diluent is not concentration-enhancing. Where the composition is comprised of a mixture of a dispersion and additional concentration-enhancing polymer, the control composition is the dispersion alone without any additional concentration-enhancing polymer.

Alternatively, the compositions containing concentration-enhancing polymers of the present invention provide in an aqueous use environment a concentration versus time Area Under The Curve (AUC), for any period of at least 90 minutes between the time of introduction into the use environment and about 270 minutes following introduction to the use environment that is at least 1.25-fold that of an appropriate control composition. More preferably, the AUC achieved with the compositions of the present invention are at least 2-fold, more preferably at least 3-fold, and most preferably at least 10-fold that of a control composition.

Alternatively, the compositions of the present invention containing concentration-enhancing polymers, when dosed orally to a human or other animal, provide an AUC in drug concentration in the blood plasma or serum that is at least 1.25-fold that observed when an appropriate control composition is dosed. Preferably, the blood AUC is at least 2-fold, more preferably at least

3-fold, that of an appropriate control composition. Thus, the compositions that, when evaluated, meet either the *in vitro* or the *in vivo*, or both, performance criteria are a part of this invention.

A typical *in vitro* test to evaluate enhanced drug concentration can be conducted by (1) administering with agitation a sufficient quantity of test composition (for example, the dispersion of the low-solubility drug and neutral polymer) in a test medium, such that if all of the drug dissolved, the theoretical concentration of drug would exceed the equilibrium concentration of the drug by a factor of at least 2; (2) adding an appropriate amount of control composition to an equivalent amount of test medium; and (3) determining whether the measured MDC and/or AUC of the test composition in the test medium is at least 1.25-fold that provided by the control composition. The concentration of dissolved drug is typically measured as a function of time by sampling the test medium and plotting drug concentration in the test medium vs. time so that the MDC and/or AUC can be ascertained. In conducting such a dissolution test, the amount of test composition or control composition used is an amount such that if all of the drug dissolved, the drug concentration would be at least 2-fold to 100-fold that of the solubility (that is, the equilibrium concentration), of the drug. For some dispersions of a very low-solubility drug and neutral polymer, it may be necessary to administer an even greater amount of the dispersion to determine the MDC.

To avoid drug particulates which would give an erroneous determination, the test solution is either filtered or centrifuged. "Dissolved drug" is typically taken as that material that either passes a 0.45 μm syringe filter or, alternatively, the material that remains in the supernatant following centrifugation. Filtration can be conducted using a 13 mm, 0.45 μm polyvinylidene difluoride syringe filter sold by Scientific Resources under the trademark TITAN®. Centrifugation is typically carried out in a polypropylene microcentrifuge tube by centrifuging at 13,000 G for 60 seconds. Other similar filtration or centrifugation methods can be employed and useful results obtained. For example, using other types of microfilters may yield values somewhat higher or lower (± 10 -40%) than that obtained with the filter specified above but will still allow identification of preferred dispersions. It is recognized that this definition of "dissolved drug" encompasses not only monomeric solvated drug molecules but also a wide range of species such as polymer/drug assemblies

that have submicron dimensions such as drug aggregates, aggregates of mixtures of polymer and drug, micelles, polymeric micelles, colloidal particles or nanocrystals, polymer/drug complexes, and other such drug-containing species that are present in the filtrate or supernatant in the specified dissolution test.

5 Alternatively, the concentration-enhancing polymer results in improved bioavailability. Relative bioavailability of the drug in the compositions of the present invention can be tested *in vivo* in animals or humans using conventional methods for making such a determination. An *in vivo* test, such as a crossover study, may be used to determine whether a composition provides an enhanced
10 relative bioavailability compared with a control. In an *in vivo* crossover study a "test composition" of drug and concentration-enhancing polymer is dosed to half a group of test subjects and, after an appropriate washout period (e.g., one week) the same subjects are dosed with a "control composition" that comprises an equivalent quantity of drug as the "test composition." The other half of the
15 group is dosed with the control composition first, followed by the test composition. The relative bioavailability is measured as the concentration in the blood (serum or plasma) versus time area under the curve (AUC) determined for the test composition divided by the AUC in the blood provided by the control composition. Preferably, this test/control ratio is determined for each subject,
20 and then the ratios are averaged over all subjects in the study. *In vivo* determinations of AUC can be made by plotting the serum or plasma concentration of drug along the ordinate (y-axis) against time along the abscissa (x-axis). Generally, the values for AUC represent a number of values taken from all of the subjects in a patient test population averaged over the entire test
25 population.

A preferred embodiment of the invention is one in which the relative bioavailability of the test composition is at least 1.25 relative to a control composition as described above. (That is, the AUC in the blood provided by the test composition is at least 1.25-fold the AUC provided by the control
30 composition.) An even more preferred embodiment of the invention is one in which the relative bioavailability of the test composition is at least 2.0, more preferably at least 3, relative to a control composition of the drug but with no polymer present, as described above. The determination of AUCs is a well-known procedure and is described, for example, in Welling, "Pharmacokinetics
35 Processes and Mathematics," ACS Monograph 185 (1986).

Often the enhancement in drug concentration or relative bioavailability that is observed increases as the drug:polymer ratio decreases from a value of about 1 (50 wt% drug) to a value of about 0.11 (10 wt% drug). The drug:polymer ratio that yields optimum results varies from drug to drug and is best determined in *in vitro* dissolution tests and/or *in vivo* bioavailability tests. However, the amount of polymer that can be used in a dosage form is often limited by the total mass requirements of the dosage form. For example, when oral dosing to a human is desired, at low drug-to-polymer ratios the total mass of drug and polymer may be unacceptably large for delivery of the desired dose in a single tablet or capsule. Thus, it is often necessary to use drug-to-polymer ratios that are less than optimum in specific dosage forms to provide a sufficient drug dose in a dosage form that is small enough to be easily delivered to a use environment.

IMPROVED CHEMICAL STABILITY

When the compositions of the present invention comprise dispersions of acid-sensitive drug and neutral polymer(s), such compositions provide improved chemical stability of the drug relative to a control composition comprised of an equivalent quantity of drug dispersed in an acidic dispersion polymer. The acidic dispersion polymer used in the control composition may be any polymer suitable for determining whether the drug is acid-sensitive that meets the definition of an "acidic polymer." Convenient acidic dispersion polymers for use in the control composition include HPMCAS, CAP and HPMCP. A composition of the present invention will meet the chemical stability requirements discussed below relative to at least one control dispersion of drug and either HPMCAS, HPMCP, or CAP.

A relative degree of improvement in chemical stability of an acid-sensitive drug in a test composition may be determined by taking the ratio of the degree of degradation of the drug in a control composition and the degree of degradation of the drug in a test composition under the same storage conditions for the same storage time period. The test composition is simply the composition of acid-sensitive drug, neutral polymer and optional additional excipients of the present invention. The control composition is the same as the test composition with the exception that an acidic dispersion polymer (that is, either HPMCAS, HPMCP or CAP) replaces the neutral dispersion polymer. For

example, where the degree of degradation of a drug in a test composition comprised of an acid-sensitive drug and a neutral polymer is 1 wt%, and the degree of degradation of a control composition comprised of an equivalent quantity of drug and HPMCAS is 5 wt%, the relative degree of improvement is 5 wt%/1 wt% equals 5.0. For dispersions of acid-sensitive drugs and neutral polymers of the present invention, the relative degree of improvement is at least 1.25. Preferably, the relative degree of improvement is at least 3, more preferably at least 5, and even more preferably at least 10. In fact, some compositions of the present invention may achieve a relative degree of improvement greater than 100.

The particular storage conditions and time of storage may be chosen as convenient depending on the degree of acid-sensitivity of the drug, the particular acidic polymer used in the control dispersion, and the ratio of drug to polymer in the dispersion. Where the drug is particularly acid-sensitive, or where the dispersion has a low ratio of drug to polymer, then shorter storage time periods may be used. Where the rate of drug degradation is linear, the relative degree of improvement will be independent of the storage time. However, where the rate of drug degradation is non-linear under controlled storage conditions, the stability test used to compare the test dispersion with the control dispersion is preferably chosen such that the degree of degradation is sufficiently large that it may be accurately measured. Typically, the time period is chosen so as to observe a degree of degradation in the control composition of at least 0.1 wt% to 0.2 wt%. However, the time period is not so long that the ratio of drug to polymer changes substantially. Typically, the time period is such that the observed degree of degradation for the test composition is less than 50 wt% and preferably less than 20 wt%. When rate of drug degradation in the control composition is relatively slow, the test is preferably conducted over a long enough period of time under controlled storage conditions to allow a meaningful comparison of the stability of the test dispersion with the control dispersion.

A stability test which may be used to test whether a dispersion meets the chemical stability criteria described above is storage of the test dispersion and the control dispersion for six months at 40°C and 75% RH. A relative degree of improvement may become apparent within a shorter time, such as three to five days, and shorter storage times may be used for some very acid-sensitive

drugs. When comparing dispersions under storage conditions which approximate ambient conditions, e.g., 30°C and 60% RH, the storage period may need to be several months up to two years.

5 In addition, it is preferred that the dispersions comprising acid-sensitive drug and neutral polymer(s) result in drug stability such that the acid-sensitive drug has a degree of degradation of less than about 2 wt%, more preferably less than about 0.5 wt%, and most preferably less than about 0.1 wt% when stored at 40°C and 75% RH for six months, or less than 2 wt%, more preferably 0.5 wt%, and more preferably 0.1 wt%, when stored at 30°C and 60% RH for
10 one year. Nevertheless, the compositions of the present invention may have a degree of degradation that is much greater than the preferred values, so long as the dispersion achieves the degree of improvement relative to a control composition as described above.

PREPARATION OF DISPERSIONS

15 Dispersions of the drugs and neutral dispersion polymer may be made according to any known process which results in at least a "major portion" (meaning at least 60 wt%) of the drug being in the amorphous state. Such processes include mechanical, thermal and solvent processes. Exemplary mechanical processes include milling and extrusion; melt processes include
20 high temperature fusion, solvent modified fusion and melt-congeal processes; and solvent processes include non-solvent precipitation, spray coating and spray-drying. See, for example, U.S. Patent No. 5,456,923 and U.S. Patent No. 5,939,099 which describe formation of dispersions via extrusion processes; U.S. Patent No. 5,340,591 and U.S. Patent No. 4,673,564 which describe forming
25 dispersions by milling processes; and U.S. Patent No. 5,707,646 and U.S. Patent No. 4,894,235 which describe the formation of dispersions via melt/congeal processes, the disclosures of which are incorporated by reference.

While the drug in its pure state may be crystalline or amorphous, at least a major portion of the drug in the dispersion is amorphous. By "amorphous" is
30 meant simply that the drug is in a non-crystalline state. As used herein, the term "a major portion" of the drug means that at least 60% of the drug in the dispersion is in the amorphous form, rather than the crystalline form. In general, a drug is more reactive in its amorphous state relative to its crystalline state and so the need to use a neutral dispersion polymer to prevent degradation of acid-
35 sensitive drug increases as the fraction of drug in the amorphous state

increases. It has also been found that the aqueous concentration of the drug in a use environment tends to improve as the fraction of drug present in the amorphous state in the dispersion increases. Accordingly, a "major portion" of the drug in the dispersion is amorphous and preferably the drug in the dispersion is substantially amorphous. As used herein, a "major portion" and "substantially amorphous" mean that the amount of the drug in crystalline form does not exceed about 40 wt% and about 25 wt%, respectively. More preferably, the drug in the dispersion is "almost completely amorphous," meaning that the amount of drug in the crystalline form does not exceed 10 wt%. Amounts of crystalline drug may be measured by powder X-ray diffraction, Scanning Electron Microscope (SEM) analysis, differential scanning calorimetry (DSC), or any other standard quantitative measurement.

The amorphous drug can exist as a pure phase, as a solid solution of drug homogeneously distributed throughout the polymer or any combination of these states or those states that lie intermediate between them. In cases where the drug is a low-solubility drug and concentration or bioavailability enhancement is desired, the dispersion is preferably "substantially homogeneous" so that the amorphous drug is dispersed as homogeneously as possible throughout the polymer. Dispersions of the present invention that are substantially homogeneous generally are more physically stable and have improved concentration-enhancing properties and, in turn improved bioavailability, relative to nonhomogeneous dispersions. As used herein, "substantially homogeneous" means that the drug present in relatively pure amorphous domains within the solid dispersion is relatively small, on the order of less than 20%, and preferably less than 10% of the total amount of drug.

While the dispersion may have some drug-rich domains, it is preferred that the dispersion itself have a single glass transition temperature (T_g) which demonstrates that the dispersion is substantially homogeneous. This contrasts with a simple physical mixture of pure amorphous drug particles and pure amorphous polymer particles which generally display two distinct T_g s, one that of the drug and one that of the polymer. T_g as used herein is the characteristic temperature where a glassy material, upon gradual heating, undergoes a relatively rapid (e.g., 10 to 100 seconds) physical change from a glass state to a rubber state. Almost completely amorphous, substantially homogeneous dispersions may be made by any of the processes discussed above.

To obtain the maximum level of concentration and bioavailability enhancement, particularly upon storage for long times prior to use, it is preferred that the drug remain, to the extent possible, in the amorphous state. The inventors have found that this is best achieved when the glass-transition temperature, T_g , of the amorphous dispersion is substantially above the storage temperature of the dispersion. In particular, it is preferable that the T_g of the amorphous dispersion is at least 40°C and preferably at least 60°C. Since the T_g is a function of the water content of the dispersion which in turn is a function of the RH to which the dispersion is exposed, these T_g values refer to the T_g of the dispersion containing water in an amount that is in equilibrium with the RH equivalent to that found during storage. For those aspects of the invention in which the dispersion is a solid, substantially amorphous dispersion of drug in the dispersion polymer and in which the drug itself has a relatively low T_g (about 70°C or less) it is preferred that the dispersion polymer have a T_g of at least 40°C, preferably at least 70°C and more preferably greater than 100°C. Exemplary high T_g polymers include hydroxypropyl cellulose and hydroxypropyl methyl cellulose. Since conversion of amorphous drug to the crystalline state is related to the relative values of (1) the T_g of the dispersion (at the storage RH) and (2) the storage temperature, solid amorphous dispersions of the present invention may tend to remain in the amorphous state for longer periods when stored at relatively low temperatures and low relative humidities. In addition, packaging of such dispersions so as to prevent absorption of water or inclusion of a water absorbing material such as a dessicant to also prevent or retard water absorption can lead to a higher T_g during storage, thereby helping to retain the amorphous state. Likewise, storage at lower temperatures can also improve the retention of the amorphous state.

Turning now to particular methods, when either the neutral polymer or the drug has a relatively low melting point, typically less than about 200°C and preferably less than about 160°C, extrusion or melt-congeal processes that provide heat and/or mechanical energy are often suitable for forming almost completely amorphous dispersions. Often, when the drug has significant solubility in the dispersion material, such methods may also make substantially homogeneous dispersions. For example, 10 wt% drug and 90 wt% polyvinyl pyrrolidone (PVP) may be dry blended, with or without the addition of water, and the blend fed to a twin-screw extrusion device. The processing temperature

may vary from about 50°C up to about 200°C depending on the melting point of the drug and polymer, which is a function of the polymer grade chosen and the amount of water, if any, added. Generally, the higher the melting point of the drug and polymer, the higher the processing temperature. Generally, the lowest
5 processing temperature that produces a satisfactory dispersion (almost completely amorphous and substantially homogeneous) is chosen.

A method for forming dispersions that is often preferred is "solvent processing," which consists of dissolution of the drug and one or more neutral polymers in a common solvent. The term "solvent" is used broadly and includes
10 mixtures of solvents. "Common" here means that the solvent, which can be a mixture of compounds, will simultaneously dissolve both the drug and the polymer(s).

After both the drug and polymer(s) have been dissolved, the solvent is removed by evaporation or by mixing with a non-solvent. Exemplary processes
15 are spray-drying, spray-coating (pan-coating, fluidized bed coating, etc.), vacuum evaporation, precipitation by mixing of the polymer and drug solution with a non-solvent such as CO₂, hexane, or heptane, and processes that combine solvents with heat and mechanical energy such as extrusion processes that include the use of a solvent as a processing aid or even wet
20 granulation processes as long as such processes result in at least a major portion of the drug being amorphous. Preferably, removal of the solvent is rapid and results in a solid dispersion which is substantially homogeneous. In substantially homogeneous dispersions, the drug is dispersed as homogeneously as possible throughout the polymer and can be thought of as a
25 solid solution of drug dispersed in the polymer(s). When the resulting dispersion constitutes a solid solution of drug in polymer, the dispersion may be thermodynamically stable, meaning that the concentration of drug in the polymer is at or below its equilibrium value, or it may be considered a supersaturated solid solution where the drug concentration in the dispersion
30 polymer(s) is above its equilibrium value.

The solvent may be removed through the process of spray-drying. The term spray-drying is used conventionally and broadly refers to processes involving breaking up liquid mixtures into small droplets (atomization) and rapidly removing solvent from the mixture in a container (spray-drying
35 apparatus) where there is a strong driving force for evaporation of solvent from

the droplets. The strong driving force for solvent evaporation is generally provided by maintaining the partial pressure of solvent in the spray-drying apparatus well below the vapor pressure of the solvent at the temperature of the drying droplets. This is accomplished by either (1) maintaining the pressure in the spray-drying apparatus at a partial vacuum (e.g., 0.01 to 0.50 atm); (2) mixing the liquid droplets with a warm drying gas; or (3) both. In addition, at least a portion of the heat required for evaporation of solvent may be provided by heating the spray solution.

Solvents suitable for spray-drying can be any organic compound in which the drug and polymer are mutually soluble. Preferably, the solvent is also volatile with a boiling point of 150°C or less. In addition, the solvent should have relatively low toxicity and be removed from the dispersion to a level that is acceptable according to The International Committee on Harmonization (ICH) guidelines. Removal of solvent to this level may require a processing step such as tray-drying or fluidized-bed drying subsequent to the spray-drying or spray-coating process. Preferred solvents include alcohols such as methanol, ethanol, n-propanol, iso-propanol, and butanol; ketones such as acetone, methyl ethyl ketone and methyl iso-butyl ketone; esters such as ethyl acetate and propylacetate; and various other solvents such as acetonitrile, methylene chloride, toluene, and 1,1,1-trichloroethane. Lower volatility solvents such as dimethyl acetamide or dimethylsulfoxide can also be used. Mixtures of solvents, such as 50% methanol and 50% acetone, can also be used, as can mixtures with water as long as the polymer and drug are sufficiently soluble to make the spray-drying process practicable. In particular, when the neutral polymer is relatively hydrophilic such as hydroxypropyl methyl cellulose (HPMC), or certain amphiphilic hydroxyl-functional vinyl copolymers, it is often preferred to use a mixture of an organic solvent and water to ensure that both drug and polymer are simultaneously soluble.

Generally, the temperature and flow rate of the drying gas is chosen so that the polymer/drug-solution droplets are dry enough by the time they reach the wall of the apparatus that they are essentially solid, and so that they form a fine powder and do not stick to the apparatus wall. The actual length of time to achieve this level of dryness depends on the size of the droplets. Droplet sizes generally range from 1 μ m to 500 μ m in diameter, with 5 to 200 μ m being more typical. The large surface-to-volume ratio of the droplets and the large driving

force for evaporation of solvent leads to actual drying times of a few seconds or less, and more typically less than 0.1 second. This rapid drying is often critical to the particles maintaining a uniform, homogeneous dispersion instead of separating into drug-rich and polymer-rich phases. As above, to get large
5 enhancements in concentration and bioavailability it is often necessary to obtain as homogeneous of a dispersion as possible. Solidification times should be less than 100 seconds, preferably less than a few seconds, and more preferably less than 1 second. In general, to achieve this rapid solidification of the drug/polymer solution, it is preferred that the size of droplets formed during the
10 spray-drying process are less than about 500 μm in diameter, and preferably less than about 300 μm . The resultant solid particles thus formed are generally less than about 300 μm in diameter.

Following solidification, the solid powder typically stays in the spray-drying chamber for about 5 to 60 seconds, further evaporating solvent from the
15 solid powder. The final solvent content of the solid dispersion as it exits the dryer should be low, since this reduces the mobility of drug molecules in the dispersion, thereby improving its stability. Generally, the solvent content of the dispersion as it leaves the spray-drying chamber should be less than about 10 wt% and preferably less than about 3 wt%. In some cases, it may be
20 preferable to spray a solvent or a solution of a polymer or other excipient into the spray-drying chamber to form granules, so long as the dispersion is not adversely affected.

Spray-drying processes and spray-drying equipment are described generally in Perry's *Chemical Engineers' Handbook*, Sixth Edition (R. H. Perry, D. W. Green, J. O. Maloney, eds.) McGraw-Hill Book Co. 1984, pages 20-54 to
25 20-57. More details on spray-drying processes and equipment are reviewed by Marshall "Atomization and Spray-Drying," 50 *Chem. Eng. Prog. Monogr. Series* 2 (1954).

The solid amorphous dispersions formed from a low-solubility drug and
30 an amphiphilic hydroxyl-functional vinyl copolymer are preferably almost completely amorphous and substantially homogenous. Such almost completely amorphous, substantially homogenous dispersions are generally more physically stable (that is, resist phase separation or crystallization upon storage), and show higher levels of concentration enhancement. As with the
35 dispersions of other embodiments of the invention, such dispersions may in

principle be formed by any known process, including mechanical, thermal and solvent processes. However, "solvent processing" is preferred for forming dispersions of amphiphilic hydroxyl-functional vinyl copolymers because when properly performed, solvent processing tends to produce dispersions in which the drug is almost completely amorphous and in which the dispersion is substantially homogeneous.

In order to achieve such preferred dispersions, a copolymer type and solvent must be chosen to yield a solution in which the amphiphilic, hydroxyl-functional vinyl copolymer and low-solubility drug are both soluble and preferably highly soluble. Preferably, both the copolymer and low-solubility drug each have solubilities in the solvent at 37°C of at least 0.5 wt% and preferably at least 2.0 wt% and more preferably at least 5.0 wt%.

In order to have the drug and copolymer both have such high solubilities in the solvent while the polymer is also sufficiently soluble in aqueous solution, the polymer molecular weight must generally not be too high and the polymer must have the correct n/m values. For example it has been found that for vinylacetate/vinylalcohol copolymers that n/m values are preferably between about 2 and 100 and molecular weights are preferably less than about 50,000 daltons.

In addition, the solvent is preferably chosen such that the lesser of either (1) the polymer solubility or (2) the product of the drug solubility and the ratio of the polymer to drug in the final dispersion, is high and is preferably maximized. To achieve this, it is often desirable to use blends of two or more solvents. In particular, a blend of water and an organic solvent is often preferred. Exemplary preferred solvents are mixtures of water and alcohols such as methanol, ethanol, n-propanol, iso-propanol and the isomers of butanol.

In addition it is preferable for the solvent processing to be conducted such that solvent is removed rapidly so that the polymer-drug solution solidifies rapidly. Thus solvent processes such as spray-coating and spray-drying are preferred.

In addition, to achieve rapid solvent removal, and to keep the residual solvent level in the dispersion low (preferably less than about 5 wt%), a relatively volatile solvent is chosen. Preferably the boiling points of the solvents are less than about 150°C and more preferably less than about 125°C. When the solvent is a mixture of solvents, up to about 40% of the solvent may

comprise a low volatility solvent: When the dispersion polymer is a vinylacetate/vinylalcohol copolymer of the preferred type described earlier, preferred solvents include mixtures of methanol and water.

Returning to dispersions generally, the amount of polymer relative to the amount of drug present in the dispersions of the present invention depends on the drug and polymer and may vary widely from a drug-to-polymer weight ratio of from 0.01 to about 4 (e.g., 1 wt% drug to 80 wt% drug). However, in most cases it is preferred that the drug-to-polymer ratio is greater than about 0.05 (4.8 wt% drug) and less than about 2.5 (71 wt% drug).

In addition to the drug and polymer(s), the dispersions of the present invention may include optional additional ingredients. One optional additional ingredient is a buffer. Buffers suitable for use in the dispersions of the present invention are preferably basic. Buffers are particularly preferred for dispersions of neutral polymers and drugs that have a high degree of acid-sensitivity. The buffers may reduce the risk of drug degradation due to the presence of acidic species in the dispersion, elsewhere in the composition, or acidic or basic species formed during storage. The compositions preferably provide improved chemical stability relative to dispersions that are free from the buffer.

Exemplary buffers that may be used in the dispersions of the present invention include sodium acetate, ammonium acetate, sodium carbonate, sodium bicarbonate, disodium hydrogen phosphate and trisodium phosphate. It is generally preferred that the buffer be chosen such that it maintains the pH at the optimum value during storage to minimize degradation reactions. For most acid-sensitive drugs, the preferred pH value is greater than about 6. Since many dispersion polymers such as cellulose are also base-sensitive, it is generally preferred that the buffer be chosen to maintain the dispersion pH between about 6 and 10. In cases where the acid-sensitive drug is also a base-sensitive drug, it is often preferred to maintain the dispersion pH between about 6 and about 8. To accomplish this, it is sometimes desirable for the buffer to be a blend of protonated and deprotonated material, where the pK_a of the protonated material is near the desired dispersion pH. Such buffers may comprise from 0.1 to 20 wt% of the dispersion.

Alternatively, the dispersion may include basic excipients such as mono-, di- or tri-ethanolamine, or glucoseamine, which may comprise from 0.1

to 20 wt% of the dispersion. Such basic excipients may be used to control the dispersion pH as described above.

MIXTURES OF DISPERSIONS AND CONCENTRATION-ENHANCING POLYMER

5 A separate embodiment of the invention comprises a combination of
(1) a dispersion of an acid-sensitive, low-solubility drug and neutral dispersion
polymer(s) and (2) a second polymer that is concentration-enhancing. The
dispersion is free from at least a portion of the second polymer, and is
preferably substantially free from the second polymer. By "substantially free" is
10 meant that the dispersion and the second polymer are not mixed at the
molecular level to form a dispersion containing the second polymer. While a
small amount of the second polymer may be incorporated into the dispersion as
a result of processing, nevertheless the drug in the dispersion meets the
chemical stability requirements discussed above for dispersions of acid-
15 sensitive drugs and neutral polymers notwithstanding the presence of the
second polymer in the composition. The neutral dispersion polymer optionally
may also be concentration-enhancing.

 The compositions of this aspect of the present invention are generally
physical combinations comprising the dispersion and the second polymer.
20 "Combination" as used herein means that the dispersion and second polymer
may be in physical contact with each other or in close proximity but without the
necessity of being physically mixed at the molecular level (i.e., a dispersion).
The dispersion and concentration-enhancing polymer may be in different
regions of the composition. For example, the solid composition may be in the
25 form of a multi-layer tablet, as known in the art, wherein one or more layers
comprises the dispersion and one or more different layers comprises the
second polymer. Yet another example may constitute a coated tablet wherein
either the dispersion or the second polymer or both may be present in the tablet
core and the coating may comprise the second polymer. Alternatively, the
30 combination can be in the form of a simple dry physical mixture wherein both
the dispersion and the second polymer are mixed in particulate form and
wherein the particles of each, regardless of size, retain the same individual
physical properties that they exhibit in bulk. Any conventional method used to
mix the dispersion and second polymer together such as physical mixing and

dry or wet granulation, which does not convert the dispersion and the second polymer to another molecular dispersion, may be used.

Alternatively, the dispersion and second polymer may be co-administered, meaning that the dispersion may be administered separately from, but within the same general time frame as, the second polymer. Thus, a dispersion may, for example, be administered in its own dosage form which is taken at approximately the same time as the second polymer which is in a separate dosage form. If administered separately, it is generally preferred to administer both the dispersion and the second polymer within 60 minutes of each other, so that the two are present together in the use environment. When not administered approximately simultaneously (e.g., within a minute or two of each other), the second polymer is preferably administered prior to the dispersion.

The second polymer may be any concentration-enhancing polymer, such as those described above in connection with the neutral dispersion polymers. As described above, the inventors have found that ionizable, cellulosic polymers, particularly those that are acidic cellulosic enteric polymers, provide superior enhancement in aqueous concentration of the drug in a use environment relative to other polymers, and are therefore preferred in the absence of their reactivity with the drug. Many of these ionizable, cellulosic polymers have acidic functional groups and therefore are inappropriate for use as a dispersion polymer with acid-sensitive drugs. However, the concentration-enhancing advantage provided by such ionizable concentration-enhancing polymers may be achieved by simply combining such a polymer as the second polymer with a pre-formed dispersion of the acid-sensitive drug and a neutral polymer in a fashion that does not alter the neutral characteristic of the pre-formed dispersion.

A preferred class of concentration-enhancing polymers suitable for use as the second polymer comprises ionizable cellulosic polymers with at least one ester- and/or ether- linked substituent in which the polymer has a degree of substitution of at least 0.05 for each substituent. Particularly preferred hydrophilic substituents for use in making ionizable polymers are those that are ether- or ester-linked acidic ionizable groups such as carboxylic acids, thiocarboxylic acids and sulfonates, and groups that may be non-acidic such as substituted phenoxy groups, amines, and phosphates. Exemplary ether-linked

ionizable substituents include: carboxylic acids, such as acetic acid, propionic acid, benzoic acid, salicylic acid, alkoxybenzoic acids such as ethoxybenzoic acid or propoxybenzoic acid, the various isomers of alkoxyphthalic acid such as ethoxyphthalic acid and ethoxyisophthalic acid, the various isomers of
5 alkoxy nicotinic acid such as ethoxy nicotinic acid, and the various isomers of picolinic acid such as ethoxypicolinic acid, etc.; thiocarboxylic acids, such as thioacetic acid; substituted phenoxy groups, such as hydroxyphenoxy, etc.; amines, such as aminoethoxy, diethylaminoethoxy, trimethylaminoethoxy, etc.; phosphates, such as phosphate ethoxy; and sulfonates, such as sulphonate
10 ethoxy. Exemplary ester linked ionizable substituents include: carboxylic acids, such as succinate, citrate, phthalate, terephthalate, isophthalate, trimellitate, and the various isomers of pyridinedicarboxylic acid, etc.; thiocarboxylic acids, such as thiosuccinate; substituted phenoxy groups, such as amino salicylic acid; amines, such as natural or synthetic amino acids, such as alanine or
15 phenylalanine; phosphates, such as acetyl phosphate; and sulfonates, such as acetyl sulfonate. For aromatic-substituted polymers to also have the requisite aqueous solubility, it is also desirable that sufficient hydrophilic groups such as hydroxypropyl or carboxylic acid functional groups be attached to the polymer to render the polymer aqueous soluble at least at pH values where any ionizable
20 groups are ionized. In some cases, the aromatic group may itself be ionizable, such as phthalate or trimellitate substituents.

Exemplary ionizable cellulosic concentration-enhancing polymers that are at least partially ionized at physiologically relevant pHs include: hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl
25 cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, hydroxyethyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, hydroxyethyl methyl cellulose acetate succinate, hydroxyethyl methyl cellulose acetate phthalate, carboxyethyl cellulose, carboxymethyl cellulose, cellulose acetate phthalate, methyl cellulose
30 acetate phthalate, ethyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, hydroxypropyl methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, hydroxypropyl methyl cellulose acetate succinate phthalate, hydroxypropyl methyl cellulose succinate phthalate, cellulose propionate phthalate, hydroxypropyl cellulose butyrate
35 phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate,

ethyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate, hydroxypropyl methyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate succinate, cellulose propionate trimellitate, cellulose butyrate trimellitate, cellulose acetate terephthalate, cellulose acetate isophthalate, cellulose acetate pyridinedicarboxylate, salicylic acid cellulose acetate, hydroxypropyl salicylic acid cellulose acetate, ethylbenzoic acid cellulose acetate, hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl phthalic acid cellulose acetate, ethyl nicotinic acid cellulose acetate, and ethyl picolinic acid cellulose acetate.

Another class of concentration-enhancing polymers suitable for use with the present invention comprises ionizable non-cellulosic polymers. Exemplary polymers of this type that are generally acidic include carboxylic acid-functionalized vinyl polymers, such as the carboxylic acid functionalized polymethacrylates and carboxylic acid functionalized polyacrylates such as the EUDRAGITS® manufactured by Rohm Tech Inc., of Malden, Massachusetts, and acidic proteins.

A preferred class of concentration-enhancing polymers comprises polymers that are "amphiphilic" in nature, meaning that the polymer has hydrophobic and hydrophilic portions. The hydrophobic portion may comprise groups such as aliphatic or aromatic hydrocarbon groups. The hydrophilic portion may comprise either ionizable or non-ionizable groups that are capable of hydrogen bonding such as hydroxyls, carboxylic acids, esters, amines or amides.

Amphiphilic polymers are preferred because it is believed that such polymers may tend to have relatively strong interactions with the drug and may promote the formation of various types of polymer/drug assemblies in the use environment. Ionizable polymers, and particularly those that have a significant level of carboxylic acid functional groups (*e.g.*, acidic polymers) typically show the greatest concentration and bioavailability enhancements and therefore are preferred for use with low-solubility drugs. However, they may not form a part of the dispersion when the drug is acid-sensitive. Thus, a particularly preferred composition is one in which an acid-sensitive drug is dispersed in a neutral or non-ionizable polymer and an ionizable or acidic polymer is added such that it is not part of the dispersion.

In addition, ionic, and in particular acidic polymers, are thought to show desirable concentration-enhancement properties due to the repulsion of the like charges of the ionized groups of such polymers. This repulsion may serve to limit the size of the polymer/drug assemblies to the nanometer or submicron scale. For example, while not wishing to be bound by a particular theory, such polymer/drug assemblies may comprise hydrophobic drug clusters surrounded by the polymer with the polymer's hydrophobic regions turned inward towards the drug and the hydrophilic regions of the polymer turned outward toward the aqueous environment. Alternatively, depending on the specific chemical nature of the drug, the ionized functional groups of the polymer may associate, for example, via ion pairing or hydrogen bonds, with ionic or polar groups of the drug. In the case of ionizable polymers, the hydrophilic regions of the polymer would include the ionized functional groups. Such polymer/drug assemblies in solution may well resemble charged polymeric micellar-like structures. In any case, regardless of the mechanism of action, the inventors have observed that such amphiphilic polymers, particularly ionizable cellulosic polymers, have been shown to improve the maximum drug concentration (MDC) and/or area under the curve (AUC) of the drug in aqueous solution relative to control compositions free from such polymers.

Surprisingly, such amphiphilic polymers can greatly enhance the maximum concentration of drug obtained when the drug is dosed to a use environment. In addition, such amphiphilic polymers may interact with the drug to prevent the precipitation or crystallization of the drug from solution despite its concentration being substantially above its equilibrium concentration.

Exemplary ionizable cellulosic polymers that meet the definition of amphiphilic, having hydrophilic and hydrophobic regions, include polymers such as cellulose acetate phthalate and cellulose acetate trimellitate where the cellulosic repeat units that have one or more acetate substituents are hydrophobic relative to those that have no acetate substituents or have one or more ionized phthalate or trimellitate substituents.

A particularly desirable subset of cellulosic ionizable concentration-enhancing polymers are those that possess both a carboxylic acid functional aromatic substituent and an alkylate substituent and thus are amphiphilic. Exemplary polymers include cellulose acetate phthalate, methyl cellulose acetate phthalate, ethyl cellulose acetate phthalate, hydroxypropyl cellulose

acetate phthalate, hydroxypropyl methyl cellulose phthalate, hydroxypropyl methyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, cellulose propionate phthalate, hydroxypropyl cellulose butyrate phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate, 5 ethyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate, hydroxypropyl methyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate succinate, cellulose propionate trimellitate, cellulose butyrate trimellitate, cellulose acetate terephthalate, cellulose acetate isophthalate, cellulose acetate pyridinedicarboxylate, salicylic acid cellulose acetate, 10 hydroxypropyl salicylic acid cellulose acetate, ethylbenzoic acid cellulose acetate, hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl phthalic acid cellulose acetate, ethyl nicotinic acid cellulose acetate, and ethyl picolinic acid cellulose acetate.

Another particularly desirable subset of cellulosic ionizable polymers are 15 those that possess a non-aromatic carboxylate substituent. Exemplary polymers include hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, and hydroxyethyl cellulose acetate succinate.

20 Non-cellulosic polymers that are amphiphilic are copolymers of a relatively hydrophilic and a relatively hydrophobic monomer. Examples include acrylate and methacrylate copolymers. Exemplary commercial grades of such copolymers include the EUDRAGITS, which are copolymers of methacrylates and acrylates. Another suitable class of amphiphilic copolymers are the 25 amphiphilic hydroxyl-functional vinyl copolymers described previously.

EXCIPIENTS AND DOSAGE FORMS

Although the key ingredients present in the compositions of the present invention are simply the dispersion and optional non-dispersion concentration-enhancing polymer(s), the inclusion of other excipients in the composition may 30 be useful. These excipients may be utilized with the drug and polymer composition in order to formulate the composition into tablets, capsules, suppositories, suspensions, powders for suspension, creams, transdermal patches, depots, and the like. The composition of drug and polymer can be added to other dosage form ingredients in essentially any manner that does not 35 substantially alter the drug. The excipients may be either physically mixed with

the dispersion and/or included within the dispersion. However, where the drug is acid-sensitive, acidic excipients should not be added to the dispersion unless neutralized.

One very useful class of excipients is surfactants. Suitable surfactants include fatty acid and alkyl sulfonates; commercial surfactants such as benzalkonium chloride (HYAMINE® 1622, available from Lonza, Inc., Fairlawn, New Jersey); dioctyl sodium sulfosuccinate, DOCUSATE SODIUM™ (available from Mallinckrodt Spec. Chem., St. Louis, Missouri); polyoxyethylene sorbitan fatty acid esters (TWEEN®, available from ICI Americas Inc., Wilmington, Delaware; LIPOSORB® P-20 available from Lipochem Inc., Patterson New Jersey; CAPMUL® POE-0 available from Abitec Corp., Janesville, Wisconsin), and natural surfactants such as sodium taurocholic acid, 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine, lecithin, and other phospholipids and mono- and diglycerides. Preferably, the surfactants are not acidic. In cases where the surfactant is acidic, such as fatty acids, the salts of the acids are preferred. Thus, sodium dodecanate is preferred relative to dodecanoic acid. Such materials can advantageously be employed to increase the rate of dissolution by facilitating wetting, thereby increasing the maximum dissolved concentration, and also to inhibit crystallization or precipitation of drug by interacting with the dissolved drug by mechanisms such as complexation, formation of inclusion complexes, formation of micelles or adsorbing to the surface of solid drug, crystalline or amorphous. These surfactants may comprise up to 5 wt% of the composition.

The addition of pH modifiers such as acids, bases, or buffers may also be beneficial, retarding the dissolution of the composition (e.g., acids such as citric acid or succinic acid when the concentration-enhancing polymer is anionic) or, alternatively, enhancing the rate of dissolution of the composition (e.g., bases such as sodium acetate or amines when the polymer is anionic). Of course, where the drug is acid-sensitive, pH modifiers should not be added to the dispersion except as discussed above in connection with including buffers or basic excipients within the dispersion.

Conventional matrix materials, complexing agents, solubilizers, fillers, disintegrating agents (disintegrants), or binders may also be added as part of the composition itself or added by granulation via wet or mechanical or other means. These materials may comprise up to 90 wt% of the composition.

Examples of matrix materials, fillers, or diluents include lactose, mannitol, xylitol, microcrystalline cellulose, calcium diphosphate, dicalcium phosphate and starch.

5 Examples of disintegrants include sodium starch glycolate, sodium alginate, carboxy methyl cellulose sodium, methyl cellulose, and croscarmellose sodium, and crosslinked forms of polyvinyl pyrrolidone such as those sold under the trade name CROSPVIDONE (available from BASF Corporation).

 Examples of binders include methyl cellulose, microcrystalline cellulose, starch, and gums such as guar gum, and tragacanth.

10 Examples of lubricants include magnesium stearate, calcium stearate, and stearic acid.

 Examples of preservatives include sulfites (an antioxidant), benzalkonium chloride, methyl paraben, propyl paraben, benzyl alcohol and sodium benzoate.

15 Examples of suspending agents or thickeners include xanthan gum, starch, guar gum, sodium alginate, carboxymethyl cellulose, sodium carboxymethyl cellulose, methyl cellulose, hydroxypropyl methyl cellulose, polyacrylic acid, silica gel, aluminum silicate, magnesium silicate, and titanium dioxide.

20 Examples of anticaking agents or fillers include silicon oxide and lactose.

 Examples of solubilizers include ethanol, propylene glycol or polyethylene glycol.

 Other conventional excipients may be employed in the compositions of this invention, including those excipients well-known in the art. Generally,
25 excipients such as pigments, lubricants, flavorants, and so forth may be used for customary purposes and in typical amounts without adversely affecting the properties of the compositions. These excipients may be utilized in order to formulate the composition into tablets, capsules, suspensions, powders for suspension, creams, transdermal patches, and the like.

30 The compositions of the present invention may be delivered by a wide variety of routes, including, but not limited to, oral, nasal, rectal, vaginal, subcutaneous, intravenous, and pulmonary. Generally, the oral route is preferred.

35 Compositions of this invention may also be used in a wide variety of dosage forms for administration of drugs. Exemplary dosage forms are

powders or granules that may be taken orally either dry or reconstituted by addition of water or other liquids to form a paste, slurry, suspension or solution; tablets; capsules; multiparticulates; and pills. Various additives may be mixed, ground, or granulated with the compositions of this invention to form a material
5 suitable for the above dosage forms.

The compositions of the present invention may be formulated in various forms such that they are delivered as a suspension of particles in a liquid vehicle. Such suspensions may be formulated as a liquid or paste at the time of manufacture, or they may be formulated as a dry powder with a liquid, typically
10 water, added at a later time but prior to oral administration. Such powders that are constituted into a suspension are often termed sachets or oral powder for constitution (OPC) formulations. Such dosage forms can be formulated and reconstituted via any known procedure. The simplest approach is to formulate the dosage form as a dry powder that is reconstituted by simply adding water
15 and agitating. Alternatively, the dosage form may be formulated as a liquid and a dry powder that are combined and agitated to form the oral suspension. In yet another embodiment, the dosage form can be formulated as two powders which are reconstituted by first adding water to one powder to form a solution to which the second powder is combined with agitation to form the suspension.

20 Generally, it is preferred that the dispersion of drug be formulated for long-term storage in the dry state as this promotes the chemical and physical stability of the drug.

A preferred additive to such formulations is additional concentration-enhancing polymer which may act as a thickener or suspending agent as well
25 as to enhance the concentration of drug in the environment of use and may also act to prevent or retard precipitation or crystallization of drug from solution. Such preferred additives are hydroxyethyl cellulose, hydroxypropyl cellulose, and hydroxypropyl methyl cellulose. In particular, the salts of carboxylic acid functional polymers such as cellulose acetate phthalate, hydroxypropyl methyl
30 cellulose acetate succinate, and carboxymethyl cellulose are useful in this regard. Such polymers may be added in their salt forms or the salt form may be formed *in situ* during reconstitution by adding a base such as trisodium phosphate and the acid form of such polymers.

In some cases, the overall dosage form or particles, granules or beads
35 that make up the dosage form may have superior performance if coated with an

enteric polymer to prevent or retard dissolution until the dosage form leaves the stomach. Exemplary enteric coating materials include hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose phthalate, cellulose acetate phthalate, cellulose acetate trimellitate, carboxylic acid-functionalized polymethacrylates, and carboxylic acid-functionalized polyacrylate.

Compositions of this invention may be administered in a controlled release dosage form. In one such dosage form, the composition of the drug and polymer is incorporated into an erodible polymeric matrix device. By an erodible matrix is meant aqueous-erodible or water-swellaable or aqueous-soluble in the sense of being either erodible or swellaable or dissolvable in pure water or requiring the presence of an acid or base to ionize the polymeric matrix sufficiently to cause erosion or dissolution. When contacted with the aqueous environment of use, the erodible polymeric matrix imbibes water and forms an aqueous-swollen gel or "matrix" that entraps the dispersion of drug and polymer. The aqueous-swollen matrix gradually erodes, swells, disintegrates or dissolves in the environment of use, thereby controlling the release of the dispersion to the environment of use. Examples of such dosage forms are disclosed more fully in commonly assigned pending U.S. Patent Application Serial No. 09/495,059 filed January 31, 2000 which claimed the benefit of priority of provisional patent application Serial No. 60/119,400 filed February 10, 1999, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the compositions of the present invention may be administered by or incorporated into a non-erodible matrix device.

Alternatively, the compositions of the invention may be delivered using a coated osmotic controlled release dosage form. This dosage form has two components: (a) the core which contains an osmotic agent and the dispersion of drug and concentration-enhancing polymer; and (b) a non-dissolving and non-eroding coating surrounding the core, the coating controlling the influx of water to the core from an aqueous environment of use so as to cause drug release by extrusion of some or all of the core to the environment of use. The osmotic agent contained in the core of this device may be an aqueous-swellaable hydrophilic polymer, osmogen, or osmagent. The coating is preferably polymeric, aqueous-permeable, and has at least one delivery port. Examples of such dosage forms are disclosed more fully in commonly assigned pending U.S. Patent Application Serial No. 09/495,061 filed January 31, 2000 which claimed

the benefit of priority of provisional Patent Application Serial No. 60/119,406 filed February 10, 1999, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the compositions may be delivered via a coated hydrogel controlled release form having at least two components: (a) a core comprising the dispersion of the present invention and a hydrogel, and (b) a coating through which the dispersion has passage when the dosage form is exposed to a use environment. Examples of such dosage forms are more fully disclosed in commonly assigned European Patent EP0378404, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the drug mixture of the invention may be delivered via a coated hydrogel controlled release dosage form having at least three components: (a) a composition containing the dispersion, (b) a water-swelling composition wherein the water-swelling composition is in a separate region within a core formed by the drug-containing composition and the water-swelling composition, and (c) a coating around the core that is water-permeable, water-insoluble, and has at least one delivery port therethrough. In use, the core imbibes water through the coating, swelling the water-swelling composition and increasing the pressure within the core, and fluidizing the dispersion-containing composition. Because the coating remains intact, the dispersion-containing composition is extruded out of the delivery port into an environment of use. Examples of such dosage forms are more fully disclosed in commonly assigned pending U.S. Patent application Serial No. 09/745,095 filed December 20, 2000, which claimed the benefit of priority of provisional patent application 60/171,968 filed December 23, 1999, the relevant disclosure of which is herein incorporated by reference.

Alternatively, the compositions may be administered as multiparticulates. Multiparticulates generally refer to dosage forms that comprise a multiplicity of particles that may range in size from about 10 μ m to about 2 mm, more typically about 100 μ m to 1 mm in diameter. Such multiparticulates may be packaged, for example, in a capsule such as a gelatin capsule or a capsule formed from an aqueous-soluble polymer such as HPMCAS, HPMC or starch or they may be dosed as a suspension or slurry in a liquid.

Such multiparticulates may be made by any known process, such as wet- and dry-granulation processes, extrusion/spheronization, roller-

compaction, melt-congealing processes or by spray-coating seed cores. For example, in wet- and dry-granulation processes, the composition of drug and concentration-enhancing polymer is prepared as described above. This composition is then granulated to form multiparticulates of the desired size.

5 Other excipients, such as a binder (e.g., microcrystalline cellulose), may be blended with the composition to aid in processing and forming the multiparticulates. In the case of wet granulation, a binder such as microcrystalline cellulose may be included in the granulation fluid to aid in forming a suitable multiparticulate.

10 In any case, the resulting particles may themselves constitute the multiparticulate dosage form or they may be coated by various film-forming materials such as enteric polymers or water-swellaable or water-soluble polymers, or they may be combined with other excipients or vehicles to aid in dosing to patients.

15 Compositions of the present invention may be used to treat any condition which is subject to treatment by administering a drug.

Other features and embodiments of the invention will become apparent from the following examples which are given for illustration of the invention rather than for limiting its intended scope.

20

EXAMPLES

Examples 1-2

This example discloses solid amorphous dispersions of an acid-sensitive, low-solubility drug and neutral polymer. For Example 1, a dispersion
25 of quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3-fluorobenzyl)-2(S),7-dihydroxy-7-methyl-octyl] amide (Drug 1) and the neutral polymer hydroxypropyl methyl cellulose (HPMC E3 Prem) was made by preparing a solution containing 0.125 wt% Drug 1 and 0.375 wt% HPMC in methanol, and spraying the solution into a drying chamber using an atomizing spray nozzle as described below. For
30 Example 2, a dispersion of Drug 1 with the neutral polymer polyvinyl pyrrolidone (PVP- Plasdone K-29/32 available from ISP Technologies Inc., Wayne, NJ) was made by preparing a solution containing 0.33 wt% Drug 1 and 1.0 wt% PVP in acetone/methanol (9/1, wt/wt), and spray-drying the solution as described below.

For Control C1, a dispersion of Drug 1 with hydroxy propyl methyl cellulose acetate succinate, LF-grade (HPMCAS-LF) (with about 14-18 wt% or 350-450 meq of succinate groups per mol of polymer) was made by preparing a solution containing 0.33 wt% Drug 1 and 1.0 wt% HPMCAS-LF in acetone, and spray-drying the solution as described below.

For Examples 1 and 2, and Control C1, the solutions were spray-dried by pumping the solution into a "mini" spray-dryer apparatus via a Cole Parmer 74900 series rate-controlling syringe pump at a rate of 1.3 mL/min. The drug/polymer solution was atomized through a Spraying Systems Co. two-fluid nozzle, Model No. SU1A using a heated stream of nitrogen (100°C). The spray solution was sprayed into an 11-cm diameter stainless steel chamber. The resulting solid amorphous dispersions containing 25 wt% Drug 1 were collected on filter paper, dried under vacuum, and stored in a dessicator. Table 1 summarizes the spray-drying variables.

Table 1

Ex.	Drug Mass (mg)	Drug No.	Polymer Mass (mg)	Polymer	Solvent Mass (g)	Solvent (wt/wt)
1	50	1	150	HPMC	40	Methanol
2	50	1	150	PVP	15	Acetone/methanol (9/1)
C1	50	1	150	HPMCAS	15	Acetone

Example 3

In this example the chemical stability of the dispersions of Examples 1 and 2 was assessed by monitoring the potency of the drug before and after exposure to increased temperatures and relative humidity (RH) in accelerated-aging studies. Dispersions of Examples 1 and 2, and Control C1, were placed in two controlled atmosphere chambers: one chamber maintained at 70°C (no RH control); the second chamber maintained at 40°C and 75% RH. Potencies of the dispersions before and after storage were determined using HPLC. A Kromasil C₄ HPLC column was used with a mobile phase of 45 vol% of 0.2 vol% H₃PO₄, and 55 vol% acetonitrile. UV detection was measured at 245 nm. Drug 1 potency was the percent of the total HPLC peak area corresponding to the theoretical amount of drug originally present in the dispersion prior to storage based on the amount of drug present in the initial solutions before spray-drying. The results are shown in Table 2 below.

Table 2

Ex.	Aqueous Soluble Polymer	Drug 1 Conc. in the Dispersion (wt%)	Pot. Before Storage (wt%)	Pot. Day 4 at 70°C (wt%)	Degree of Degradation Day 4 at 70°C (wt%)	Pot. Day 14 at 40°C/75% RH (wt%)	Degree of Degradation Day 14 at 40°C/75% RH (wt%)
1	HPMC	25	100	100	<1	100	<1
2	PVP	25	100	100	<1	99	1
C1	HPMCAS-LF	25	94	<1	>93	<1	>93

5

As the data show, the dispersions of Examples 1 and 2 formed with neutral polymers were much more chemically stable than the control dispersion C1 formed from HPMCAS. In fact, the dispersions of both Example 1 and Example 2 had a relative degree of improvement when compared with the control C1 of greater than 93 after only two weeks at 40°C/75% RH.

Example 4

In this example the dispersions of Examples 1 and 2 were tested to show that the dispersions provided concentration-enhancement of the drug in aqueous solution. For Control C2, the crystalline form of the drug alone was used without further processing. For this test, 7.2 mg of the dispersions of Examples 1 and 2, and 3.6 mg of Control C2, was added to respective microcentrifuge tubes. The tubes were placed in a 37°C temperature-controlled bath, and 1.8 mL phosphate buffered saline (PBS) at pH 6.5 and 290 mOsm/kg was added to each. The samples were quickly mixed using a vortex mixer for about 60 seconds. The samples were centrifuged at 13,000 G at 37°C for 1 minute. The resulting supernatant solutions were then sampled and diluted 1:6 (by volume) with methanol and then analyzed by high-performance liquid chromatography (HPLC). The contents of the tubes were mixed on the vortex mixer and allowed to stand undisturbed at 37°C until the next sample was

taken. Samples were collected at 4, 10, 20, 40, 90, and 1200 minutes. The results are shown in Table 3.

Table 3

5

Example	Time (min)	[Drug 1] ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
1	0	0	0
	4	863	1,700
	10	895	7,000
	20	885	15,900
	40	925	34,000
	90	907	79,800
	1200	926	1,097,100
2	0	0	0
	4	923	1,800
	10	928	7,400
	20	913	16,600
	40	947	35,200
	90	927	82,100
	1200	929	1,112,100
C2	0	0	0
	4	274	600
	10	266	2,200
	20	338	5,200
	40	289	11,500
	90	300	26,200
	1200	303	360,800

The concentrations of Drug 1 obtained in these samples were used to determine the values of the maximum drug concentration between 0 and 90 minutes ($C_{\text{max}90}$) and the area under the curve from 0 to 90 minutes (AUC_{90}).

10 The results are shown in Table 4.

Table 4

Example	$C_{\text{max}90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
1	925	79,800
2	947	82,000
C2	338	26,200

15 As can be seen from the data, the neutral polymers HPMC and PVP are both concentration-enhancing polymers in the dispersions of Examples 1 and 2. The Drug 1 and HPMC dispersion of Example 1 provided a $C_{\text{max}90}$ that was 2.7-fold,

and an AUC₉₀ that was 3.0-fold, that of the control C2, while the PVP Drug 1 dispersion of Example 2 provided a C_{max90} that was 2.8-fold, and an AUC₉₀ that was 3.1-fold, that of Control C2.

Example 5

The dispersion of Example 2 was stored for 1.5 years at room temperature (sealed), and then analyzed for Drug 1 potency by HPLC using the method described in Example 3. Results are shown in Table 5.

Table 5

Ex.	Polymer	Drug 1 Conc. in the Dispersion (wt%)	Potency Before Storage (wt%)	Potency After 1.5 yrs at room temp./sealed (wt%)
2	PVP	25	100	100

As the data show, the dispersion of Example 2 formed with PVP is chemically stable for extended duration under ambient conditions. This data is consistent with the results of accelerated storage stability tests for the dispersion of Example 2 (stored at 70°C and 40°C/75%RH) shown in Example 3 (Table 2).

In vitro dissolution tests were performed for the dispersion of Example 2 before and after 1.5 years storage, using the procedure described in Example 4, with the following exceptions: 7.2 mg of the dispersion of Example 2 was used in initial dissolution tests, and 5.4 mg of Example 2 was used in dissolution tests after 1.5 years storage. The results are reported as a percentage of the total dose dissolved, to facilitate comparison. The results are shown in Table 6.

Table 6

Time (min)	Initial Drug 1 Concentration (% dissolved)	Drug 1 Concentration After 1.5 yrs Storage (% dissolved)
0	0	0
4	91	95
10	92	90
20	90	87
40	94	88
90	92	98
1200	92	89

- 5 As can be seen from the data, the dissolution performance of the dispersion of Example 2 was maintained after 1.5 years storage, indicating that the dispersion is physically stable and Drug 1 is chemically stable for at least 1.5 years at ambient conditions.

Examples 6-9

- 10 These examples disclose additional dispersions of Drug 1 prepared with the dispersion polymers HPMC, Eudragit® E100 (available from Rohm America Inc., Piscataway, NJ), and an HPMC/Eudragit® E100 mixture. Eudragit E100 is a polymethacrylate polymer which contains basic groups. Examples 6 through 9 were prepared by spray-drying a solution of Drug 1 and polymer, as described
- 15 in Example 1. For Example 6, the solution consisted of 1.25 wt% Drug 1 and 3.75 wt% HPMC in methanol/water (9/1, wt/wt). For Example 7, the solution consisted of 0.25 wt% Drug 1 and 4.75 wt% HPMC in methanol/water (9/1, wt/wt). For Example 8, the solution consisted of 1.25 wt% Drug 1 and 3.75 wt% Eudragit® E100 in methanol/acetone (1/1, wt/wt). For Example 9, the solution
- 20 consisted of 1.25 wt% Drug 1 and 3.75 wt% HPMC/Eudragit (1/1 wt/wt) in methanol/water (9/1, wt/wt). The solution variables are summarized in Table 7.

For Control C3, the solution consisted of 1.25 wt% Drug 1 and 3.75 wt% HPMCAS-HF (HF grade) and methanol.

25

Table 7

Ex.	Drug Mass (mg)	Drug No.	Polymer Mass (mg)	Polymer	Solvent Mass (g)	Solvent (wt/wt)
6	50	1	150	HPMC	4	methanol/ water (9/1)
7	50	1	950	HPMC	30	methanol/ water (9/1)
8	125	1	375	Eudragit	10	acetone/ methanol (1/1)
9	250	1	375/375	HPMC/ Eudragit	20	methanol/ water (9/1)
C3	50	1	150	HPMCAS- HF	4	methanol

5

Example 10

In this example the chemical stability of the dispersions of Examples 6-9 was assessed. Dispersions were stored under elevated temperature and humidity conditions to increase the rate of physical changes occurring in the materials in order to simulate a longer storage interval in a typical storage environment. Drug purity was determined using HPLC as in Example 3. Results of purity analysis of the dispersions of Drug 1 after various storage intervals under various storage conditions are shown in Table 8.

10

Table 8

Ex.	Aqueous-Soluble Polymer	Drug 1 Conc. in the Dispersion (wt%)	Potency Day 5 at 40°C/75 RH (wt%)	Degree of Degradation Day 5 (wt%)
6	HPMC	25	97.3	2.7
7	HPMC	5	96.3	3.7
8	Eudragit	25	95.6	4.4
9	HPMC/Eudragit	25	97.9	2.1
C3	HPMCAS-HF	25	60	40

5 The chemical stability of the acid-sensitive drug in dispersions of Examples 6-9 containing neutral polymers were all significantly improved in comparison to the stability of dispersions of the control C3 having the acidic polymer HPMCAS-HF. The relative degree of improvement compared with the control C3 was 14.8 for Example 6, 10.8 for Example 7, 9.1 for Example 8, and 19.0 for Example 9.

Examples 11-12

10 These examples disclose dispersions of Drug 1 in a neutral polymer with added buffer. Examples 11 and 12 were prepared with Drug 1, as described in Example 1, with the following exceptions: for Example 11, the solution to be spray-dried consisted of 1.25 wt% Drug 1, 0.513 wt% sodium acetate, and 3.75 wt% HPMC in methanol/water (9/1); and for Example 12, the solution
 15 consisted of 1.25 wt% Drug 1, 0.32 wt% sodium bicarbonate, and 3.75 wt% HPMC in methanol/water (9/1). The solution variables are summarized in Table 9.

Table 9

Ex.	Drug Mass (mg)	Excipient	Excipient Mass (mg)	Polymer	Polymer Mass (mg)	Solvent (wt/wt)	Solvent Mass (g)
11	125	NaOAc	51.4	HPMC	375	MeOH/ water (9/1)	10
12	125	NaHCO ₃	32	HPMC	375	MeOH/ water (9/1)	10

5

Example 13

Chemical stability of the acid-sensitive Drug 1 in the dispersions of Examples 11 and 12 containing buffers was determined by measuring the drug purity before and after storage for Examples 11 and 12, and Control C3. Dispersions were stored under elevated temperature and humidity conditions to increase the rate of changes occurring in the materials in order to simulate a longer storage interval in a typical storage environment. Drug purity was determined using HPLC. Results of analysis of dispersions of Examples 11 and 12 after five days at 40°C/75% RH are shown in Table 10.

10

Table 10

15

Ex.	Aqueous-Soluble Polymer/Buffer	Drug 1 Conc. in the Dispersion (wt%)	Potency Day 5 at 40°C/75%RH (%)	Degree of Degradation (%)
11	HPMC/NaOAc	23	97.79	2.2
12	HPMC/NaHCO ₃	24	95.65	4.3
C3	HPMCAS-HF	25	60	40

The stability of the dispersions with neutral polymers containing buffers was significantly improved in comparison to the stability of the dispersion with HPMCAS-HF. The relative degree of improvement after five days compared with the Control C3 was 18.2 for Example 11, and 9.3 for Example 12.

20

Example 14

This example discloses dispersions of a second acid-sensitive drug, quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide (Drug 2). For Example 14, a dispersion of Drug 2 and PVP was made by first preparing a solution containing 0.030 wt% drug and 0.27 wt% PVP in methanol/acetone (1/9 wt/wt), and then spraying the solution into a drying chamber using an atomizing spray nozzle in the manner described in Example 1.

For Control C4, a dispersion of Drug 2 and HPMCAS-LF was made by preparing a solution containing 0.030 wt% Drug 2 and 0.27 wt% HPMCAS-LF in acetone, and spray-drying as described above. For Control C5, a dispersion of Drug 2 with the acidic polymer hydroxypropyl methyl cellulose phthalate (HPMCP) was made by preparing a solution containing 0.030 wt% drug and 0.27 wt% HPMCP in acetone, and spray-drying as described above. Table 11 summarizes the solution variables.

Table 11

Ex.	Drug Mass (mg)	Drug No.	Polymer Mass (mg)	Polymer	Solvent Mass (g)	Solvent (wt/wt)
14	3	2	27	PVP	10	methanol/acetone (1/9)
C4	3	2	27	HPMCAS	10	acetone
C5	3	2	27	HPMCP	10	acetone

Example 15

This example assesses the stability of the dispersion of Example 14. The dispersions of Example 14 and Controls C4 and C5 were stored in controlled atmosphere chambers for 7 days at room temperature and 0% RH. Potencies of the dispersions before and after storage were determined using HPLC as in Example 3, with the following exceptions. An HP ODS Hypersil column was used with a mobile phase of 60 vol% 0.02M KH_2PO_4 (pH 3.0) and 40 vol% acetonitrile. UV detection was measured at 248 nm. The results are shown in Table 12 below.

Table 12

Ex.	Aqueous-Soluble Polymer	Drug 2 Conc. In the Dispersion (wt%)	Potency Day 7 at room temp/0%RH (wt%)	Degree of Degradation (wt%)
14	PVP	10	84	16
C4	HPMCAS-LF	10	45	55
C5	HPMCP	10	53	47

5

As the data show, the dispersion of Example 14 comprised of the acid-sensitive drug and neutral polymer PVP had improved chemical stability relative to dispersions formed using either of the acidic polymers HPMCAS-LF or HPMCP. Example 14 had a relative degree of improvement when compared with control C4 of 3.4, and a relative degree of improvement when compared with control C5 of 2.9.

10

Example 16

This example measures the concentration-enhancing properties of the PVP dispersion of Example 14. Dissolution tests were performed, as described in Example 4 with the following exceptions: the PBS solution contained 7.3 mM sodium taurocholic acid and 1.4 mM 1-palmitoyl-2-oleyl-sn-glycero-3-phosphocholine (model fasted duodenal (MFD) solution). 1.8 mg of dispersion was added to 1.8 mL of test media. For Control C6, 0.18 mg of the crystalline form of the Drug 2 alone was used without further processing and added to 1.8 mL of MFD solution. The results are shown in Table 13 below.

15

20

Table 13

Example	Time (min)	Drug 2 ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
14	0	0	0
	4	57	100
	10	55	500
	20	55	1,000
	40	52	2,100
	90	54	4,700
C6	0	0	0
	4	2	0
	10	4	0
	20	4	100
	40	4	100
	90	7	400

The concentrations of Drug 2 obtained in these samples were used to determine the values of $C_{\max 90}$ and AUC_{90} . The results are shown in Table 14.

5 As can be seen from the data, dispersions of Example 14 provided an 8-fold improvement in $C_{\max 90}$ and a 12-fold improvement in AUC_{90} .

Table 14

Example	$C_{\max 90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
14	57	4700
C6	7	400

10

Examples 17-18

These examples disclose dispersions of a third acid-sensitive drug, quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide (Drug 3). For Example 17, a
15 dispersion of Drug 3 and HPMC was made by preparing a solution containing 0.050 wt% Drug 3 and 0.45 wt% HPMC in methanol, and spraying the solution into a drying chamber using an atomizing spray nozzle, as described in Example 1. For Example 18, a dispersion of Drug 3 with PVP was made by
20 preparing a solution containing 0.030 wt% Drug 3 and 0.27 wt% PVP in methanol/acetone (1/9 wt/wt), and spray-drying as described above.

For Control C7, a dispersion of Drug 3 and HPMCAS-LF was made by preparing a solution containing 0.030 wt% drug and 0.27 wt% HPMCAS-LF in acetone, and spray-drying as described above. For Control C8, a dispersion of
25 Drug 3 and HPMCP was made by preparing a solution containing 0.030 wt% drug and 0.27 wt% HPMCP in acetone, and spray-drying as described above. Table 15 summarizes the spray-drying variables.

30

35

Table 15

Ex.	Drug Mass (mg)	Drug No.	Polymer Mass (mg)	Polymer	Solvent Mass (g)	Solvent (wt/wt)
17	5	3	45	HPMC	10	methanol
18	3	3	27	PVP	10	acetone/ methanol (9/1)
C7	3	3	27	HPMCAS-LF	10	acetone
C8	3	3	27	HPMCP	10	acetone

5

Example 19

This example demonstrates the relative chemical stability of the dispersions of Examples 17 and 18. The dispersions of Examples 17 and 18, and Controls C7 and C8, were placed in controlled atmosphere chambers for 6 days at room temperature and 0% RH (the HPMC dispersion of Example 16 was stored for 4 days). Potencies of the dispersions before and after storage were determined using HPLC, using the method described in Example 15 above. The results are shown in Table 16 below.

10

Table 16

15

Ex.	Aqueous-Soluble Polymer	Drug 3 Conc. in the Dispersion (wt%)	Potency Day 6 at room temp/0%RH (wt%)	Degree of Degradation (wt%)
17	HPMC	10	92 (4 days)	8
18	PVP	10	95	5
C7	HPMCAS-LF	10	46	54
C8	HPMCP	10	56	44

The data show that dispersions of the acid-sensitive drug and neutral polymers were more stable than dispersions formed with the acidic polymers HPMCAS-LF and HPMCP. The dispersion of Example 18 had a relative degree of improvement of 10.8 when compared with the control C7, and 8.8 when compared with the control C8.

20

188

Example 20

This example demonstrates the concentration enhancement provided by the dispersion of Example 18. Dissolution tests were performed, as described in Example 4 with the following exceptions: samples were tested in MFD Solution, and 1.8 mg of dispersion was added to 1.8 mLs of MFD Solution. Control C9 was simply 0.18 mg of the crystalline Drug 3 alone add to 1.8 mL of MFD solution. The results are shown in Table 17 below.

Table 17

Example	Time (min)	Drug 3 ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
17	0	0	0
	4	88	200
	10	83	700
	20	80	1,500
	40	79	3,100
	90	77	7,000
C9	0	0	0
	4	4	0
	10	6	0
	20	8	100
	40	11	300
	90	13	900

The concentrations of Drug 3 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown in Table 18.

Table 18

Example	$C_{\text{max}90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
18	88	7100
C9	13	900

As can be seen from the data, dispersions of Drug 3, an acid-sensitive drug, and the neutral polymer, PVP, provided concentration-enhancement in an aqueous environment of use. The dispersion of Example 18 provided a $C_{\text{max}90}$ that was 6.8-fold, and an AUC_{90} that was 7.9-fold that of the control C9.

Examples 21-29

These examples disclose dispersions formed from various low-solubility drugs and neutral amphiphilic, hydroxyl-functional vinyl copolymers, particularly

vinyl acetate/vinyl alcohol copolymers (hereinafter "VAVAC"). For Examples 21 to 24, the dispersions were made using Drug 1. For Examples 25 to 27, the dispersions were made using 5-chloro-1H-indole-2-carboxylic acid [(1S) - benzyl-3-((3R, 4S)-dihydroxypyrrolidin-1-yl)-(2R)-hydroxy-3-oxypropyl] amide ("Drug 4"). For Examples 28 to 29, the dispersions were made using (+)-N-{3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl}-N-hydroxyurea ("Drug 5").

Several different grades of VAVAC were used. For Examples 23, 25 and 28, the VAVAC, referred to herein as "VAVAC-20%," was 80% hydrolyzed (meaning that about 20% of the vinyl repeat units were acetylated), and had an average molecular weight of 9,000-10,000 daltons. The VAVAC-20% was supplied by Aldrich Chemical Company of lot #36,062-7.

For Examples 21, 26 and 29, the VAVAC, referred to herein as "VAVAC-12%," was approximately 88% hydrolyzed (approximately 11-13% of the repeat units were acetylated), and had an average molecular weight of 13,000-23,000 daltons. VAVAC-12% was supplied by Aldrich Chemical Company, lot #36,317-0.

For Examples 22, 24 and 27, the VAVAC, referred to herein as "VAVAC-2%," was 98% hydrolyzed (approximately 2% of the repeat units were acetylated) and had an average molecular weight of 13,000-23,000 daltons. VAVAC-2% was supplied by Aldrich Chemical Company, lot #34,840-6.

To form the dispersions of Examples 21-29, solutions containing drug and polymer in a solvent were spray-dried by pumping each solution into a "mini" spray-drier apparatus as described in Example 1.

Table 19 summarizes the variables for the solutions of Examples 21-29.

190

Table 19

Ex.	Drug	Drug Mass (mg)	Drug Conc.* (%)	Polymer	Polymer Mass (mg)	Spray Solvent (wt/wt)	Solvent Mass (g)
21	1	62	25	VAVAC-12%	186	MeOH/H ₂ O (4/1)	18
22	1	692	75	VAVAC-2%	231	MeOH/H ₂ O (1/1)	50
23	1	664	50	VAVAC-20%	664	MeOH/H ₂ O (4/1)	105
24	1	522	25	VAVAC-2%	1565	MeOH/H ₂ O (4/1)	110
25	4	110	25	VAVAC-20%	330	MeOH/H ₂ O (4/1)	33.8
26	4	126	25	VAVAC-12%	376	MeOH/H ₂ O (1.8/1)	50
27	4	78	25	VAVAC-2%	233	MeOH/H ₂ O (1/1)	20.6
28	5	37	10	VAVAC-20%	330	MeOH/H ₂ O (4/1)	33.8
29	5	42	10	VAVAC-12%	376	MeOH/H ₂ O (1.8/1)	50

* Drug concentration in the final dispersion.

5

Example 30

In this example the chemical stability of the dispersions of Examples 23 and 24 was assessed. The dispersions were stored for three weeks at 40°C and 75% RH, then analyzed for Drug 1 potency by HPLC using the method described in Example 3. Results are shown in Table 20, as are the results for Control C1, a dispersion of Drug 1 and the acidic polymer HPMCAS-LF (prepared in Example 1).

15

Table 20

Ex.	Polymer	Drug 1 Conc. in the Disper- sion (wt%)	Pot. Before Storage (%)	Potency After 3 weeks @ 40°C/75 RH (%)	Degree of Degrad- ation @ 3 wks (wt%)	Relative Degree of Improve- ment
23	VAVAC- 20%	50	98	97	3	>33
24	VAVAC- 2%	25	98	96	4	>25
C1	HPMCAS- LF	25	94	<1*	>99*	-

*Values were obtained after 2 weeks.

5 As the data show, the dispersions of Examples 23 and 24 formed with VAVAC have improved chemical stability when compared with Control C1, showing a relative degree of improvement of >33 for Example 23 and >25 for Example 24.

Example 31

10 *In vitro* dissolution tests were performed for Examples 21-24, using the procedure described in Example 4, with the following exceptions: 14.4 mg of the dispersions of Example 21, 4.8 mg of the dispersion Example 22, 7.2 mg of the dispersion of Example 23, and 14.4 mg of the dispersion of Example 24 were added to respective tubes. 1.8 mL of PBS was added to each tube to achieve a theoretical total Drug 1 concentration of approximately 2000 µg/mL if all of the drug were to dissolve completely. For comparison, a Control C10 of 1.8 mg of crystalline Drug 1 was added to 1.8 mL of PBS. The results are shown in Table 21.

Table 21

Example	Time (min)	Drug 1 Concentration ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
21	0	0	0
	4	1554	3,100
	10	1860	13,300
	20	1474	30,000
	40	744	52,200
	90	685	87,900
	180	632	147,200
	1200	580	790,000
22	0	0	0
	4	809	1,600
	10	1019	7,100
	20	1162	18,000
	40	1211	41,700
	90	1273	103,835
	1200	1117	1,430,292
23	0	0	0
	4	1698	3,400
	10	1685	13,500
	20	1076	27,400
	40	608	44,200
	90	552	73,200
	1200	499	656,500
24	0	0	0
	4	1081	2,200
	10	1279	9,200
	20	1334	22,300
	40	1342	49,100
	90	662	99,200
	1200	478	731,900
C10	0	0	0
	4	241	500
	10	256	2,000
	20	299	4,700
	40	260	10,300
	90	295	25,500
	180	298	49,400
	1200	313	361,000

5 The concentrations of Drug 1 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown below in Table 22.

Table 22

Example	C _{max90} (µg/mL)	AUC ₉₀ (min*µg/mL)
21	1860	87,900
22	1273	103,800
23	1698	73,200
24	1342	99,200
C10	299	25,500

As can be seen from the data, the dispersions of Examples 21-24 provided concentration-enhancement over that of Control C10 (crystalline Drug 1 alone). Example 21 provided a C_{max90} that was 6.2-fold and an AUC₉₀ that was 3.4-fold that of Control C10; Example 22 provided a C_{max90} that was 4.3-fold, and an AUC₉₀ that was 4.1-fold that of the Control C10; Example 23 provided a C_{max90} that was 5.7-fold and an AUC₉₀ that was 2.9-fold that of Control C10; and Example 24 provided a C_{max90} that was 4.5-fold and an AUC₉₀ that was 3.9-fold that of Control C10.

Example 32

This example demonstrates a composition comprising a dispersion of an acid-sensitive drug and a neutral polymer mixed with an acidic concentration-enhancing polymer, namely HPMCAS-HF. Example 32 was prepared by mixing the dispersion of Example 21 with HPMCAS-HF. An *in vitro* dissolution test was performed using the procedure described in Example 4, with the following exceptions: 14.4 mg of the dispersion of Example 21 and 14.4 mg of HPMCAS-HF were added to 1.8 mL of PBS solution for a total theoretical Drug 1 concentration of approximately 2000 µg/mL if all of the drug were to dissolve completely. The results are shown in Table 23.

Table 23

Example	Time (min)	Drug 1 Concentration ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
32	0	0	0
	4	1365	2,700
	10	1441	11,100
	20	1399	25,300
	40	1577	55,100
	90	1541	133,100
	180	1648	276,600
	1200	1863	2,021,700

5

The concentrations of Drug 1 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown below in Table 24.

10

Table 24

Example	$C_{\text{max}90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
32	1577	133,100
21	1860	87,900

As can be seen from the data, the composition of Example 32 (i.e., the dispersion of Example 21 and HPMCAS) had an improved AUC_{90} relative to the dispersion of Example 21 alone. Results of dissolution tests for Example 21 were shown previously in Table 22, however, they are shown again in Table 24 for comparison.

15

Example 33

20

In vitro dissolution tests were performed for Examples 25-27 using the procedure described in Example 4, with the following exceptions: 14.4 mg each of the dispersions of Examples 25-27 were added to separate, respective centrifuge tubes. 1.8 mL PBS was added to each tube. For comparison, Control C11 was prepared consisting of 3.6 mg of crystalline Drug 4 added to 1.8 mL PBS. Drug 4 concentration in the samples was determined using HPLC. A Zorbax SB C_{18} HPLC column was used with a mobile phase of 35 vol.% water

25

and 65 vol.% methanol. UV detection was measured at 297 nm. The results are shown in Table 25.

5

Table 25

Example	Time (min)	[Drug 4] ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
25	0	0	0
	4	1580	3,200
	10	1581	12,600
	20	1570	28,400
	40	1594	60,000
	90	1476	136,000
	1200	248	1,093,600
26	0	0	0
	4	1957	3,900
	10	1993	15,800
	20	1948	35,500
	40	1980	74,700
	90	1959	173,200
	1200	163	1,350,900
27	0	0	0
	4	735	1,500
	10	733	5,900
	20	707	13,100
	40	691	27,100
	90	676	61,200
	1200	147	518,000
C11	0	0	0
	4	131	300
	10	114	1,000
	20	124	2,200
	40	107	4,500
	90	126	10,300
	1200	72	120,200

The concentrations of Drug 4 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown below in Table 26.

10

15

Table 26

Example	C _{max90} (µg/mL)	AUC ₉₀ (min*µg/mL)
25	1594	136,800
26	1993	173,200
27	735	61,200
C11	131	10,300

As can be seen from the data, the dispersions of Examples 25-27 provided concentration-enhancement over that of Control C11, the crystalline Drug 4 alone. The dispersion of Example 25 provided a C_{max90} that was 12-fold and an AUC₉₀ that was 13-fold that of Control C11; Example 26 provided a C_{max90} that was 15-fold and an AUC₉₀ that was 17-fold that provided by Control C11; and Example 27 provided a C_{max90} that was 6-fold, and an AUC₉₀ that was 6-fold that of Control C11.

Example 34

In vitro dissolution tests were performed for Examples 28-29, using the procedure described in Example 4, with the following exceptions: 9 mg each of the dispersions of Examples 28-29 were added to separate respective centrifuge tubes. 1.8 mL of MFD solution was then added to each tube. For comparison, Control C12 was prepared consisting of 0.36 mg of crystalline Drug 5 added to 1.8 mL of MFD solution for a total theoretical concentration of 200 µg/mL if all of the drug dissolved. Drug 5 concentration in the samples was determined using HPLC. A Waters Symmetry C₁₈ HPLC column was used with a mobile phase of 50 vol.% (0.3 vol.% glacial acetic acid, 0.2 vol.% TEA in water) and 50 vol.% acetonitrile. UV detection was measured at 260 nm. The results are shown in Table 27.

Table 27

Example	Time (min)	Drug 5 Concentration ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
28	0	0	0
	4	211	400
	10	180	1,600
	20	168	3,300
	40	155	6,600
	90	144	14,000
	1200	139	171,100
29	0	0	0
	4	210	400
	10	192	1,600
	20	186	3,500
	40	175	7,100
	90	169	15,700
	1200	155	195,500
C12	0	0	0
	4	9	0
	10	15	100
	20	21	300
	40	27	800
	90	32	2,200
	1200	42	43,300

- 5 The concentrations of Drug 5 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results are shown below in Table 28. While the amount of Drug 5 used in the Control C12 was less than the amount of Drug 5 in the test compositions of Examples 28 and 29, nevertheless a sufficient amount of Drug 5 in excess of the solubility of Drug 5
- 10 was used in Control C12 in order to provide an accurate measure of the performance of Examples 28 and 29 relative to crystalline drug.

Table 28

Example	$C_{\text{max}90}$ ($\mu\text{g/mL}$)	AUC_{90} ($\text{min} \cdot \mu\text{g/mL}$)
28	211	14,000
29	210	15,700
C12	32	2,200

15 As can be seen from the data, the dispersions of Examples 28 - 29 provided concentration-enhancement over that of crystalline Drug 5 alone. Example 28

provided a C_{max90} that was 6.6-fold and an AUC_{90} that was 6.4-fold that of Control C12; and Example 29 provided a C_{max90} that was 6.6-fold and an AUC_{90} that was 7.1-fold that provided by Control C12.

Examples 35-38

5 Examples 35 and 36 were dispersions of Drug 4 prepared with the dispersion polymers PVP and HPMC respectively. For Examples 37 and 38, dispersions of Drug 5 and the polymers PVP and HPMC were prepared. To form the dispersions of Examples 35-38, solutions containing drug and polymer in a solvent were spray-dried by pumping each solution into a "mini" spray-drier apparatus as described in Example 1. Table 29 summaries the variables for the solutions used to prepare Examples 35-38.

Table 29

Ex.	Drug	Drug Mass (mg)	Drug Conc.* (%)	Polymer	Polymer Mass (mg)	Spray Solvent (wt/wt)	Solvent Mass (g)
35	4	50	25	PVP	150	acetone	15
36	4	50	25	HPMC	150	MeOH/ water (9/1)	15
37	5	50	10	PVP	450	Acetone	80
38	5	50	10	HPMC	450	MeOH/ acetone (1/1)	80

* Drug concentration in the final dispersion

15

Example 39

In vitro dissolution tests were performed for Examples 35-38, using the procedures described in Example 4, with the following exceptions: 14.4 mg each of Example 35 and 36 were added to separate respective test tubes to which 1.8 mL of PBS solution was added, so that the total theoretical Drug 4 concentration was approximately 2000 $\mu\text{g/mL}$ if all of Drug 4 were to dissolve completely; 3.6 mg each of Examples 37 and 38 were added in separate respective test tubes to 1.8 mL of MFD solution, to provide a total theoretical Drug 5 concentration of approximately 200 $\mu\text{g/mL}$ if all of Drug 5 were to dissolve completely. Drug 4 concentration in the samples was determined as in Example 33. Drug 5 concentration in the samples was determined as in Example 34. The results are shown in Table 30.

25

Table 30

Example	Time (min)	[Drug 4] ($\mu\text{g/mL}$)	AUC ($\text{min} \cdot \mu\text{g/mL}$)
35	0	0	0
	4	771	1,500
	10	773	6,200
	20	781	13,900
	40	753	29,300
	90	703	65,700
	1200	540	755,500
36	0	0	0
	4	420	1,000
	10	422	3,400
	20	429	7,600
	40	447	16,400
	90	447	38,700
	1200	319	463,900
37	0	0	0
	4	121	200
	10	113	900
	20	111	7,100
	40	111	4,300
	90	120	10,100
	1200	106	135,500
38	0	0	0
	4	429	900
	10	347	3,200
	20	280	6,300
	40	229	11,400
	90	257	23,600
	1200	164	237,200

5 The concentrations of Drug 4 and Drug 5 obtained in these samples were used to determine the values of $C_{\text{max}90}$ and AUC_{90} . The results for dispersions containing Drug 4 are shown below in Table 31, and for dispersions containing Drug 5 in Table 32.

Table 31

Exempl e	$C_{\max 90}$	AUC_{90}
35	781	65,700
36	447	38,700
C11	131	10,300

5

Table 32

Example	$C_{\max 90}$	AUC_{90}
37	121	10,100
38	429	23,600
C12	32	2,200

As can be seen from the data, the dispersions of Examples 35 and 36 provided superior concentration-enhancement relative to the Control C11 and Examples 37 and 38 provided superior concentration-enhancement relative to the Control C12.

The terms and expressions which have been employed in the foregoing specification are used therein as terms of description and not of limitation, and there is no intention, in the use of such terms and expressions, of excluding equivalents of the features shown and described or portions thereof, it being recognized that the scope of the invention is defined and limited only by the claims which follow.

What is claimed is:

1. A pharmaceutical composition comprising a solid amorphous dispersion of an acid-sensitive drug and a neutral dispersion polymer, wherein said
5 composition provides improved chemical stability relative to a control acidic dispersion comprised of an equivalent quantity of said drug and an acidic polymer.
2. A pharmaceutical composition comprising a solid amorphous dispersion
10 comprising a low-solubility drug, a neutral dispersion polymer, and an excipient selected from the group consisting of a base and a buffer, wherein said composition provides improved chemical stability relative to a control composition, and wherein said control composition comprises an equivalent
15 quantity of a dispersion of said drug and said neutral dispersion polymer but free from said base and said buffer.
3. The composition of any one of claims 1-2 wherein said drug when
20 present in a control acidic dispersion and stored for a period of six months at 40°C and 75% relative humidity has a degree of degradation of at least 0.01%.
4. The composition of any one of claims 1-2 wherein said drug has one or
25 more functional groups selected from the group consisting of sulfonyl ureas, hydroxamic acids, hydroxy amides, carbamates, acetals, hydroxy ureas, esters, and amides.
5. The composition of any one of claims 1-2 wherein said drug is selected
30 from the group consisting of quinoxaline-2-carboxylic acid [4(R)-carbamoyl-1(S)-3-fluorobenzyl-2(S),7-dihydroxy-7-methyl-octyl]amide, quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide, quinoxaline-2-carboxylic acid [1-benzyl-4-(4,4-difluoro-1-hydroxy-cyclohexyl)-2-hydroxy-4-hydroxycarbamoyl-butyl]-amide, (+)-N-{3-[3-(4-fluorophenoxy)phenyl]-2-cyclopenten-1-yl}-N-hydroxyurea, omeprazole, etoposide, famotidine, erythromycin, quinapril, lansoprazole, and progabide.
6. The composition of any one of claims 1-2 wherein said dispersion
35 polymer is selected from the group consisting of hydroxypropyl methyl cellulose

acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, and hydroxyethyl ethyl cellulose.

- 5 7. The composition of any one of claims 1-2 wherein said dispersion polymer is selected from the group consisting of vinyl polymers and copolymers having one or more substituents selected from the group consisting of hydroxyl-containing repeat units, alkylacyloxy-containing repeat units, or cyclicamido-containing repeat units, polyvinyl alcohols that have at least a portion of their
- 10 repeat units in the unhydrolyzed form, polyvinyl alcohol polyvinyl acetate copolymers, polyethylene glycol, polyethylene glycol polypropylene glycol copolymers, polyvinyl pyrrolidone, polyethylene polyvinyl alcohol copolymers, and polyoxyethylene-polyoxypropylene block copolymers.
- 15 8. The composition of any one of claims 1-2 wherein said dispersion polymer is concentration-enhancing and said drug has a minimum solubility in aqueous solution in the absence of said dispersion polymer of less than 1 mg/mL at any pH of from about 1 to about 8.
- 20 9. The composition of any one of claims 1-2 wherein said dispersion polymer is present in an amount sufficient so that said dispersion provides to a use environment at least one of:
- (a) a maximum concentration of said drug in said use environment that is at least 1.25-fold that provided by a second control composition comprising
- 25 an equivalent quantity of said drug and free from said dispersion polymer;
- (b) an area under the concentration versus time curve for any period of at least 90 minutes between the time of introduction into said use environment and about 270 minutes following introduction to said use
- 30 environment that is at least 1.25-fold that provided by said second control composition comprising an equivalent quantity of said drug and free from said dispersion polymer; and
- (c) a relative bioavailability that is at least 1.25 relative to said second control composition comprising an equivalent quantity of said drug and
- 35 free from said dispersion polymer.

10. A pharmaceutical composition comprising a solid amorphous dispersion of a low-solubility drug and a neutral dispersion polymer, wherein said neutral dispersion polymer comprises a vinyl copolymer having hydrophilic hydroxyl-containing repeat units and hydrophobic repeat units.

5

11. The composition of any one of claims 1-2 and 10 wherein said dispersion has a pH from about 6 to about 10.

12. The composition of any one of claims 1-2 and 10 wherein said
10 composition further comprises a second polymer, said dispersion is free from at least a portion of said second polymer, and said second polymer is selected from the group consisting of hydroxypropyl methyl cellulose acetate succinate, hydroxypropyl methyl cellulose succinate, hydroxypropyl cellulose acetate succinate, hydroxyethyl methyl cellulose succinate, hydroxyethyl cellulose
15 acetate succinate, hydroxypropyl methyl cellulose phthalate, hydroxyethyl methyl cellulose acetate succinate, hydroxyethyl methyl cellulose acetate phthalate, carboxyethyl cellulose, carboxymethyl cellulose, cellulose acetate phthalate, methyl cellulose acetate phthalate, ethyl cellulose acetate phthalate, hydroxypropyl cellulose acetate phthalate, hydroxypropyl methyl cellulose
20 acetate phthalate, hydroxypropyl cellulose acetate phthalate succinate, hydroxypropyl methyl cellulose acetate succinate phthalate, hydroxypropyl methyl cellulose succinate phthalate, cellulose propionate phthalate, hydroxypropyl cellulose butyrate phthalate, cellulose acetate trimellitate, methyl cellulose acetate trimellitate, ethyl cellulose acetate trimellitate, hydroxypropyl
25 cellulose acetate trimellitate, hydroxypropyl methyl cellulose acetate trimellitate, hydroxypropyl cellulose acetate trimellitate succinate, cellulose propionate trimellitate, cellulose butyrate trimellitate, cellulose acetate terephthalate, cellulose acetate isophthalate, cellulose acetate pyridinedicarboxylate, salicylic acid cellulose acetate, hydroxypropyl salicylic acid cellulose acetate,
30 ethylbenzoic acid cellulose acetate, hydroxypropyl ethylbenzoic acid cellulose acetate, ethyl phthalic acid cellulose acetate, ethyl nicotinic acid cellulose acetate, ethyl picolinic acid cellulose acetate, carboxylic acid functionalized polymethacrylates, carboxylic acid functionalized polyacrylates, amine-functionalized polyacrylates, amine-fuctinoalized polymethacrylates, proteins,
35 and carboxylic acid functionalized starches.

13. The composition of any one of claims 1-2 and 10 wherein said composition further comprises a second polymer, said dispersion is free from at least a portion of said second polymer, and said second polymer is selected from the group consisting of hydroxypropyl methyl cellulose acetate, hydroxypropyl methyl cellulose, hydroxypropyl cellulose, methyl cellulose, hydroxyethyl methyl cellulose, hydroxyethyl cellulose acetate, hydroxyethyl ethyl cellulose, and vinyl polymers and copolymers having one or more substituents selected from the group consisting of hydroxyl-containing repeat units, alkylacyloxy-containing repeat units, or cyclicamido-containing repeat units, polyvinyl alcohols that have at least a portion of their repeat units in the unhydrolyzed form, polyvinyl alcohol polyvinyl acetate copolymers, polyethylene glycol, polyethylene glycol polypropylene glycol copolymers, polyvinyl pyrrolidone, polyethylene polyvinyl alcohol copolymers, and polyoxyethylene-polyoxypropylene block copolymers.
14. The composition of any one of claims 1, 2 and 10-13 wherein said drug in said composition has a relative degree of improvement in chemical stability of at least 1.25.
15. The composition of any of claims 1, 2 and 10-13 wherein the drug comprises [2R,4S] 4-[(3,5-bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid ethyl ester; [2R,4S] 4-[acetyl-(3,5-bis-trifluoromethyl-benzyl)-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester; or [2R, 4S] 4-[(3,5-Bis-trifluoromethyl-benzyl)-methoxycarbonyl-amino]-2-ethyl-6-trifluoromethyl-3,4-dihydro-2H-quinoline-1-carboxylic acid isopropyl ester.

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K9/14 A61K9/16		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SHAH J C ET AL: "PREFORMULATION STUDY OF ETOPOSIDE: II. INCREASED SOLUBILITY AND DISSOLUTION RATE BY SOLID-SOLID DISPERSIONS" INTERNATIONAL JOURNAL OF PHARMACEUTICS, AMSTERDAM, NL, vol. 113, no. 1, 2 January 1995 (1995-01-02), pages 103-111, XP001040972 ISSN: 0378-5173 page 104-105, paragraph entitled "2.2. Preparation of coprecipitates"; page 109-110, paragraph entitled "4. Discussion" paragraph bridging page 103 and 104 <div style="text-align: center;">--- -/--</div>	1-9, 11, 14 1, 3-9, 11-14
A		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents:		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*&* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">1 August 2002</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">08/08/2002</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Borst, M</div>

Form PCT/ISA/210 (second sheet) (July 1992)

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 64414 A (BREITENBACH JOERG ; BASF AG (DE); BINDER RUDOLF (DE); LIEPOLD BERND) 2 November 2000 (2000-11-02) page 2, line 31 - page 3, line 21; example 2, 3	1-9, 11, 14
A	page 1, line 30-37;	1, 3-9, 11-14
X	--- SARISUTA N ET AL: "PHYSICO-CHEMICAL CHARACTERIZATION OF INTERACTIONS BETWEEN ERYTHROMYCIN AND VARIOUS FILM POLYMERS" INTERNATIONAL JOURNAL OF PHARMACEUTICS, AMSTERDAM, NL, vol. 186, no. 2, 20 September 1999 (1999-09-20), pages 109-118, XP000974319 ISSN: 0378-5173 page 110-111, paragraph entitled "2.3. Preparation of the drug-polymer mixtures"	1-6, 8, 9, 11, 14
X	--- US 4 673 564 A (KAWATA HIROITSU ET AL) 16 June 1987 (1987-06-16) cited in the application column 1, line 47-51; example 1-3, 9	1-4, 6-9, 11-14
X	--- EP 1 027 886 A (PFIZER PROD INC) 16 August 2000 (2000-08-16) '0027!', '0082!', '0083!'	1, 2, 6, 7, 10
X	--- WO 97 08950 A (FUISZ TECHNOLOGIES LTD) 13 March 1997 (1997-03-13) page 6, line 4-9; examples 1-7	1-4, 7-9, 11, 14
P, X	--- WO 02 11710 A (CURATOLO WILLIAM JOHN ; FRIESEN DWAYNE THOMAS (US); GUMKOWSKI MICHA) 14 February 2002 (2002-02-14) example 14, 16	1-4, 6-9, 11, 14, 15
A	--- WO 99 14204 A (SEARLE & CO ; SIKORSKI JAMES A (US)) 25 March 1999 (1999-03-25) page 1, line 1-15; page 84, line 27-29	1-15
X	--- WO 00 17166 A (WESTER RONALD THURE ; PFIZER PROD INC (US); DENINNO MICHAEL PAUL (U) 30 March 2000 (2000-03-30) page 2, line 22 - page 6, line 2	15
A	--- WO 99 40061 A (KATH JOHN C ; BROWN MATTHEW F (US); PFIZER PROD INC (US); POSS CHRI) 12 August 1999 (1999-08-12) page 2, line 21 - page 3, line 19	5
	--- -/--	

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LEUNER C ET AL: "Improving drug solubility for oral delivery using solid dispersions" EUROPEAN JOURNAL OF PHARMACEUTICS AND BIOPHARMACEUTICS, ELSEVIER SCIENCE PUBLISHERS B.V., AMSTERDAM, NL, vol. 50, no. 1, 3 July 2000 (2000-07-03), pages 47-60, XP004257179 ISSN: 0939-6411 whole document -----	1-15

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0064414	A	02-11-2000	DE 19918434 A1	26-10-2000
			WO 0064414 A2	02-11-2000
			EP 1173152 A2	23-01-2002
US 4673564	A	16-06-1987	JP 1272484 C	11-07-1985
			JP 56133217 A	19-10-1981
			JP 59048810 B	29-11-1984
			JP 1591530 C	30-11-1990
			JP 2009007 B	28-02-1990
			JP 56049314 A	02-05-1981
			CA 1146866 A1	24-05-1983
			CH 648484 A5	29-03-1985
			DE 3024858 A1	22-01-1981
			ES 493152 D0	16-11-1981
			ES 8200557 A1	01-02-1982
			ES 499854 D0	01-06-1982
			ES 8205125 A1	16-09-1982
			FR 2460667 A1	30-01-1981
			FR 2489146 A1	05-03-1982
			GB 2053681 A , B	11-02-1981
			IT 1132169 B	25-06-1986
			SE 448342 B	16-02-1987
			SE 8004938 A	06-01-1981
			US 4343789 A	10-08-1982
			US 4404183 A	13-09-1983
EP 1027886	A	16-08-2000	BR 0000346 A	21-08-2001
			EP 1027886 A2	16-08-2000
			JP 2000229887 A	22-08-2000
WO 9708950	A	13-03-1997	AU 7106996 A	27-03-1997
			CA 2231050 A1	13-03-1997
			EP 0952770 A1	03-11-1999
			JP 11514979 T	21-12-1999
			WO 9708950 A1	13-03-1997
			US 6391338 B1	21-05-2002
WO 0211710	A	14-02-2002	AU 2914202 A	18-02-2002
			WO 0211710 A2	14-02-2002
WO 9914204	A	25-03-1999	AU 9297798 A	05-04-1999
			WO 9914204 A1	25-03-1999
WO 0017166	A	30-03-2000	AU 5439899 A	10-04-2000
			BR 9913842 A	12-06-2001
			EP 1114033 A1	11-07-2001
			WO 0017166 A1	30-03-2000
			US 6147090 A	14-11-2000
			US 6395751 B1	28-05-2002
WO 9940061	A	12-08-1999	AP 992 A	06-08-2001
			AU 1778999 A	23-08-1999
			BG 104726 A	30-04-2001
			BR 9907655 A	24-10-2000
			CA 2320388 A1	12-08-1999
			CN 1290255 T	04-04-2001
			EP 1051405 A2	15-11-2000
			HR 20000529 A1	31-08-2001

PCT/IB 02/01783

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9940061	A	WO 9940061 A2	12-08-1999
		JP 2002502839 T	29-01-2002
		NO 20003965 A	03-10-2000
		PL 342817 A1	02-07-2001
		SK 11262000 A3	18-01-2001
		TR 200002248 T2	21-12-2000
		ZA 9900873 A	04-08-2000